

Sub Chronic Inhalation Toxicity Studies of 2,2-Dichlorovinyl Dimethyl Phosphate (DDVP) in Albino Rats

¹S.O.O. Ajiboso, ¹M. Gbate, ¹O.I. Ajari and ²S.O. Adeyemo

¹Department of Science Laboratory Technology,
The Federal Polytechnic, P.M.B. 55 Bida Niger State

²Department of Biochemistry, IBB University, Lapai, Niger State Nigeria, West Africa

Abstract: 60 adult *Rattus norvegicus* (both sexes) of mean body weight of 150.0±3.5g were distributed into six groups of ten rats per group (same sex). The groups A, B, C, D, E and F were exposed to adjusted dosages of 0%, 20%, 40%, 60%, 80% and 100% of DDVP respectively by inhalation from day 1 to day 28. Biochemical analysis indicated that DDVP caused significant decrease ($p<0.05$) in GPT activity, WBC and lymphocytes levels in exposed groups while GOT activity, creatinine, haemoglobin, PCV and neutrophil levels were significantly increased. ALP activity was unchanged in groups exposed to 20%; decreased in groups exposed to 40%, 60% 80% while significant increase was observed in group exposed to 100% DDVP. Albumin was also significantly increased in groups exposed to 80% and 100% DDVP. Groups exposed to 0%, 20% and 40% DDVP showed GOT/GPT ratios <1 while >1 was the value obtained for groups exposed to 60%, 80% and 100%. Groups exposed to 0%, 20% and 40% DDVP showed weight gain while weight loss was recorded in group exposed to 100% DDVP throughout the experimental period. Investigated parameters were generally higher in group F (exposed to 100%) than other groups. Severe toxicity was observed in group exposed to 100% DDVP after 21st day of continuous exposure. Cholinergic signs observed in severely affected group that was exposed to 100% DDVP included anorexia, muscular tremors, mucous nasal discharges, increased frequency of salivation and urine staining, decreased body weight gain and diarrhoea. No adverse effects or cholinergic signs were observed in the functional observational battery and locomotors of the rats exposed to 0% to 80%; the NOEL was determined to be 20%. No mortality was recorded and none was killed because of intoxication. The leukopenia, lymphocytopenia, neutrophilia, increased haemoglobin, albumin and creatinine established in this study resulted from increased DDVP concentration and were attributed to malnutrition from low feed intake and dehydration.

Key words: DDVP • Liver • Serum • Toxicity • Dehydration

INTRODUCTION

In Nigeria, 2, 2- Dichlorovinyl Dimethyl Phosphate (DDVP or Dichlorvos) is one of the classes of insecticide referred to as organophosphate which is commonly used as insecticide in agriculture and also used to control household and stored products insects. It is effective against mushroom flies, aphids, spider mite, caterpillars, thrips and white flies in greenhouse, outdoor fruits and vegetable crops [1-4], it is traded under the brand name 'sniper' or 'ota piapia' or 'ogun efo say yanmuyanmu'. According to Tomlin [5] dichlorvos is produced as a

metabolite of the insecticide named 1,2-dibromo-2,2-dichloroethyl dimethyl phosphate. Dichlorvos may also be produced as an initial degradation product of the insecticide trichlorfon (dimethyl 2,2,2-trichloro-1-hydroxyethylphosphonate) under neutral to alkaline conditions and may in fact be responsible for the insecticidal effect of trichlorfon.

Howard [6] commented that dichlorvos is one of the more volatile organophosphates. Formulations in which dichlorvos has been available therefore include a pressurised gas formulation with CO₂ propellant, ready to use liquid formulations, plastic or naphthalene matrices

providing sustained release as well as emulsifiable concentrate and liquid concentrate formulations. Dichlorvos has been commercially manufactured and used throughout the world since 1961 [7]. 2,2-dichlorovinyl dimethyl phosphate and about 40 other organophosphate insecticides such as parathion, methyl parathion are available nationally in Nigeria.

Organophosphates are potent toxicants, the potential toxicity to humans and other animals include abdominal pain, dizziness, headaches, nausea, vomiting, as well as skin and eye problems. Additionally, many studies have indicated that pesticide exposure is associated with long-term health problems such as respiratory problems, memory disorders, dermatologic conditions, cancer, depression, neurological deficits, miscarriages and birth defects [8]. There are also benefits to the use of pesticides in the organophosphate class such as 2, 2-dichlorovinyl dimethyl phosphate (DDVP) which have saved 7 million human lives since 1945 by preventing the transmission of diseases such as malaria, bubonic plague, sleeping sickness and typhus. Dichlorvos exhibits respiratory, contact and stomach action and gives rapid knockdown; it acts by inhibiting the activity of cholinesterase enzymes which normally block nerve transmissions after they are sent, its vapour pressure is sufficiently high to give it high insecticidal activity in the vapour phase and its main importance is based on its fumigant-type action and knockdown ability [9].

In Nigeria, 2, 2-dichlorovinyl dimethyl phosphate (DDVP) is commonly produced and used as an effective and potent insecticide, its effectiveness and potency surpassed other competitive household insecticides (such as mosquito coil and flints) in Nigeria, the use and handling of DDVP as a household insecticide by Nigerians are indiscriminate. Therefore, this study aims at carrying out sub chronic toxicity studies by exposing albino rats to different concentrations of 2,2-dichlorovinyl dimethyl phosphate (DDVP) for 28 days via inhalation and also to determine the effects of exposure (sub chronic toxicity) on some biochemical parameters in exposed rats.

MATERIALS AND METHODS

Sources and Management of Experimental Animals: 60 male and female albino rats (*Rattus norvegicus*) with a mean body weight of 150.0 ± 3.5 g were obtained from the small animal holding unit of the Department of Science Laboratory Technology, the Federal Polytechnic Bida, Niger state, Nigeria-West Africa. They were allowed free access to food and water.

Collection of 2,3-Dichlorovinyl Dimethyl Phosphate (DDVP): Three bottles (100ml each) of Sniper (2, 2-dichlorovinyl dimethyl phosphate- DDVP) was purchased from a standard agro-allied store in Minna city, Niger state Nigeria.

Animal Grouping and DDVP Administration: This study was carried out after approval from the Departmental Ethical Committee on the use and care of experimental animals. The animals were handled humanely in accordance with the guidelines of European convention for the protection of vertebrate animals and other scientific purposes. 60 albino rats of both sexes were distributed into six groups (A to F) of 10 rats in each group. Group A rats were control group that were exposed to 0% DDVP solution. Groups B to F animals were exposed to different amounts of DDVP as indicated below:

- A = Exposed to 0% (0ml DDVP/100ml distilled water-v/v)
- B = Exposed to 20% (20ml DDVP/80ml distilled water-v/v)
- C = Exposed to 40% (40ml DDVP/60ml distilled water-v/v)
- D = Exposed to 60% (60ml DDVP/40ml distilled water-v/v)
- E = Exposed to 80% (80ml DDVP/20ml distilled water-v/v)
- F = Exposed to 100% (100ml DDVP/0ml distilled water-v/v)

Exposure of Animals to DDVP Solutions: 20ml of 20% DDVP solution was poured into a sealed container that was perforated by the side for easy diffusion of its odour in and out of the container and inhalation by the animals. The container was fixed in a corner in group B cage in such a way that the animals were unable to pour away the content of the container in order to prevent body contact. This procedure was repeated for other groups using the same volume (20ml). All the rats were allowed free food (rat chow) and water throughout the 28 days of experiment in the sealed wooden cages of netted doors. At the end of 28th day, rats were sacrificed by cutting their jugular veins to collect blood. Some of the blood was collected in plain bottles for the preparation of serum while some other part was collected into EDTA coated bottles and refrigerated immediately. Liver was also extracted for liver function test.

Preparation of Serum: The blood sample in a clean, dry centrifuge tube was allowed to clot and centrifuged at 224g x 10 minutes [10]. The clear supernatant (serum) was separated from the pellet, kept frozen and used within 24 hours.

Preparation of Liver Homogenate: The extracted liver was suspended in ice-cold 0.25M sucrose solution [10] buffered with Tris pH 7.4 and was cut into tiny pieces and homogenized until a fine homogenate was produced using Teflon homogenizer.

Biochemical Analyses

Transaminases and Alkaline Phosphatase Activities: The transaminases and alkaline phosphatase activities were determined in the serum and liver homogenate according to the methods described by Bergmeyer [11].

GOT

Test Principle:

L-aspartate + 2-Oxoglutarate → Oxalacetate + L-glutamate

Procedure: GOT is measured by monitoring the concentration of Oxaloacetate hydrazine formed with 2, 4-dinitrophenyl hydrazine.

GPT

Test Principle:

L-alanine + Oxoglutarate → Pyruvate + L-Glutamate (α-ketoglutarate)

Procedure: GPT is measured by monitoring the concentration Pyruvate hydrazine formed with 2, 4-dinitrophenyl hydrazine.

The enzymes activities were measured at 546 nanometer using Spectronic 20 (4001/4 model, USA).

Alkaline Phosphatase:

Test principle

p-Npp + H₂O → p-Nitrophenol + H₃PO₄

Procedure: p-Nitrophenol is kinetically measured after 3 minutes as a measure of the activity of alp. ALP activity was measured at 405 nanometer using Spectronic 20 (4001/4 model, USA).

Albumin: The method of Bromocresol Green (BCG) was used in determining serum albumin level as described by Wester [12].

Creatinine: Creatinine was determined according to Jaffe's reaction method described by Ferry *et al.* [13].

Determination of Body Weight: The method described by Ajiboso *et al.* [14] was used to determine body weight of experimental rats. Individual rat was monitored for weekly gain in body weight using digital electronic balance (Gilbertini, Italy). Gain in weight was obtained from the relationship given below:

Weekly gain in weight = Final Week Weight – Initial Week Weight

Determination of Haematological Indices: The methods described by Akpanabiatu *et al.* [15] and Uboh *et al.* [16] were used to determine the haematological indices of experimental rats.

Statistical Analysis: Data are presented as Means ± SD and analyzed using ANOVA and Duncan post hoc test and significance was determined at p < 0.05.

RESULTS

A 28-day sub chronic toxicity study of DDVP was carried out in *Rattus norvegicus*. The mean results obtained are shown in Tables 1 to 4.

Liver Enzymes: As shown in Table 1 at p<0.05, exposure of groups B and C to 20% and 40% respectively of DDVP solutions did not alter the alkaline phosphatase activity of the groups while 60% and 80% DDVP solutions caused significant decrease in exposed groups D & E respectively. However, there was a significant increase in alkaline phosphatase activity of group F that was exposed to 100% DDVP. GOT and GPT were significantly increased and decreased respectively in exposed groups B to F, increased GOT and decreased GPT activity observed in exposed groups were dose related. However, there was no significant difference in GPT activity of groups exposed to 40% and 60%; 80% and 100% DDVP solutions. The GOT/GPT ratios revealed values that were less than 1 in groups A, B and C and greater than 1 in groups D, E and F.

Serum Enzymes and Selected Bimolecules: As shown in Table 2, at p<0.05 activity of serum alkaline phosphatase was unchanged in group B that was exposed to 20% DDVP solution; serum ALP activity was significantly decreased in groups C, D and E (that were exposed to

Table 1: Result of 28-day toxicity studies of DDVP inhalation on some liver enzymes of rats

Treatment group	ALP (U/L)	GOT (U/L)	GPT (U/L)	GOT/GPT
A	165.6±3.4 ^a	8.4±0.2 ^a	34.0±0.4 ^a	0.2±0.0 ^a
B	165.6±2.3 ^a	14.1±0.7 ^b	32.2±2.0 ^b	0.4±0.0 ^a
C	165.4±5.0 ^a	19.0±1.0 ^c	20.3±1.7 ^c	0.9±0.0 ^a
D	138.2±3.0 ^b	19.0±0.3 ^c	17.0±0.1 ^d	1.1±0.0 ^{ab}
E	138.0±4.2 ^b	27.0±1.1 ^d	17.1±0.0 ^d	1.6±0.0 ^b
F	207.0±8.3 ^c	52.6±6.4 ^e	20.0±0.4 ^c	2.6±0.3 ^c

Results are mean values of triplicate determinations and expressed as Mean± SEM

At p<0.05, same letter across the column shows no significant difference

A= Exposed to 0%, B= Exposed to 20%, C= Exposed to 40%, D= Exposed to 60%, E= Exposed to 80% and F= Exposed to 100%

Table 2: Result of 28-day toxicity studies of DDVP inhalation on some selected serum enzymes and bimolecules of rats

Treatment group	ALP (U/L)	GOT (U/L)	GPT (U/L)	ALB (mg/L)	CREATININE (mg/L)
A	36.3±5.1 ^a	21.3±0.7 ^a	32.0±4.0 ^a	27.0±0.5 ^a	59.1±3.1 ^a
B	36.0±0.0 ^a	31.0±1.0 ^b	27.3±0.3 ^b	39.0±2.1 ^b	75.9±7.8 ^b
C	34.2±4.6 ^b	57.1±7.3 ^c	33.0±3.2 ^a	48.1±0.0 ^c	126.4±9.2 ^c
D	27.0±0.1 ^c	105.7±6.1 ^d	34.1±3.5 ^a	48.0±0.6 ^c	170.5±8.6 ^d
E	32.8±2.8 ^d	108.3±9.7 ^e	43.7±3.1 ^c	50.2±4.2 ^d	190.0±5.0 ^e
F	53.0±7.8 ^e	130.0±9.1 ^f	83.5±4.7 ^d	50.0±0.8 ^d	220.0±9.1 ^f

Results are mean values of triplicate determinations and expressed as Mean± SEM

At p<0.05, same letter across the column shows no significant difference

A= Exposed to 0%, B= Exposed to 20%, C= Exposed to 40%, D= Exposed to 60%, E= Exposed to 80% and F= Exposed to 100%

Table 3: Result of 28-day toxicity studies of DDVP inhalation on haematological indices of rats

Treatment group	Hb	PCV	WBC X 10 ³	Neutrophil	Lymphocyte	Monocyte	Eosinophils	Basophils	RBC Morphology
A	10.0±0.1 ^a	13.3±1.0 ^a	20.0±3.4 ^a	50.3±2.5 ^a	74.0±5.6 ^a	0.0±0.0 ^a	0.0±0.0 ^a	0.0±0.0 ^a	Normal
B	10.3±0.3 ^a	23.0±0.0 ^b	10.0±0.0 ^b	70.0±1.3 ^b	44.0±1.0 ^b	0.0±0.0 ^a	0.0±0.0 ^a	0.0±0.0 ^a	Normal
C	11.0±1.3 ^a	27.0±0.5 ^c	9.0±0.1 ^b	73.1±4.7 ^c	43.0±1.4 ^b	0.0±0.0 ^a	0.0±0.0 ^a	0.0±0.0 ^a	Normal
D	12.0±1.0 ^b	30.0±0.0 ^d	9.0±0.7 ^b	75.0±3.0 ^d	40.0±3.2 ^c	1.0±0.0 ^b	0.0±0.0 ^a	0.0±0.0 ^a	Normal
E	15.3±1.2 ^c	33.0±3.0 ^e	7.0±1.4 ^c	81.5±1.0 ^e	37.0±0.1 ^d	1.0±0.0 ^b	0.0±0.0 ^a	0.0±0.0 ^a	Slightly crenated
F	37.1±2.9 ^d	41.7±1.3 ^f	2.3±0.7 ^d	97.0±5.1 ^f	13.3±0.6 ^c	2.0±0.0 ^c	2.0±0.0 ^b	2.0±0.0 ^b	Highly crenated

Results are mean values of triplicate determinations and expressed as Mean± SEM

At p<0.05, same letter across the column shows no significant difference

A= Exposed to 0%, B= Exposed to 20%, C= Exposed to 40%, D= Exposed to 60%, E= Exposed to 80% and F= Exposed to 100%

Table 4: Result of 28-day toxicity studies of DDVP inhalation on mean body weight of rats (g)

	Day				
	0	7	14	21	28
A	151.3±9.2	156.2±6.0 (3.2%)	163.0±7.3 (4.4%)	168.4±2.0 (3.3%)	175.1±4.3 (4.0%)
B	152.7±13.0	156.3±7.1 (2.4%)	161.7±2.3 (3.5%)	165.9±4.1 (2.6%)	171.2±9.0 (3.2%)
C	150.1±11.3	155.0±1.3 (3.3%)	160.1±5.0 (3.3%)	165.7±6.3 (3.5%)	170.5±8.3 (2.9%)
D	147.1±12.1	151.1±0.7 (2.7%)	153.3±4.7 (1.5%)	150.8±7.2 (-1.6%)	147.0±6.4 (-2.5%)
E	150.3±9.1	153.5±1.1 (2.1%)	154.2±8.4 (1.1%)	147.1±9.3 (-2.7%)	142.2±5.6 (-3.3%)
F	150.9±6.7	150.2±7.4 (-0.5%)	143.5±3.1 (-4.5%)	136.0±4.0 (-5.2%)	127.1±8.3 (-6.5%)

Results are mean values of triplicate determinations and expressed as Mean± SEM

At p<0.05, same letter across the column shows no significant difference

A= Exposed to 0%, B= Exposed to 20%, C= Exposed to 40%, D= Exposed to 60%, E= Exposed to 80% and F= Exposed to 100%

40%, 60% and 80% respectively of DDVP solutions) while significant increase in serum ALP activity was observed in group F. Serum GOT activity, albumin and creatinine were significantly increased in exposed groups

B to F; serum GOT activity was 5 times and 6.5 times of control group A values in groups E and F respectively. Serum GPT activity was significantly increased in groups C to F and significantly decreased in group B. Activity of

serum GPT in group F was 3 times of control group A value and there was significant difference in ALP activity of groups D, E and F. There were no significant differences in the albumin values of group C and D; E and F, higher albumin values were recorded for groups E and F. The significant increase of creatinine was dose-related since increase in DDVP dosage (concentration) resulted into increase in creatinine level. Creatinine level of Group F was 4 times of control group A value. The liver and renal functions markers investigated were generally higher in group F than other groups.

Haematological Indices: As shown in Table 3, at $p < 0.05$ there was a significant increase in haemoglobin, PCV and neutrophil levels of exposed groups B to F. Haemoglobin was higher in group F than other groups. Haemoglobin level of group F was 3.7 times of control group A value. PCV and neutrophil levels of group F were 4 times and 2 times respectively of control group A values. WBC and lymphocytes were significantly decreased in exposed groups B to F with marked decreased of about 9 times and 7 times respectively of control group A values in group F. Monocytes were absent in groups A to C; eosinophil and basophil were also absent in groups A to E. However, monocytes levels were the same in groups D and E while 2 times amount of monocytes in groups D and E was seen in group F. Eosinophil and basophil were also detected high in group F. The results of microscopic examination of red blood cell morphology revealed normal shape of red blood cells in groups A, B, C and D (0%, 20%, 40% and 60% DDVP solutions); red blood cells of group D (exposed to 80% DDVP solution) were slightly crenated while group F (exposed to 100% DDVP) showed highly crenated red blood cells.

Body Weight: As shown in Table 4, 147.1g to 152.7g body weights range of rats were used for this study on day 0. From computed percentages on gain and loss in body weights of experimental rats on days 7, 14, 21 and 28. 2.1% to 3.3% of day 0 body weights were gained by group A to E while 0.5% body weight loss was recorded in group F on day 7, least body weight gain of 2.1% was recorded in group E. On day 14, 1.1% to 4.4% of day 7 body weights were gained by groups A to E, least body weight gain of 1.1% was observed in group D while 4.5% body weight loss was also recorded for group F. Weight gain of 2.6% to 3.5% of day 14 body weights was observed in groups A to C on day 21. Loss in body weight of 1.6% to 5.2% was also recorded for groups D to F. On day 28, weight gain of 2.9% to 4.0% of day 21 body weights was

observed in groups A to C, loss of 2.5% to 6.5% body weights was also recorded for groups D to F. General gain in body weights was recorded in groups A to C throughout the experimental period. However, change in body weights of experimental rats was dose related in which increase in DDVP concentration resulted into loss in body weights.

DISCUSSION

The liver has a variety of transaminases to synthesize and break down amino acids and to interconvert energy storage molecules. The concentrations of these in the serum (the non-cellular portion of blood) are normally low. However, if the liver is damaged, the hepatocytic cell membrane becomes more permeable and some of the enzymes leak out into the blood stream. The two transaminases commonly measured are alanine transaminase (ALT) and aspartate transaminase (AST). These levels were also called the serum glutamate-pyruvate transaminase (SGPT) and the serum glutamate-oxaloacetate transaminase (SGOT). GPT is not commonly found outside the liver; GOT too is most commonly found in the liver. In general, very high elevations of the transaminases suggest severe liver damage, such as viral hepatitis, liver injury from lack of blood flow, or injury from drugs or toxins. Most disease processes cause GPT to rise higher than GOT; GOT levels double or triple that of GPT are consistent with alcoholic liver disease [17].

The increase in activity of liver and serum GOT observed with increased DDVP concentration in this study suggests toxic and harmful effects of DDVP on the liver. This effect was more pronounced on the groups that were exposed to higher DDVP concentrations indicating that GOT activity in exposed groups was dose-dependent in this study. The increase in GOT activity of groups exposed to DDVP in this study agrees with earlier investigation of Atef [18] on effects of a class of organophosphate insecticide on physiological and histopathological investigations in rats. High levels of GPT usually indicate a damaged liver while most low GPT level indicates malnutrition as a major cause of low blood GPT levels [17]. The mild decrease in activity of liver GPT in groups D to F is due to malnutrition from withdrawal from food by groups exposed to higher concentration. According to Lum [19] malnutrition is one of the conditions that lead to low serum ALP.

The decreased liver and serum activity of ALP in this study may be attributed to malnutrition that resulted from withdrawal from food by groups D to F due to continuous

and prolonged exposure to DDVP. ALP activity was not altered in the liver of groups B and C and serum of group B. Impairment in alkaline phosphatase of rats exposed to other classes of insecticides such as DDT, malathion, phosalone and elsan as reported by Saigal *et al.* [20] was also observed in this study on altered activity of both liver and serum ALP of groups exposed to 20% and 40% DDVP solutions. Albumin is the main protein of plasma; its main function is to regulate the colloidal osmotic pressure of blood. The increased albumin observed in the exposed groups may be attributed to dehydration [17] from the withdrawal from food by exposed groups. The kidney maintains the blood creatinine in a normal range, creatinine has been found to be a fairly reliable indicator of kidney function and elevated creatinine level suggests impaired kidney function or kidney disease [21-23].

Increase in creatinine levels of exposed groups shows the toxic and idiosyncratic properties of DDVP, the increase in creatinine levels of groups exposed to DDVP in this study agrees with the result obtained by Atef [18] on effects of malathion on creatinine levels of rats. According to Gimson [17] GOT/GPT ratio is useful in determining extent of damage to liver cell integrity and also for differentiating between causes of liver damage (hepatotoxicity). The GOT/GPT ratio < 1 of groups B and C indicates the non-toxic effects of DDVP solutions on the exposed groups B and C while the GOT/GPT ratio value < 1 of unexposed group A also indicates normal liver cell integrity. Groups D, E and F GOT/GPT ratio > 1 suggests damaged liver, this may be attributed to increased DDVP concentration in groups D, E and undiluted DDVP solutions.

Blood analysis is crucial in many fields of toxicology research and in environmental monitoring as possible indicator of physiological or pathological changes in diseases investigation. In warm-blooded animals, changes in the blood parameters occur because of injuries or infections of some tissues or organs, leading to the dysfunction or injuries of organs or tissues [24]. According to Gu [25] dehydration is an important factor that results in increase in hemoglobin and PVC levels while seizure leads to neutrophilia (increased neutrophil). Leukopenia (decreased WBC) and lymphocytopenia (reduced lymphocytes) recorded in this present study may be attributed to destruction of WBC and lymphocytes cells [25] due to continuous and prolonged exposure to DDVP. Absence of monocytes prevent the occurrence of atherosclerosis [25] while presence of monocytes in groups exposed to increase concentration of DDVP suggests alteration in lipid metabolism

particularly in group F that was exposed to 100% DDVP. Increased eosinophilia and basophilia suggest tissue inflammation in group exposed to 100% DDVP suggest tissue inflammation in the group.

Significant increase in body weights of the unexposed and exposed groups to 20% and 40% DDVP solutions throughout the experimental period may be attributed to an increase in the feed intake of the groups. The decrease in body weights observed in groups exposed to 60% and 80% DDVP on days 21 to 28 and the weekly decrease in body weights of the groups exposed to undiluted DDVP indicate decrease in feed intake, this observation confirms the report of El-Hilaly [26] on changes in body weight and feed intake as an indicator of the adverse effects of drugs and chemicals [27-28]. Volatilisation of DDVP solutions established in this study from the observed toxic effects on exposed groups via inhalation was in accordance with the earlier reports of Mensink *et al.* [29] on the volatility of DDVP (dichlorvos) from dry surfaces, making dichlorvos one of the more volatile organophosphates and also on Gu [22] report of higher mobility of highly water soluble metabolites of dichlorvos than the parent substance.

Spray Drift Task Force [30] also reported the potential of Dichlorvos to reach non-target areas directly via sprayed dichlorvos solutions as vapours or indirectly through movement of directly released vapours or dichlorvos volatilising from sprayed surfaces in spray drift. However, the methods of use employed in this study involved neither direct spray nor release from sprayed surfaces but release from non-contact perforated containers. Thus, an indication that DDVP has mobility potential whether sprayed or not (in use and not in use) and this mobility is dose-related (concentration dependent). Severe toxicity was observed after the 21st day of continuous exposure of group F rats to 100% DDVP. Cholinergic signs observed in group F rats included anorexia, muscular tremors, mucous nasal discharges, increased frequency of salivation and urine staining, decreased body weight gain and diarrhoea. No adverse effects or cholinergic signs were observed in the functional observational battery and locomotor of the exposed rats to 20% to 80% DDVP, the NOEL was determined to be 20% diluted DDVP.

CONCLUSION

The 28-day subchronic toxicity studies of continuous exposure of rats by inhalation to different concentrations of DDVP showed dose-related toxic effects on the

investigated parameters. DDVP showed high mobility potential as a highly volatile compound in both diluted state and parent concentrate whether sprayed or in containing container. 100% DDVP showed severe toxicity while no observed effect level (NOEL) in this study was 20% (20ml DDVP in 80ml water, v/v).

Recommendations: The following recommendations are hereby made:

- The use of DDVP in diluted form (by mixing 20ml DDVP with 80ml water) as household insecticide.
- When not in use it should always be kept in tight, sealed and non perforated container.
- It should be kept in a safe place that is out of reach of children.
- It should be kept in a cool and not hot place to avoid explosion being a volatile compound.
- It should be applied in corners of room away from food stuffs.
- Avoid usage in kitchen and contact with drinking water tank and other food items.
- Hand gloves should be worn during preparation of diluted solutions.
- Hands should be washed after DDVP application.

REFERENCES

1. Murphy, S., 1986. Toxic Effects of Pesticides. In: Casarett and Doull's Toxicology, the Basic Science of Poisons, 3rd Edition. MacMillan Publishing Co. Inc. New York, pp: 519-581.
2. Kabiri, M. and B. Amiri-Besheli, 2012. Toxicity of Palizin©, Mospilan© and Consult© on *Agonoscena pistaciae* Burckhardt and Lauterer (Hemiptera: Psyllidae), *Oenopia conglobata* L. (Coleoptera: Coccinellidae) and *Psyllaephagus pistaciae* Ferrière (Hymenoptera: Encyrtidae). Academic Journal of Entomology, 5(2): 99-107.
3. Ahmed, S.A., L.C. Dutta and M.C. Sarmah, 2012. Bio-Efficacy of some Insecticides against Leaf Eating Caterpillar *Cricula trifenestrata* Helfer (Lepidoptera: Saturniidae) Infesting Som *Persea bombycina* Kost. Plantation. Academic Journal of Entomology, 5(2): 94-98.
4. Saur, G., 2005. Efficacy of kaolin particle film and selected synthetic insecticides against pistachio psyllid *Agonoscena targionii* (Homoptera: Psyllidae) infestation. Crop Protection, 24(8): 711-717.
5. Tomlin, C.D.S., 1997. The pesticide manual: a world compendium. Eleventh edition, British Crop Protection Council, Surrey UK, pp: 372-374.
6. Howard, P., 1991. Handbook of Environmental fate and Exposure data for organic chemicals. Volume III, *Pesticides*, Lewis Publishers Chelsea, Michigan, USA, pp: 251-258.
7. WHO, Pesticide residues in food - 1993. Toxicology evaluations. Joint meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Expert Group on Pesticide Residues. Geneva, 20-29 September, 1993.
8. Gupta, R.C., G.T. Patterson and W.D. Dettbarn, 1991. Comparison of cholinergic and neuromuscular toxicity following acute exposure to sarin and VX in Rat. *Fundamental and Applied Toxicology*, 16: 449-458.
9. Brimijoin, Enzymology and biology of cholinesterases. In the Proceedings of the 1992 U.S. EPA Workshop on Cholinesterase Methodology, pp: 4-5.
10. Ajiboso, S.O.O., 2000. Effects of *Methyl angolensate* in Rats: An in vivo study. In the Proceedings of the 2004 1st Biennial Conference of the Nigerian Society for Experimental Biology, pp: 52.
11. Bergmeyer, U.H., 1986. Methods of enzymatic analysis 3rd Edn, Verlag Chemie Press Germany. Vol II and IV. pp: 864, 1786.
12. Webster, D., 1974. A study of the interaction of BCG with isolated serum globulin fractions. *Clinical Chim Acta*, 53: 109-15.
13. Ferry, N., A. Caillette, J. Goudable, C. Denicola and N. Pozet, 1996. Creatinine determination in peritoneal dialysis: what method should be used? *Nephrology Dialysis Transplant*, 11: 2285-2287.
14. Ajiboso, S.O.O., M. Gbate, O.I. Adejumo and S.O. Adeyemo, 2007. A study on the performance of grain residues rations in ANAK 2000 chicks. *Scientific Research and Essay*, 2(8): 353-357.
15. Akpanabiatu, M.I., O. Otitoloju, E.E. Effiong, J.I. Ndem, A.F. Uwah and U.F. Ufot, 2012. Vitamin E supplementation *Rauwolfia vomitoria* root bark extract improves haematological indices. *North American Journal of Medical Science*, 4(2): 86-89.
16. Uboh, E.F., E.I. Okon and B.M. Ekong, 2010. Effect of aqueous extract of *Psidium guajava* leaves on liver enzymes, histological integrity and haematological indices in rats. *Gastroenterology Research*, 3(1): 32-38.

17. Gimson, A.E., 1996. Fulminant and late onset hepatic failure. *British Journal of Anaesthesia*, 77(1): 90-8.
18. Atef, M.A., 2010. Physiological and Histopathological Investigations on the Effects of α -Lipoic Acid in Rats Exposed to Malathion. *Journal of Biomedicine and Biotechnology*. Article ID 203503, pp: 1-8.
19. Lum, G., 1995. significance of low serum alkaline phosphatase activity in a predominantly adult male population. *Clinical Chemistry*, 41(4): 515-8.
20. Saigal, S., V. K. Bhatnagar and A. N. Malviya, 1982. Effect of selected pesticides on alkaline and acid phosphatase in the rat. *Toxicology Letter*, 12(3): 177-80.
21. Al-Jassabi, S., A. Al-Omari, Ali Saad and Mohd Sofian Azirun, 2011. Tamsulosin- induced hepatotoxicity and nephrotoxicity and its prevention by potato peel extract. *American-Eurasian Journal of Toxicological Sciences*, 3(2): 52-58.
22. Chauhan, P.K., I.P. Pandey and Vinod Kumar Dhatwalia, 2010. Evaluation of the Anti-diabetic Effect of Ethanolic and Methanolic Extracts of *Centella asiatica* Leaves Extract on Alloxan Induced Diabetic Rats. *Advances in Biological Research*, 4(1): 27-30.
23. Olojo, E.A.A., K.B. Olurin, G. Mbaka and A.D. Oluwemimo, 2005. Histopathology of the gill and liver tissues of the African catfish *Clarias gariepinus* exposed to lead. *African Journal of Biotechnology*, 4: 117-122.
24. Folmar, 1993. Acute Effects Of Diazinon On Blood Paramters In The African Catfish (*Clarias Gariepinus*). *The Internet Journal of Hematology*, 5(2): 58-65.
25. Gu, L., 1998. Absence of monocytes reduces atherosclerosis. *Molecular. Cell*, 2(2): 275-81.
26. El-Hilaly, J., Z.H. Israili and B. Lyoussi 2004. Acute and chronic toxicological studies of *Ajuga iva* in experimental animals. *Journal of Ethnopharmacology*, 91: 43-50.
27. Shobhit, K., K.G. Satish and P.K. Sharma, 2012. A Review on Recent Trends in Oral Drug Delivery-Fast Dissolving Formulation Technology. *Advances in Biological Research*, 6(1): 06-13.
28. Hamed, N., E. Gharedaashi, M. Hosseinzadeh and M. R. Imanpour 2012. Determination of LC50 of Lead Nitrate and Copper Sulphate in Common Carp (*Ciprinus carpio*). *American-Eurasian Journal of Toxicological Sciences*, 4(2): 60-63.
29. Mensink, B.J.W.G., M. Montforts, L.H.T. Wijkhuizen-Maslankiewicz and J.B.H.J. Linders, 1995. Manual for summarising and evaluating the environmental aspects of pesticides. The Netherlands National Institute of Public Health and the Environment. Report No. 679101022.
30. Spray Drift Task Force, 1997. AgDRIFT™ spray model: a tiered approach for the assessment of spray drifts of pesticides, Version 1.05, November 1997. Spray Drift Task Force, c/o Stewart Agricultural Research Services Inc., Macon, Missouri, USA. *Lamellidens marginalis*. *Journal of Environmental Biology*, 7: 101.