

A Comparative Study on Composition Analysis of Plants Consumed in India and Iran

Ali Aberoumand

Department of Food Science and Technology,
Behbahan Branch, Islamic Azad University, Behbahan, Iran

Abstract: The proximate composition and mineral constituents of *Asparagus officinalis* stem and *Alocacia indica* stem were evaluated in order to standard methods. *Asparagus officinalis* stem contained a ashes: 10.70%, crude protein: 32.69%, crude lipid: 3.44%, crude fiber: 18.50% and carbohydrates: 34.67%. The stem also have high energy value (384.27kcal/100g) dry. *Alocacia indica* stem contained a ashes: 9.1%, crude protein: 5.44%, crude lipid: 3.25%, crude fiber: 22.9% and carbohydrates: 59.31%. The stem also have high energy value (288.25kcal/100g) dry weight. Comparing proximate and minerals contents of the stem and the fruit, the results indicated that *Asparagus officinalis* stem could be a good supplement for some nutrients such as protein, potassium and carbohydrates. The stem could be promoted as a carbohydrate supplement for cereal-based diets in poor rural communities, while *Alocacia indica* stem could be a good supplement for some nutrients such as Fibre, Potassium, Zinc, lipid and Carbohydrates. *A. officinalis* is suitable for high-temperature food processes.

Key words: *Asparagus officinalis* DC. • *Alocacia indica* stem • Micronutrients • Proximate and Mineral composition

INTRODUCTION

In developing nations, numerous types of edible wild plants are exploited as sources of food hence provide an adequate level of nutrition to the inhabitants. Recent studies on agro pastoral societies in Africa indicate that these, plant resources play a significant role in nutrition; food security and income generation [1].

Furthermore, FAO report, at least one billion people are thought to use wild foods in their diet [2]. In Ghana along, the leaves of over 300 species of wild plants and fruits are consumed. In Swaziland, wild plants provide a greater share of the diet than domesticated cultivars. In India, Malaysia and Thailand, about 150 wild plants species have been identified as sources of emergency food [3]. Similarly, in South Africa about 1400 edible plant species are used. In Sahel region of Africa, over 200 wild foods were identified to be used by the rural communities [4]. In most of these reports, it was emphasized that nutritionally, these unconventional plants foods could be comparable to or even sometimes superior to the introduced cultivars (Edmonds and [5]. It is, therefore, worthwhile to note that the incorporation of edible wild and semi-cultivated plant resources could be beneficial to nutritionally marginal populations or to certain

vulnerable groups within populations, especially in developing countries where poverty and climatic changes are causing havoc to the rural populace. In this context, analyses were carried out to evaluate the nutritional content of *Asparagus officinalis* stem and *Alocacia indica* stem with hope that it would be incorporated into the food basket of the country [3, 6-8] Aim of analysis of nutrients in the plant foods is preliminary assessment of nutritional value of the plant-based diet.

Plant Material: *A. officinalis* stem and *A. indica* stem used as experimental material were collected from farm and Agricultural lands (garden) in around Behbahan, South Iran, in October 2007. The collected plant material was placed in a polyethylene bag to prevent loss of moisture during transportation to the laboratory.

Preparation of the Plant Materials for Chemical Analyses: *A. officinalis* stem and *A. indica* stem were washed with distilled water and dried at room temperature to remove residual moisture, then placed in paper envelope and oven-dried at 55°C for 24 hours [9, 10]. The dried stem were ground into powder using pestle and mortar and sieved through 20-mesh sieve. The stem powder was used for the nutrients analyzes.

Plant Foods Chemical Analysis: The methods recommended by the Association of Official Analytical Chemists were used to determine ash (#942.05), crude lipid (#920.39), crude fibre (#962.09) and nitrogen content (#984.13) [11].

Determination of Crude Lipid and Crude Fibre Content: Two grams of dried stem or fruits were weighed in a porous thimble of a Soxhlet apparatus, with its mouthed cotton wool plugged. The thimble was placed in an extraction chamber which was suspended above a pre-weighed receiving flask containing petroleum ether (b.p. 40-60°C). The flask was heated on a heating mantle for eight hours to extract the crude lipid. After the extraction, the thimble was removed from the Soxhlet apparatus and the solvent distilled off. The flask containing the crude lipid was heated in the oven at 100°C for 30 minutes to evaporate the solvent, then cooled in a dessicator and reweighed. The difference in weight was expressed as percentage crude lipid content.

Crude fibre was estimated by acid-base digestion with 1.25% H₂SO₄ (prepared by diluting 7.2 ml of 94% conc. acid of specific gravity 1.835g ml⁻¹ per 1000 ml distilled water) and 1.25% NaOH (12.5 g per 1000 ml distilled water) solutions. The residue after crude lipid extraction was put into a 600 ml beaker and 200 ml of boiling 1.25% H₂SO₄ added. The contents were boiled for 30 minutes, cooled, filtered through a filter paper and the residue washed three times with 50 ml aliquots of boiling water. The washed residue was returned to the original beaker and further digested by boiling in 200 ml of 1.25% NaOH for 30 minutes. The digest was filtered to obtain the residue. This was washed three times with 50 ml aliquots of boiling water and finally with 25 ml ethanol.

The washed residue was dried in an oven at 130°C to constant weight and cooled in a dessicator. The residue was scraped into a pre-weighed porcelain crucible, weighed, ashed at 550°C for two hours, cooled in a dessicator and reweighed. Crude fibre content was expressed as percentage loss in weight on ignition, [11, 12].

Determination of Nitrogen Content and Estimation of Crude Protein: Macro-Kjeldahl method was used to determine the nitrogen content of the stem 2g of dried stem were digested in a 100 ml Kjeldahl digestion flask by boiling with 10 ml of concentrated tetraoxosulphate (VI) acid and a Kjeldahl digestion tablet (a catalyst) until the mixture was clear. The digest was filtered into a 100 ml

volumetric flask and the solution made up to 100 ml with distilled water. Ammonia in the digest was steam distilled from 10 ml of the digest to which had been added 20 ml of 45% sodium hydroxide solution. The ammonia liberated was collected in 50 ml of 20% boric acid solution containing a mixed indicator. Ammonia was estimated by titrating with standard 0.01 mol L⁻¹ HCl solution. Blank determination was carried out in a similar manner. Crude protein was estimated by multiplying the value obtained for percentage nitrogen content by a factor of 6.25, [11].

Estimation of Carbohydrates and Energy Values: Available carbohydrate was estimated by difference, by subtracting the total sum of percent crude protein, crude lipid, crude fibre and ash from 100% DW of the plant the plant calorific value (in kJ) was estimated by multiplying the percentages of crude protein, crude lipid and carbohydrate by the factors 16.7, 37.7 and 16.7 respectively [11].

Mineral Analysis: The mineral elements Na, K, Ca, Fe and Zn were determined on 0.3g plant powder by the methods of Funtua. using Energy Dispersive X-ray Fluorescence (EDXRF) transmission emission spectrometer carrying an annular 25 mCi 109Cd isotopic excitation source that emits Ag-K X-ray (22.1 keV) and a Mo X-ray tube (50KV, 5mA) with thick foil of pure Mo used as target material for absorption correction. The system had a Canberra Si (Li) detector with a resolution of 170eV at 5.9keV line and was coupled to a computer controlled ADCCard (Trump 8K).

Measurements were carried out in duplicate. Na was analyzed after wet digestion of one gramme of the stem powder with nitric/perchloric/sulphuric acid (9:2:1 v/v/v) mixture. Sodium was analyzed with a Corning 400 flame photometer [11].

RESULTS AND DISCUSSION

Proximate Analysis: The results of proximate composition of *A. officinalis* stem and *A. indica* stem are shown in Tables 1 and 2. The ash content, which is an index of mineral contents was less than to the values reported for other edible leaves such as *Momordica balsamina* leaves ($18.00 \pm 1.27\%$ DW) where as for *Momordica dioica* fruits the value of 9.1% DW was less than the reported values [1, 3, 10]. It is apparent that *A. officinalis* stem are a good source of Potassium while *Alocacia indica* stem are a good source of Potassium and

Table 1: Proximate composition of *Asparagus officinalis* stem

Parameters	Concentration (% DW) *
Ash	10.70± 0.80
Crude protein	32.69± 0.27
Crude lipid	3.44± 0.50
Crude fibre	18.50± 0.35
Carbohydrates	34.67±0.68
Calorific value (kcal/100g)	384.27±5.31

* The data are mean values± deviation (SD) of three replicates.

* Values expressed as % wet weight.

Table 2: Proximate composition of *Alocacia indica* S. stem

Parameters	Concentration (% DW) *
Ash	7.3± 0.80*
Crude protein	5.7± 0.27*
Crude lipid	3.29± 0.50*
Crude fibre	11.05± 0.35*
Carbohydrates	72.66±0.68*
Calorific value (kcal/100g)	343.05±5.31

* The data are mean values± deviation (SD) of three replicates.

* Values except Calorific value expressed as % DW

Table 3: Mineral composition of *Asparagus officinalis* stem

Recommended Dietary Allowances (mg/day) †					
Mineral	Available Quantity in mg/100gDW*	Children 7-10 Years	Adult male	Adult female	Pregnant and Lactating mother
Potassium	10.94±0.02	800	800	800	1200
Calcium	0.67±0.15	1600	2000	2000	2000
Sodium	1.840±0.08	400	500	400	500
Iron	0.19±0.01	10	10	15	13
Zinc	2.60±0.07	10	15	12	19

* The data are mean values± deviation (SD) of three replicates. † Sources: Thangadari *et al.* (2001)

Table 4: Mineral composition of *Alocacia indica* S. stem

Recommended Dietary Allowances (mg/day) **					
Minerals	Available quantity in mg/100gDW*	Children 7-10years	Adult male	Adult female	Pregnant and Lactating Mothers
Calcium	0.88±0.02	800	800	800	1200
Potassium	3.40±0.15	1600	2000	2000	2000
Sodium	4.40±0.08	400	500	400	500
Iron	0.48±0.01	10	10	15	13
Zinc	1.21±0.07	10	15	12	19

Zinc. The crude protein content of both edible plants was more than what is reported for some lesser known wild leafy vegetables such as *Momordica balsamina* ($11.29 \pm 0.07\%$) , *Moringa oleifera* (20.72%) , *Lesianthera africana* leaves (13.10-14.90%) and *Leptadenia hastate* (19.10%) [13, 14]. Plant food that provide more than 12% of their calorific value from protein are a good source of protein. In that context, *A. officinalis* stem (32.69%) and *A.indica* stem (19.38%) are good sources of protein. The crude lipid contents of both plant foods were less than the range (8.3-27.0% DW) reported for some vegetables consumed in Nigeria and Republic of Nigerian, [15].

The estimated carbohydrate contents (34.67%) in *A. officinalis* stem and in (47.92%) in *A. indica* stem were stand to be higher than that for *Senna obtusifolia* leaves (20%) and *Amaranthus incurvatus* leaves (23.7%). On the other hand, *A. officinalis* stem and *A. indica* stem contain comparable amount of carbohydrate for *Momordica balsamina* ($39.05 \pm 2.01\%$) . The crude fibre content in both plant foods were more than the reported values (8.50-20.90%) for some Nigeria vegetables, Isong and Idiong [15]. One discussed drawback to the use of vegetables in human nutrition is their high fibre content, which may cause intestinal irritation and a decrease of

nutrient bioavailability. The fibre RDA values for children, adults, pregnant and breast-feeding mothers are 19-25%, 21-38%, 28% and 29% respectively. The calorific value of *A. officinalis* stem was estimated to be 384.27 kcal/100g (DW) and 311.5 kcal/100g (DW) respectively was estimated to be 311.5 kcal/100g (DW), which is an indication that it could be an important source of dietary calorie. High calorific content of the stem could be attributed to high carbohydrates and protein contents.

Mineral Content: Table 3 and 4 show the results of the mineral concentrations of *A. officinalis* stem and *A. officinalis* stem. Nutritional significant of elements is compared with the standard recommended dietary allowance. When compared with standard values as showed in Table 3 and 4, *A. officinalis* stem and *A. officinalis* stem less than adequate level of K, Fe, Zn, Ca and Na, but the plant stem could be good source of K.

Concluding Remarks: The results of the nutritional analysis shown that *A. officinalis* stem is good sources of plant potassium, protein and carbohydrates where as *A. officinalis* stem is good sources of plant fibre, potassium, zinc, lipid, protein and carbohydrates. The results suggests that the plant fruits if consumed in sufficient amount could contribute greatly towards meeting human nutritional requirement for normal growth and adequate protection against diseases arising from malnutrition. From the result, *A. officinalis* stem and *A. officinalis* stem are recommend for continues used for nutritional purposes, considering to the amount and diversity of nutrients it contains. Chemical analysis alone however, should not be the exclusive criteria for judging the nutritional significance of a plant parts. Thus, it becomes necessary to consider other aspects such as presence antinutritional/ toxicological factors and biological evaluation of nutrient content, [16].

In conclusion, the edible plant suitable for different technological processes:

- *A. officinalis* plant is suitable for high-temperature food processes, because it has very low carbohydrate concentrations-thereby reducing the possibility of Maillard reaction and then acryl amide formation [17].
- *A. officinalis* should not be used in high-temperature processes and should be cooked without the peel. We suggest its use for domestic purposes and home cooking.

Abbreviations Used: AOAC, Association of Official Analytical Chemists; FAO, Food and Agricultural Organization; RDA, recommended dietary allowances; DW, dried weight.

ACKNOWLEDGEMENTS

The authors are grateful to Head Department of Food Science Technology of Ramin Agricultural University of Iran. for providing necessary laboratory facilities and for encouragement.

REFERENCES

1. Asibey-Berko, E. and F. Tayie, 1999. The antibacterial properties of some plants found in Hawaii, Ghana J. Sci., 39: 91-92.
2. Burlingame, B., 2000. Comparison of total lipids, fatty acids, sugars and nonvolatile organic acids in nuts from *Castanea* species. J. Food Composition and Analysis, 13: 99-100.
3. Faruq, U., A. Sani and L. Hassan, 2002. Composition and distribution of deadly nightshade. Niger J. Basic Application Sci., 11: 157-164.
4. Sena, L., D. VanderJagt, C. Rivera, A. Tsin, I. Muhammadu, O. Mahamadou, M. Milson, A. Pastosyn and R. Glew, 1998. Nutritional profile of some edible plants from Mexico, Plant Foods for Human Nutrition, 52: 17-30.
5. Edmonds, J. and J. Chweya, 1995. Black nightshades, *Solanum nigrum L.* and related species. Promoting the conservation and use of underutilized and neglected crops. Taylor and Francis, London, pp: 221-234.
6. Funtua, I., 2004. Minerals in foods: Dietary sources, chemical forms, interactions, bioavailability, Instrumentation Sci. Technol., 32: 529-536.
7. Funtua, I. and J. Trace, 1999. Quantitative variability in *Pisum* seed globulins: its assessment and significance. Plant Foods for Human Nutrition, 17: 293-297.
8. Ifon, E. and O. Bassir, 1980. Determination of carbohydrates in foods. II-unavailable carbohydrates. Food Chemistry, 5: 231-235.
9. Abuye, C., K. Urga, H. Knapp, D. Selmar, A. Omwega, J. Imungi and P.A. Winterhalter, 2003. survey of wild, green, leafy vegetables and their potential in combating micronutrient deficiencies in rural populations, East African Medicine J., 80: 247-252.

10. Aletor, V. and O. Adeogun, 1995. Chemical analysis of the fruit of *Vitex doniana* (Verbenaceae) . Food Chemistry, 53: 375-379.
11. AOAC, 1990. Official methods of analysis, 14th edition, Association of Official Analytical Chemists, Washington DC. Arlington, Virginia, USA.
12. Nesamvuni, C., N. Steyn and M. Potgieter, 2001. nutrients analysis of selected western African foods, South African J. Sci., 97: 51-54.
13. Pearson, D., 1999. Nutrient and chemical composition of 13 wild plant foods of Niger. Ghana J. Sci., 39: 91-92.
14. Plessi, M., D. Bertelli, A. Phonzani, M. Simonetti, A. Neri and P. Damiani, 1999. Role of indigenous leafy vegetables in combating hunger and malnutrition, J. Food Composition and Analytical, 12: 91-96.
15. Isong, E. and U. Idiong, 1997. Nutrient content of the edible leaves of seven wild plants from Nigerian. Plant Foods for Human Nutrition, 51: 79-84.
16. Aberoumand, A. And S.S. Deokule, 2008. Comparison of compounds of some edible plants of Iran and India, Pakistan J. Nutrition, 7(4): 582-585.
17. Mottram, D.S., B.L. Wedzicha and A.T. Dodson, 2002. Acrylamide is formed in the Maillard reaction. Nature, 419: 449-450.