

Factor Analysis for Performance and Other Characteristics in Durum Wheat under Drought Stress and Without Stress

¹Roza Gholamin, ²Mohammad Zaefizadeh and ¹Majid Khayatnezhad

¹Young Researchers Club, Islamic Azad University, Ardabil Branch, Ardabil, Iran

²Islamic Azad University, Ardabil Branch, Ardabil, Iran

Abstract: Since the correlation coefficients may complete information on the relationship between different traits and not to provide benefits according to several multivariate statistical analysis to understand the deep structure of data, factor analysis can be used. In order to assess this potential performance grain durum wheat genotypes in drought conditions and review some of the traits associated with yield and some selected superior genotypes, 30 genotypes of durum wheat with a test originated in Iran and Azerbaijan in 2008-09 crop year, agricultural research stations, Azad University of Ardebil was performed. The analysis of variance showed significant differences between the traits evaluated in terms of stress and there was no tension. Also among genotypes in terms of height, main spike length, grain weight and there was a significant difference in yield. Performing analysis Factor, through analysis, principal 5 components 82/58 percent of total operating changes were justified. The results indicate the importance of factor coefficients characteristics of fertile tillers, grain weight original lavender, seed weight and harvest index selected genotypes is desirable for dry conditions.

Key words: Drought • Durum wheat • Factor analysis

INTRODUCTION

Tetraploid durum wheats (*T. durum*) or hard wheat mainly used for flour to produce Semolina, especially pasta, pasta and grits be Cultivation. Although the cultivation of tetraploid wheats is low but their resistance to disease and environmental stress than wheat is more hexaploid [1]. Considering that performance is a polygenic adjective and its heritability is high to achieve high yield, selection used by Performance components [1]. Dryness of the most important factor limiting production of crops including wheat in the world and Iran. This Topic is more important in dry and semi-arid regions of the world [2]. Importance of this subject is determined when we know which more than 1/4 part ground is dry and estimated that about 1/3 of the world's cultivable land under water shortage conditions are in range [2]. Yield and the adjective little are controlled by many genes. Heritability of this trait also due to the interaction of genotype and environment, so choose based on lower yield in order to improve it may not be very effective [3]. Especially early generations of the large number of genotypes and genotypes assessed as having repeated testing does not

return if no good genetic [4]. Morphological traits simply be measured with great precision and quality relatively high heritability for plant communities and improve screener performance is [5]. Decay correlation coefficients between different traits with grain yield to decisions about the relative importance of these attributes and their values as selection criteria helps [6]. According to several reports, between grain yield in wheat and grain weight, fertile tillers or spikes per plant, spikelet's per spike and spike has a significant correlation [7, 8]. Solidarity, especially in plant height and heading time, different results (depending on variety and planting systems used) is seen [9]. Renold and co-workers[10] with different grain Simit review concluded that a whole wheat linear relationship between stress and the yield is. Gupta *et al.* 17 attribute the generation of 40 advanced lines of wheat with 11 controls in a randomized complete block design were evaluated. Factor analysis, 15 traits associated with yield and grain quality to address five main characteristics of spike, grain characteristics and quality protein and reduced tillering [11]. Dawari and Luthra[12] in his studies on bread wheat cultivars showed that the harvest index, kernels per ear per plant and spike length were important

components of performance and selection It could be the basis for improved performance to be effective. given that a significant proportion of land under wheat cultivation in arid and semiarid regions has been the aim of this research employing statistical methods on factor analysis the resulting data, to review the structure of complex traits and determine the relative importance of traits associated with performance, to identify genotypes resistant to drought, for use in future breeding programs for high yield per unit area in dry conditions.

MATERIAL AND METHODS

Twenty two durum wheat cultivars (*Triticum durum* Desf.) with Iran and Azerbaijan republic region were chosen for the study based on their reputed differences in yield performance under irrigated and non-irrigated conditions (Table 1).

Experiments were conducted at the experimental field of Islamic Azad University of Ardabil, in Ardabil province (Northwest of Iran) in 2008-2009. Seeds were hand drilled and each genotype was sown in five rows of 1.5 m, with row to row distance of 0.20 m. The experiment was laid out in randomized complete block design (RCBD) with two replications. Two levels of stress treatments including:

- Full irrigation (100 percent water based on plant needs wheat cultivars at different growth stages).
- Limited irrigation (Supply plant water needs until pollination stage and then Format water until the end of wheat growth and development).

Every line in 5 rows and 20 cm intervals and 150 cm in width were planted. Immediately after planting the field was irrigated to soil moisture profiles in root development

and saturated and identical for all treatments in addition to the germination easily is done. Irrigation was done with leaking method. After harvest to evaluate the factors affecting the performance traits, plant height, tiller number total, fertile tillers, number of internodes, peduncle length, length of main spike, spike original weight, awn length, total dry weight, number of seeds per main spike and main spike grain weight were measured.

Data were analyzed using SPSS16 for analysis of variance and Duncan's multiple range tests was employed for the mean comparisons.

RESULTS

The results of analyses of variance for grain yield and other related traits in both stress and non-stress environments are given in Table 2. There was a significant difference among stress conditions for grain yield and other traits. Except as total number of tillers and fertile tillers remaining traits were Significant in 0.01 percent probability level. The genotypes showed significant differences in grain yield and other traits. Traits of total number of tillers, total plant weight and seed number per main spike were non-significant, traits of fertile tillers and harvest index 0.05 percent level and other traits were significant in 0.01 percent level. Thus, indirect selection for a drought-prone environment based on the results of optimum conditions will not be efficient. These results are in agreement with those of Sio-Se Mardeh *et al.* [13] and Bruckner and Froberg [14].

Table 2 Average, minimum, maximum, standard deviation and coefficient of variation shows phenotypic traits. Of all traits, a large variation between wheat cultivars studied further was found. Highest percentage of phenotypic changes related to the main spike weight and total weight the plant.

Table 1: Origin and taxonomy of durum wheat landraces tasted

no	name	Landraces	origin	no	name	Landraces	origin
1	Hordeiforme	Miyane	Iran	12	leucumelan	Naxcivan	Azerbaijan
2	Africanum	Sanandaj	Iran	13	albiprovinciale	Qu	Azerbaijan
3	lecurum	kermanshah	Iran	14	murceinse	Naxcivan	Azerbaijan
4	melanopus	Ahar	Iran	15	lecurum	Lerik	Azerbaijan
5	hordeiforme	Maragheh	Iran	16	leucumelan	Naxcivan	Azerbaijan
6	lecurum	Sarab	Iran	17	apulicum	11010	Iran
7	lecurum	Tabriz	Iran	18	melanopus	hasanbaruq	Azerbaijan
8	melanopus	Cheiltoxm	Azerbaijan	19	hordeiforme	Langan	Azerbaijan
9	hordeiforme	shamxi	Azerbaijan	20	apulicum	Ardabi	Iran
10	apulicum	xanlar	Azerbaijan	21	boeffi	Ardabi	Iran
11	boeffi	shaxi	Azerbaijan	22	aboscurum	Ardabi	Iran

Table 2: Results of Analysis of variance for studied traits

MS							
S.O.V	df	Plant height	Total tillers	Fertile tillers	Peduncle length	Main spike length	Main spike weight
Rep	1	739.96**	2.13	0.031	649.80**	1.77*	0.012
Condition	1	7040.78**	0.13	1.65	2465.03**	10.09**	3.64**
Genotype	21	713.54**	3.041	4.15*	179.01**	1.71**	0.406**
C*G	21	278.74**	3.78*	2.906	150.54**	0.36	0.07
Error	43	113.36	2.205	2.46	37.63	0.61	0.15
MS							
S.O.V	df	Total plant weight	grains per main spike	Grain weight per main spike	1000 grain weight	yield	Harvest index
Rep	1	77.4	1.11	0.013	50.16	15.72	64.96
Condition	1	747.17**	120.11**	4.11**	217.54**	1216.901**	647.09**
Genotype	21	71.84	15.44	0.23**	32.41**	1516.86**	45.35*
C*G	21	58.17	12.72	0.05	23.13	953.4*	46.48*
Error	43	64.36	17.16	0.07	18.63	474.41	27.84

** And * significant at the 0.01 and 0.05 levels, respectively

Table 2: minimum, maximum, mean, standard deviation and percentage changes in phenotypic traits in genotypes

Traits	Mean	minimum	maximum	standard deviation	Percentage of phenotypic changes
Plant height	118/01	81/07	155/45	3/04	2/57
Total tillers	6/77	4/17	10/87	0/25	3/69
Fertile tillers	5/27	3/78	8/34	0/2	3/79
Main spike length	6/18	5/15	7/41	0/13	2/10
Main spike weight	2/35	1/68	4/35	0/09	3/82
Total plant weight	18/77	10/13	28/66	0/72	3/83
grains per main spike	26/16	21/91	33/55	0/51	1/94
Grain weight per main spike	1/56	1/07	2/25	0/05	3/20
1000 grain weight	55	47/25	59	0/62	1/12
yield	88/63	60/73	126/6	2/73	3/08

Table 3: Special Roots in factor analysis by considering all the traits and eliminating performance

Operating	Without yield		With yield	
	Cumulative percentage variance	Percent variance	Cumulative percentage variance	Percent variance
1	28/73	28/73	26/75	26/79
2	52/18	23/44	48/54	21/75
3	66/05	13/87	61/40	12/85
4	75/71	9/66	73/68	12/28
5	85/08	9/36	82/58	8/89

Since the correlation coefficients may complete information on the relationship between different traits and not to provide benefits according to several multivariate statistical analysis to understand the deep structure of data, factor analysis was used. Table 3 to the analysis Operating shows. Note that in terms of entering or not yield the factor analysis of differences of opinion among experts there [15], so in order to compare the two views presented in two modes, factor analysis For existing

data was conducted. Damania and Jackson [16] as examples in the factor analysis did not yield the intervention. While most researchers entering the performance together with the other characters in the factor analysis were emphasized [17,18]. As seen in Table 3, based on Eigen values greater than one in both factor analysis with regard to five factors was performed. The total factor 82/58 percent (performance) and 85 / 08 percent (no function) changes in the received data.

Table 4: Factor Analysis of water considering all the traits

Traits	Operating				
	1	2	3	4	5
Plant height	0/475	-0/29	-0/64	0/36	-0/24
Total tillers	0/536	-0/62	0/35	-0/27	0/17
Fertile tillers	0/622	-0/64	0/29	-0/098	-0/011
Main spike length	0/071	0/38	-0/63	-0/16	0/48
Main spike weight	0/44	0/66	-0/10	-0/14	0/041
Total plant weight	0/88	-0/16	-0/13	0/042	0/19
Original number of seeds lavender	0/28	0/63	0/077	-0/58	-0/063
Main lavender seed weight	0/53	0/68	0/16	0/074	-0/016
1000-Grain weight	0/45	0/55	0/15	0/50	-0/10
Yield	0/23	0/14	0/26	0/78	-0/020
Harvest index	-0/35	0/28	0/65	0/090	0/43

Table 5: Operating simple correlation coefficients with grain yield

Operating	Yield
1	0/23
2	0/14
3	0/26
4	0/78**
5	-0/027

First factor the highest volume (26/79 percent) of changes in the data was a positive and large coefficients for weight symbols, total plant weight and tiller number were we can as a factor in plant weight and performance considered. The second factor (21/75 percent) of changes in the data fills large and positive coefficients for grain weight per main lavender, seed number and weight of the original lavender lavender was the original as we can factor in the spike characteristics Rate Nanm said. damania and Jackson [16] in the third factor as spike features were introduced. These coefficients indicate that the genotypes have high levels of second factor regardless of other characteristics has a long and fertile spike with more grain number and grain weight would be greater. Spike components (length, spikelet number, number of fertile florets and grain number) have more impact on performance. If there are moisture and appropriate and accessible food, genotypes that have high spike length, number of spikelet's and fertile florets and spike them is greater, will yield a large (1.) feet tall as well as selecting varieties resistant to lodging and plant height (75 to 100 centimeters) in dry conditions can be useful for the following reasons [15]:

- More photosynthetic material can be stored in the stem
- Mechanized harvesting, especially in the drought conditions that cause extreme short leg is easily possible

- Is a straw in the soil can remain in storage at the field level during the descending sky fall and winter to keep moisture to be effective and save.

The third factor in having (12/85 percent) of the changes has positive and large coefficients for the total number of tillers and fertile tillers were. Having the fourth factor (12.28 percent) of the changes has positive and large coefficients for height and yield was. damania and Jackson [16] reported in the third factor. Evaluation of advanced bread wheat genotypes has shown that more figures with a height of Early access and enjoy high performance [9]. Having the fifth factor (8 / 89 percent) of the change has large and positive coefficients for the length lavender harvest index was the original. rostaei [9] the relationship between grain yield quantitative traits through factor analysis in wheat expressed changes from entering or failure in performance analysis results were not significant impact on factor was achieved. Bramel[17] while race (quoting 3) by performing factor analysis with and without grain yield reported in the second case (remove the grain) the number of operating changes are less justified. With reference to Table 3 the factor coefficients related traits in the analysis, regardless of operating performance shows we can get results largely inconsistent with the results and when the factor analysis without yield higher percentage was calculated to justify the changes. So, it looks like their selection based on performance efficiency can be less so based on genotype factor values in factor analysis, regardless of performance will be possible desirable genotypes in terms of selected indicators intended to be selective. In this case, simple correlation coefficients calculated with the yield factor, which is reflected in Table 5 indicate that a significant correlation between the fourth factor (0 / 78 **) with the performance show (Table 5).

REFERENCES

1. Khayatneghad, M., M. Zaefizadeh, R. Gholamin and Sh. Jamaati-e somarein, 2010. Study of Genetic Diversity and Path Analysis for Yield in Durum Wheat Genotypes under Water and Dry Conditions. *World Appl. Sci. J.*, 9(6): 655-665.
2. Kirigwi, F.M., M. van Ginkel, R.G. Trethowan, R.G. Sears, S. Rajaram and G.M. Paulsen, 2004. Evaluation of selection strategies for wheat adaptation across water regimes. *Euphytica*, 135: 361-371.
3. Richards, R.A., 1996. Defining selection criteria improve yield under drought plant Growth Regulation, 20: 157-166.
4. Keim, D.L. and W.E. Kronstand, 1981. Drought responses of winter wheat cultivars grown under field stress condition. *Crop Sci.*, 21: 11-14.
5. Yap, T.C. and B.L. Harvey, 1972. Inheritance of yield components and morpho-physiological traits in barley (*Hordeum vulgare* L.) *Crop Sci.*, 12: 283-286.
6. Agrama, H.A.S., 1996. Sequential path analysis of grain yield and its components in maize. *Plant breeding*, 115: 343-346.
7. Shanahan, J.F., K.J. Donnelly, D.H. Smith and D.E. Sminka, 1985. Shoot developmental properties associated with grain yield in Winter wheat. *Crop Sci.*, 25: 770-775.
8. Mohiuddin, S.H. and L.I. Cory, 1980. Flag leaf and peduncle area duration in relation to winter wheat grain yield. *Agron. J.*, 72: 299-301.
9. Rostaei, M., V. Sadeghzadeh and Y. Arshad, 2003. Relationship traits affecting grain yield by using factor analysis in dry conditions. *Knowledge of Agric.*, 13(1).
10. Renold, M.B., R.T. Skovmand, Ve Thowar and W. Pfeiffe, 2000. Wheat program, CIMMYT.
11. Gupta, A.K., R.K. Mittal and A.Z. Ziauddin, 1999. Association and factor analysis in spring wheat. *Ann. Agric. Res.*, 20: 481-485.
12. Dawari, N.H. and O.P. Luthra, 1991. Character association studies under high and low environments in wheat (*Triticum aestivum* L.). *Indian J. Agric. Res.*, 25: 515-518.
13. Sio-Se Mardeh, A., A. Ahmadi, K. Poustini and V. Mohammadi, 2006. Evaluation of drought resistance indices under various environmental conditions. *Field Crop Res.*, 98: 222-229.
14. Bruckner, P.L. and R.C. Frohberg, 1987. Stress tolerance and adaptation in spring wheat. *Crop Sci.*, 2: 31-36.
15. Tosi mojarad, M., M. Ghanadha, M. Khodarahmi and S. Shahabi, 2005. Factor analysis for grain yield and other characteristics. *Res. Construction*, 67: 9-16.
16. Damania, A.B. and M.T Jackson, 1986. An application of factor analysis to morphological data of wheat and barley landraces from the Bheri River Valley, Nepal. *Rachis*, 5: 25-30.
17. Bramel, P.J., P.N. Hinnz, D.E. Green and R.M. Shibles, 1984. Use of principal factor analysis in the study of three stem termination types of soy bean. *Euphytica*, 33: 387-400.
18. Selier, G.J. and R.E. Stafford, 1985. Factor analysis of components of yield in guar. *Crop Sci.*, pp: 905-908.