

Rainfall, Phenology of Leaf Flushing, Flower Initiation and Fruit Maturation in Dry Deciduous and Evergreen Forests of Bhadra Wildlife Sanctuary, Karnataka, Southern India

^{1,2}A. Nanda,, ¹H.M. Prakasha, ¹Y.L. Krishna Murthy and ³H.S. Suresh

¹Department of PG studies and Research in Applied Botany, Bioscience Complex,

Jnana Sahyadri, Kuvempu University, Shankaraghatta, Shimoga, 577 451, Karnataka, India

²Environmental Study Centre, Annappa Garden Santhe Kadur, Shimoga, 577 222, Karnataka India

³Center for Ecological Sciences, Indian Institute of Science, Bangalore, 560 012, Karnataka, India

Abstract: Varying with altitude and rainfall of two forest types of tropical forest were studied. Community wide pattern in both vegetative and reproductive phenophases among various tree species of Bhadra wildlife sanctuary, Karnataka is reported here. Leaf initiation peaks in the month of April in dry forest, after leaf less period of 1-2 months, in evergreen forest leaf initiation peaks in the month of January simultaneously with leaf senescence. Flower initiation begins from January till June with a peak in April. Evergreen forest starts from November to march with a peak in January and February. Fruit maturation in dry forest starts from March to December with a peak in August and November, in evergreen forest from January to July with a peak in April.

Key words: Community-wide • Tree Phenophases • Tropical forest • Varying altitude • Western ghats

INTRODUCTION

Tropical forest plant communities display various phenological patterns. Different forest types are considered to be indicators of the amount and annual distribution of rainfall Walter [1] because seasonal variation in tree water status constitutes a major determinant of tropical tree phenology Borchert [2], Borchert [3]. A variety of factors have been proposed to drive the phenology. These include abiotic factors such as rainfall, Opler [4] day length, Van Schaik [5] irradiance, Janzen [6], temperature Singh and Singh [7] and relative humidity Borchert and Wright and Van Schaik [8,9]. Biotic factors are pollinators Frankie [10] or seed dispersers Rathke and Lacey [11], herbivory (Murali and Sukumar [12], Aide [13], Williams-Linera [14]. Seasonal changes in abiotic and biotic factors can be expected to have consistent effects on phenology of tropical forests Wright [15]. There are several community wide studies on phenology. Sundarapandian [16], Singh and Kushwaha [17], Murali and Sukumar [12, 18], Prasad and Hegde [19], Bhat [20], Justiniano and Fredericksen [21], Newstrom [22], Lieberman [23], Borchert [24], McLaren and McDonald [25], Lott [26]. Yet our understanding of phenophases in different forests types is scanty.

Understanding of community wide phenophase patterns is essential for understanding the functioning of ecosystem and scientific management of natural resources. This type of study will help to understand various phenophases in the global climatic change scenario at local level.

Study Area: The study was conducted in Bhadra Wildlife Sanctuary located in Chikmagalur and Shimoga districts (13°25' and 13°50' N, 75°15' and 75°50' E) of Karnataka. The dry deciduous forest (Umbleailu, 13°.46' to 13°52' N, 75°36' to 75°. 42' E). The altitude 690 to 750 m ASL. The evergreen forest (Kemmanundi, 13°32' to 13°40' N, 75°44' to 75°45' E). These forests area common around the hills and valleys of where the altitude ranges from 1400 to 2000 m ASL. These forests are on the windward side of the Western Ghats. The rainfall of the Sanctuary ranges from 500 to 2000mm. The terrain is gently undulating with valleys and steep hillocks. Detailed geological account of the sanctuary is given by Parameswar, [27]. Rainfall data for the dry forest was collected from meteorological station, Bhadra River Project. Whereas for evergreen forest rainfall data from the coffee estate. Annual rainfall in two forest types varies in evergreen forest 1561.1 mm and in dry forest 542 mm during the study period is shown in (Fig. 1).




Fig. 1: Annual total rainfall in dry and evergreen forest during the study period (june-2004 to may-2006) of Bhadra Wildlife Sanctuary

Vegetation: Vegetation of the sanctuary varies from dry deciduous to evergreen forest. According to Champion and Seth [28] the dry deciduous forests of Umblebailu are classified as ‘southern dry mixed deciduous forests’. The characteristic tree species of this type are *Terminalia paniculata*, *Anogeissus latifolia*, *Xylia xylocarpa*, *Cassia fistula*, *Albizia lebbek*, *Terminalia crenulata*, *Tectona grandis*, *Diospyros montana*, *Mitragyna parviflora* and *Pterocarpus marsupium*.

Kemngundi forests are classified as ‘Tropical wet evergreen forests’. The characteristic tree species of this forest are *Artocarpus integrifolia*, *Cinnamomum* spp., *Myristica malabarica*, *Litsea* spp., *Neolitsea zeylanica*, *Syzygium* spp., *Macaranga peltata*, *Trema orientalis*, *Actinodaphne hookeri* and *Isonandra perrottetiana*.

Detailed description of the study area can be found elsewhere Raju and Hegde [29], Parameswar [27]. The present study describes the certain phenological patterns of tropical tree communities in dry deciduous forest and evergreen forest at Bhadra Wildlife Sanctuary for 2 years during June 2004 to May 2006.

MATERILAS AND METODS

One transects each in evergreen forest (10 Km) long and in dry deciduous forest (2 Km) long was laid along the periphery of sanctuary for clear visibility of the tree canopy. A total of 177 individuals comprising of 47 tree species in site-I and 277 individuals comprising of 45 tree species in site-II all above 20 cm d.b.h (diameter at breast

height) were identified using various regional floras Gamble and Fischer [30] madras, Saldhana [31] karnataka, Yoganarasimhan [32] Chikkamagalur, Ramaswamy [33] shimoga, Neginhal [34] south India. The representative plant specimens were collected, deposited in the Herbarium of the department of Applied Botany, Kuvempu University. Identified individuals were marked with a unique tag numbers with clear visibility to facilitate re-location. These marked individuals were monitored for both vegetative (foliar) and reproductive phenological events such has for leaf flushing (LF2), flower budding (FL2) and fruit maturation (FR4) phenologies. Each stage in different categories of phenology was scored qualitatively with respect to both spread and intensity on canopy on a 0 to 100 scale.

Data Analysis: Correlations analysis relating to rainfall parameter was computed. Differences in the patterns of frequency of leaf flushing and flower initiation in a month between two forest types were analysed using the Kolmogorov-Smirnov test (KS test) and Chi-Square test by Zar [35].

RESULTS

Leaf initiation phenology: Leaf initiation peaks in the month of April in dry forest, after leaf less period of 1-2 months. Whereas leaf initiation peaks in the month of January simultaneously along with leaf senescence in evergreen forest (Fig. 2).




Fig. 2: Leaf flushing pattern in dry and evergreen forest of Bhadra Wildlife Sanctuary




Fig. 3: Leaf flushing pattern and rainfall in dry forest of Bhadra Wildlife Sanctuary




Fig. 4: Leaf flushing pattern and rainfall in evergreen forest of Bhadra Wildlife Sanctuary




Fig. 5: Flower budding in dry and evergreen forest of Bhadra Wildlife Sanctuary




Fig. 6: Leaf flushing and flower budding in dry forest of Bhadra Wildlife Sanctuary




Fig. 7: Leaf flushing and flower budding in evergreen forest of Bhadra Wildlife Sanctuary




Fig. 8: Fruit maturation in dry and evergreen forest of Bhadra Wildlife Sanctuary




Fig. 9: Fruit maturation and rainfall in dry forest of Bhadra Wildlife Sanctuary




Fig. 10: Fruit maturation and rainfall in evergreen forest of Bhadra Wildlife Sanctuary

Factors influencing leaf flushing phenology: In dry deciduous forest, Rainfall had significant negative influence on leaf flush during two month lag period ($r = -0.62$, $p < 0.001$) along with number of rainy days ($r = -0.61$, $p < 0.002$) It is clear from the (Fig. 3) that peak leaf flush happens two months before the peak in rainfall. However in evergreen forest, Rainfall had significantly negative influence on leaf flush during the corresponding months ($r = -0.41$, $p < 0.04$). Time lag correlations with rainfall were not significant either during one-month lag period or two-month lag period. Leaf flush peaks five months before the rainfall (Fig. 4).

Flowering phenology: Flower initiation begins from January till June with a peak in April. i.e. most species initiate flowering in leafless period and few species simultaneously leafing and flowering in summer period. But the pattern in evergreen forest starts from November to march with a peak in January and February in post-winter to pre-summer (Fig. 5).

Factors influencing flower initiation phenology: In dry deciduous forest flower initiation during two month lag period. Rainfall ($r = -0.49$, $p < 0.02$) and number of rainy days ($r = -0.57$, $p < 0.005$) during corresponding month had a significant negative influence.

Most of the species initiate flower during dry season in dry deciduous forest. It is also clear from the figure that the initiation begins in January and continues till June with peaks during April to June. Leaf initiation and flower initiation coincide (Fig. 6). But some of the species do initiate flower while having senescent leaves or during the leafless period. In dry deciduous forest some trees had leaf initiation and flower initiation coincide with each other (KS test $D = 0.2916$ $P > 0.10$). In dry deciduous forest leafing and flowering events had taken place simultaneously. Chi-Square ($\chi^2 = 83.74$ $df = 23$ $p < 0.0001$) is highly significant.

Whereas in evergreen forest rainfall during the corresponding months has a significant negative influence on the initiation of flowers among the species ($r = -0.57$, $p < 0.003$). Time lag correlations are not significant. Flower initiation and leaf flush happens simultaneously among the species (KS test, $D = 0.333$, $p > 0.10$), though in some months there are more number of species initiating flowers than having young leaves Chi-Square ($\chi^2 = 71.87$, $p < 0.0001$, $N = 23$) is highly significant (Fig. 7).

Fruiting phenology: Fruit maturation in dry forest starts from March to December with a peak in August and November. From summer periods to end of rainy season.

Whereas in evergreen forest from January to July with a peak in April i.e., fruit maturation in evergreen forest starts from onset of winter to a peak rainfall (Fig. 8).

Factors influencing fruit maturation phenology: In dry forest number of rainy days had a significant negative influence on maturity of fruits during corresponding months. Multiple regression during one month lag period was significant ($r = 0.65$, $F = 3.43$, $p < 0.02$) with number of rainy days and rainfall influencing the event. Step wise regression was also significant ($r = 0.65$, $F = 4.82$, $p < 0.01$) with number of rainy days and rainfall influencing the event. Though rainfall influences maturity of fruits, it did not have significant impact on the maturity of fruits (Fig. 9).

Similar pattern was observed with maturity of fruits in evergreen forest also. Correlation during corresponding month was not significant with respect to fruit maturity (Fig. 10). Time lag correlations were negatively significant ($r = -0.66$, $p < 0.004$, one-month lag) ($r = -0.64$, $p < 0.001$, two-month lag).

DISCUSSION

Dry periods in tropical dry forests are characterized by intense physiological activities. Most of the phenological activities happen during the early of the dry season in evergreen forest and in dry season in dry forest. In this present study leaf flushing and leaf fall happens before the onset of rains in evergreen forest of Kemmangundi. This synchronization between senescence and flushing ensures that trees were never totally leafless. This pattern is similar to other studies (Ralhan [36], Saha [37], Shukla and Ramakrishnan [38], Bhat [20]).

However in dry forest leaf flushing is in dry season after a 1 - 2 month of leafless period. i.e. before the onset of rains is in agreement with other observation, Murali and Sukumar [12] mudumalai, Singh and Singh [7] vindhyana plateau, Prasad and Hegde [19] bandipur, Kushwaha and Singh [39] Hathinala forest of vindhyana plateau, Justiniano and Fredericksen [21] Bolivia.

Flower initiation also differs with respect to forest type in this study. As evergreen trees initiated flowering in the winter period is similar to other studies. Bhat [20] Uttara Kannada, Karnataka, Sundarapandian [16] kodayar in the Western Ghats, Tamilnadu and Saha [37] Darjeeling. Whereas in dry forest trees initiated flowering during the beginning of the dry season at the time most of the trees were leafless or leaf flushing stage. As flower initiation can advertise to pollinators as they get pollinated as seen

in other tropical forest (Murali and Sukumar [18], Sivaraj and Krishnaswamy [40], Sundarapandian [16]. Opler [4] found that in Costa Rica rainfall (wet season) play a role in flower initiation.

Reproductive phenological studies in seasonal tropical forest ecosystems have indicated rainfall seasonality as being the major abiotic factor controlling the timing, Koelmeyer [41], Smythe [42], Frankie *et al.* [10], intensity, Lieberman [23], Lott *et al.* [26], De Lampe *et al.* [43], Newstrom *et al.* [22] and duration White [44], Sun *et al.* [45], Justiniano and Fredericksen [21], Borchert *et al.* [24] of flowering and fruiting periodicities. Van Schaik *et al.* [5] deduced that leaf flushing and flowering fell within 1 month before the onset of the rainiest period in strongly seasonal forests. In deciduous trees, anthesis can be induced by the temporal rehydration of the trees after leaf fall by isolated rainfall during the dry season or by the onset of the wet season. Reich and Borchert 1984 [46].

For these reasons, it has been assumed that water availability is both the proximate and the ultimate factor controlling phenology in tropical dry forest plants Reich and Borchert [46]. The timing of synchronous flowering in individual species of tropical trees and the resulting flowering periodicity at the community level are widely thought to have evolved as a result of biotic interactions between trees and their pollinators. Borchert [24].

The duration of fruit maturation was longest in dry forest from August to April (monsoon to pre-monsoon period) with a peak in November to January. This pattern is in agreement with other tropical studies. Murali and Sukumar [18], Sundarapandian [16], Prasad and Hegde [19]. However in evergreen forest the Fruit maturation was from January to July with a peak in the month of April. Maturation was more in pre-monsoon and lessens after a monsoon.

However, tropical wet forests exhibit a wide range of fruiting patterns, including unimodal or bimodal fruiting peaks. Foster [47], Zhang and Wang [48] 1995, Hamann [49]. Some tropical dry forest plant phenology studies describe a single fruiting peak related to the dry season or several peaks in the wet and dry seasons. Frankie *et al.* [10], Opler *et al.* [4], Bullock and Solis-Magallanes [50].

ACKNOWLEDGEMENTS

We thank Department of Environment, Ministry for Environment and Forests, Government of India for funding this project. We thank Forest department of Karnataka for permission to work in Bhadra sanctuary.

REFERENCES

1. Walter, H., 1971. Ecology of tropical and subtropical vegetation. Oliver and Boyd, Edinburgh.
2. Borchert, R., 1994a. Water storage in soil or tree stems determines phenology and distribution of tropical dry forest trees. *Ecology*, 75: 1437-1449.
3. Borchert, R., G. Rivera and W. Hagnauer, 2002. Modification of vegetative phenology in a tropical semi deciduous forest by abnormal drought and rain. *Biotropica*, 34: 27-39.
4. Opler, P.A., G.W. Frankie and H.G. Baker, 1976. Rainfall as a factor in the release, timing and synchronization of anthesis by tropical trees and shrubs. *Journal of Biogeography*, 3: 231-236.
5. Van Schaik, C.P., J.W. Terborgh and S.J. Wright, 1993. The phenology tropical forests: significance and consequences for primary consumers. *Annual Review of Ecology and Systematics*, 24: 333-357.
6. Janzen, D.H., 1967. Synchronization of sexual reproduction of trees with in dry season in Central America. *Evolution*, 21: 620-637.
7. Singh, J.S. and V.K. Singh, 1992. Phenology of seasonally dry tropical forest. *Current Science*, 63: 684-689.
8. Borchert, R., 1994. Induction of rehydration and bud break by irrigation or rain in deciduous trees of a tropical dry forest in Costa Rica. *Trees*, 8: 198-204.
9. Wright, S.J. and C.P. Van Schaik, 1994. Light and the phenology of tropical trees. *American Naturalist*, 143: 192-199.
10. Frankie, G.W., H.G. Baker and P.A. Opler, 1974. Comparative phenological studies of trees in tropical wet and dry forests in the lowlands of Costa Rica *Journal of Ecology*, 62: 881-919.
11. Rathcke, B. and E.P. Lacey, 1985. Phenological patterns of terrestrial plants. *Annual Review of Ecology and Systematics*, 16: 179-214.
12. Murali, K.S. and R. Sukumar, 1993. Leaf flushing phenology and herbivory in a tropical dry deciduous forest, southern India. *Oecologia*, 94: 114-120.
13. Aide, T.M., 1993. Patterns of leaf development and herbivory in a tropical understorey community. *Ecology*, 74: 456-466.
14. Williams-Linera, G., 1999. Leaf dynamics in a tropical cloud forest: phenology, herbivory and life span. *Selbyana*, 20: 98-105.
15. Wright, S.J., 1996. Phenological responses to seasonality in tropical forest plants. *Tropical forest ecophysiology*. N-York, Chapman and Hall. pp: 440-460.

16. Sundarapandian, S.M., S. Chandrashekar and P.S. Swamy, 2005. Phenological behaviour of selected tree species in tropical forests at kodayar in the Western Ghats, Tamilnadu, India. *Current Science*, 88: 805-810.
17. Singh, K.P. and C.P. Kushwaha, 2005. Emerging paradigms of tree phenology in dry tropics. *Current Science*, 89: 964-975.
18. Murali, K.S. and R. Sukumar, 1994. Reproductive phenology of a tropical dry forest in Mudumalai, southern India. *Journal of Ecology*, 82: 759-767.
19. Prasad, S.N. and M. Hegde, 1986. Phenology and seasonality in the tropical deciduous forest of Bandipur, South India. *Proceedings of Indian Academy of Sciences. (Plant Sci.)* 96: 121-133.
20. Bhat, D.M., 1992. Phenology of tree species of tropical moist deciduous forest of Uttara Kannada district, Karnataka, India *Journal of Bioscience* 17: 325-352.
21. Justiniano, M.J. and T.S. Fredericksen, 2000. Phenology of tree species in Bolivian dry forests. *Biotropica*, 32: 276-281.
22. Newstrom, L.E., G.W. Frankie and H.G. Baker, 1994. A new classification for plant phenology based on flowering patterns in lowland tropical rain forest trees at La Selva, Costa Rica. *Biotropica*, 26: 141-159.
23. Lieberman, D., 1982. Seasonality and phenology in a dry tropical forest in Ghana. *Journal of Ecology*, 70: 791-806.
24. Borchert, R., S.A. Meyer, R.S. Felger and L. Porter-Bolland, 2004. Environmental control of flowering periodicity in Costa Rican and Mexican tropical dry forests. *Global Ecology and Biogeography*, 13: 409-425.
25. McLaren, K.P. and M.A. McDonald, 2003. Seedling dynamics after different intensities of human disturbance in a tropical dry limestone forest in Jamaica. *Journal of Tropical Ecology*, 19: 567-578.
26. Lott, E.J., S.H. Bullock and J.A. Solis-Magallanes, 1990. Phenology of canopy trees of a tropical deciduous forest in Mexico. *Biotropica*, 22: 22-35.
27. Parameshwar, G., 1996. Draft Management Plan for Bhadra Wildlife Sanctuary (1996-2001), Publication: Wildlife Division, Chikkamagalur, Karnataka Forest Department, pp: 43.
28. Champion, H.G. and S.K. Seth, 1968. A Revised Survey of the forest types of India, Govt. of India Press.
29. Raju, R. and S.N. Heggde, 1995. Bhadra Wildlife Sanctuary, a Fragile Ecosystem. Inidan Forester, pp: 938-948.
30. Gamble, J.S. and C.E.C. Fischer, 1998. Flora of the Presidency of madras, Adlard and Son, Ltd, 21, Hart Street, W.C., pp: 1-3.
31. Saldanha, C.J., 1984-1996. Flora of Karnataka Vol-I-IV. Oxford and IBH Publishing Co. Pvt. Ltd., New Delhi.
32. Yoganarasimhan, S.N., K. Subramanyam and B.A. Razi, 1982. Flora of Chikkamagalur District, Karnataka, India. International Book Distributors, Deharadun.
33. Ramaswamy, S.N., M.R. Rao and D.A. Govindappa, 2001. Flora of Shimoga District, Karnataka', Prasaranga University of Mysore, Manasagangotri, Mysore.
34. Neginhal, S.G., 2004. Forest trees of South India. Navbharath Press, Bangalore.
35. Zar, J.H., 2007. Biostatistical analysis. Pearson Education (Singapore) Pte. Ltd.
36. Ralhan, P.K., R.K. Khanna, S.P. Singh and J.S. Singh, 1985. Phenological characteristics of the tree layer of Kumaun Himalayan forests. *Vegetatio*, 60: 91-101.
37. Saha, S., 2007. Study of leafing, flowering and fruiting activities in three species of temperate forest at the Singalila range, Darjeeling. *Indian Journal of Ecology*, 34: 15-18.
38. Shukla, R.P. and P.S. Ramakrishnan, 1982. Phenology of trees in sub-tropical humid forest in north-eastern India. *Vegetatio*, 49: 103-109.
39. Kushwaha, C.P. and K.P. Singh, 2005. Diversity of leaf phenology in a tropical deciduous forest in India. *Journal of Tropical Ecology*, 24: 47-56.
40. Sivaraj, N. and V.K. Krishnamurthy, 1989. Flowering phenology in the vegetation of Shervoy's, south India. *Vegetatio*, 79: 85-88.
41. Koelmeyer, K.O., 1959. The periodicity of leaf change and flowering in the principal forest communities of Ceylon. *Ceylon Forester*, 4: 157-189.
42. Symthe, N., 1970. Relationships between fruiting seasons and seed dispersal methods in a neotropical forest. *American Naturalist*, 104: 25-36.
43. De Lampe, M.G., Y. Bergeron, R. Mcneil and A. Leduc, 1992. Seasonal flowering and fruiting patterns in tropical semi-arid vegetation of Northeastern Venezuela *Biotropica*, 24: 64-76.
44. White, L.J.T., 1994. Patterns of fruit-fall in the Lope Reserve, Gabon. *Journal of Tropical Ecology*, 10: 289-312.
45. Sun, C., B.A. Kaplin, K.A. Kristensen, V. Munyaligoga, J. Mvukiyumwami, K.K. Kajondo and T.C. Moermond, 1996. Tree phenology in a tropical montane forest in Rwanda. *Biotropica*, 28: 668-681.
46. Reich, P.B. and R. Borchert, 1984. Water stress and tree phenology in a tropical dry forest in the lowlands of Costa Rica. *Journal of Ecology*, 72: 61-74.

47. Foster, R.B., 1982. The seasonal rhythms of fruit fall on Barro Colorado Island, In E.G. Leigh, A.S. Rand and D.M.Windsor (Eds.). The ecology of a tropical forest: Seasonal rhythms and long-term changes. Smithsonian Institution Press, Washington, D.C., USA.
48. Zhang, S. and L. Wang, 1995. Comparison of three fruit census methods in French Guiana. *Journal of Tropical Ecology*, 11: 281-294.
49. Hamann, A., 2004. Flowering and fruiting phenology of a Philippine submontane rain forest: Climatic factors as proximate and ultimate causes. *Journal of Ecology*, 92: 24-31.
50. Bullock, S.H. and J.A. Solis-Magallanes, 1990. Phenology of canopy trees of a tropical deciduous forest in México. *Biotropica*, 22: 22-35.