Soil Quality Modeling of a Highly Acidic Eutric - Tropofluent Soil

Uzoije Atulegwu Patrick

Department of Environmental Engineering,
School of Engineering and Engineering Technology,
Federal University of Technology, Owerri, Imo State, Nigeria

Abstract: Waste from garri processing centre of Umuagwo, Ohaji L.G.A of Imo state has impacted enormous pollution on the soil ecosystem of the centre. Huge resources and quality man hour have been spent on determining the level of such pollution in similar circumstances using laboratory techniques. In view of the aforementioned, the study developed several regression models, establishing the relationship between Ph and other selected soil physiochemical properties, to predict the values of the these selected properties in the polluted soils. The study took place in three sampling pedons A, B and C using the technique of stratified random sapling for sample collection. Pedon A is the background or the control unit, pedon B or the discharge point is the cassava mill effluent (CME) receiving unit and pedon C is the down stream, 500m away from pedon B. the result showed high values of soil physiochemical properties(mainly the heavy metals) at pedon B and the trend reduced away from pedon B and lowest in pedon A. In terms of pedons the heavy metal concentrations took this trend; pedon B>pedon C>pedonA. pH relationship with heavy metals in pedons A and B showed positive linear relationships with their coefficient of linearity(R²) ranging between 0.70-0.92 while other soil physiochemical properties obeyed polynomial relationships at reasonably high coefficient of fitness. Polynomial models fitted the pH relationships virtually in all the selected physiochemical properties of pedon C. The models have provided quick assessment of these physiochemical materials by mere determining the pH value and substituting the pH values into the models to predicting other properties.

Key words:Regression models • Cassava • Effluent • Pollution • Pedons • Soil • Physiochemical
• Properties • Pedons • Heavy metals

INTRODUCTION

In Nigeria, Cassava(*Manihot esculent*) can be processed into different kinds of food such as fufu, garri, tapioca etc. [1]. Garri production is most practiced in Nigeria and it is carried out at varying scales; in a small, medium and large scales. Most garri processing plants in Nigeria produce between 7-10 million tones of garri annually. [2]. Processing of the this product requires large volume of water which in addition to some other solid substances forms part of the waste. This waste contains measurable quantity of heavy metals in a low acidic range. [3]. Apart from constituting foul smell and unattractive sight, the cassava mill effluent (CME) can also upset the ecological balance of the receiving bodies [4]. Francis *et al.* 2008 [5] raveled total extinction of benthic macro invertebrate specie and abundance of

oligocheats and dipterans in a marine ecosystem by CME due to its high acidic medium. Also in marine environment, low acidic value of the effluent lowers the pH value of the stream leading to difficulty in fish breeding [6]. Low pH value of the CME also influences number of edaphic factors such as clay, organic matter, cation exchange capacity and some other soil phsiochemical properties such as heavy metals [7-9]. Giving the apparent influence of pH on these physiochemical properties, developing the modeling in respect of the soil properties in such a low acidic CME polluted medium will create an enabling condition to predict their values for assessment of their pollution potential and remediation. Therefore this study models the physiochemical properties of the CME receiving soil of Ohaji cassava processing center and its environs.

Corresponding Author: Uzoije Atulegwu Patrick, Department of Environmental Engineering,
School of Engineering and Engineering Technology, Federal University of Technology,
Owerri, Imo State, Nigeria E-mail:patuzong@yahoo.com.hk.

MATERIALS AND METHODS

Description of Study Area: The study was carried out at central garri processing center at Umuagwo, in Ohaji Egbema Local Government of Imo state, South- Eastern Nigeria. The area lies within the latitude 50 12'N to 50 48'N and at the temperature range of between 250C-300C [10]. The people of the area experience high rainfall of about 2000mm-2500mm. Umuagwo is located in the sandy benin formation and therefore the geology of the region is characterized by quarternary, alluvium, meander belt, wooded back swamps as well as fresh water swamps [11]. The soil of the area is classified as Eutric Tropofluent [12]. Vegetation of the study area is that of a rain forest and the soil supports arable crop production. That explained why the major socio-economic activity of the area is cassava cultivation and processing. The garri processing centre which has been in operation for over ten years, takes up in commercial quantity. Processings ranging from cassava grinding to fermentation and frying are the major steps of the center. Liquid and some solid waste such as fibers from the processing centre are channeled to the nearby farmland and ordinary land that has been left fallow for years.

Soil Sampling Techniques and Analytical Methods:

The entire study area was divided into three morphological units or pedons; namely the background (BA) or control, the waste receiving area or point of discharge (PD) and down stream (DS) of the discharged waste or the waste drainage channel. The three morphological units represent pedon A, poedon B and pedon C respectively. A transect was drawn to link the three units. Given the high mobility of most substances constituting the cassava mill effluent in the soil, soil sampling was extended up to 100cm in depth. Along the transect, three pedons were dug at the inter-pedon distance of 500m. At each pedon, soil samples were collected from different soil layers corresponding to 0-15, 15-30, 30-70 and 70-100cm in depth with a sterilized soil auger which was rinsed with a lot of distilled water and dry cleaned after every sampling to avoid contamination. The layers were designated as L1, L2, L3 and L4 respectively. The samples collected from each unit and layer were done in three replicates at the interval of 20m, making a total sample of 36 soil samples for this study. The soil samples were subjected to various laboratory analysis using the following analytical

methods after processes of air-drying, crushing and sieving using 2mm sieve; Hydrometer method as conducted by Gee and Or, (2002) [13] and was used to analyzed for the particle size distribution, the SOLAAR UNICAM 969 atomic absorption spectrometer (AAS) was used to analyze all the metals [14,15]. Core method of Grossman and Reinsch, 2002 [16], was used for the analysis of bulk density. Moisture content was determined using the gravimetric method as carried out by Obi, 1999 [17]. Organic carbon was determined directly by furnace combustion at 3790C.

Model Approach: The models used for this study are the linear regression models of the least square method. Regression model was favored due to its reliability even in the face of limited data and ease of calibration and application. The model was developed based on the following conditions;

- That the flow of the CME down the soil horizon assumed one-dimensional flow
- That the concentration of the effluent on each horizon is homogenous
- That the samples collected at various sampling points be the representative of the study area.
- The models were calibrated with the average concentration of heavy metals and other physiochemical soil properties obtained from the analysis.

The linear regression or the predicted model was presented after Nwaogzie (1998) [18] as

$$y = \alpha_0 + \alpha_1 x \quad (1)$$

Where α_0 and α_1 are the regression coefficients. Y and X are the dependent and independent variables respectively. Method of least squares was used to estimate α_0 and α_1 in order to form the line of best fit to the data. For each data point there is always a difference between observed value of y, y_{obs} and the predicted value y_{pre} . The difference measures the error (e_i) or the residue and is denoted as

$$y_{obs} - y_{pre} = e_i \quad (2)$$

To measure the goodness of fit for a given data, $e_1 + e_2 + e_3 + \dots + e_n$ must be minimum. This is mathematically represented as

$$S = \sum_{i=1}^n e_i^2 = \text{minimum} \quad (3)$$

Substituting equation 2 into 3, it transforms to

$$S = \sum_{i=1}^n (y_{obs} - y_{pre})_i^2 = \text{minimum} \quad (4)$$

Substituting the y_{pre} as in equation 1 into equation 4, it becomes

$$S = \sum_{i=1}^n (y_{obs} - a_0 + a_1 x_i)^2 = \text{minimum} \quad (5)$$

To solve for α_0 and α_1 partial derivative was made of equation 5 with respect to α_0 and α_1 to get

$$\frac{\partial S}{\partial a_0} = \sum 2(y_{obs} - a_0 - a_1 x_i)(-1) = 0 \quad (6)$$

$$\frac{\partial S}{\partial a_1} = \sum 2(y_{obs} - a_0 - a_1 x_i)(-x_i) = 0 \quad (7)$$

Equations 6 and 7 can be rearranged to get the following pair of equation.

$$na_0 + a_1 \sum x_i = \sum y_i \quad (8)$$

$$a_0 \sum x_i + a_1 \sum x_i^2 = \sum x_i y_i \quad (9)$$

Equations 8 and 9 can be solved simultaneously to get α_0 and α_1 as

$$a_i = \frac{n \sum x_i y_i - \sum x_i \sum y_i}{n \sum x_i^2 - (\sum x_i)^2} \quad (10)$$

Substituting equation 10 into 8 and solving for α_0 results to

$$a_0 = \frac{\sum y_i - a_1 \sum x_i}{n} \quad (11)$$

The quadratic regression model which is given as

$$y = a_0 + a_1 x + a_2 x^2 \quad (12)$$

Will also be used to predict the values of the selected physiochemical properties of the soil.

The least square estimates of a_0 , a_1 and a_2 were carried out in the same way as that of linear regression model. In the same manner, the error was estimated by determining the difference between the observed and the predicted value and this is expressed as

$$S = \sum (y - a_0 - a_1 x - a_2 x^2)^2 \quad (13)$$

The partial differential of equation 13 with respect to a_0 , a_1 and a_2 yielded the following equations

$$\sum y = a_0 n + a_1 \sum x + a_2 \sum x^2 \quad (14)$$

$$\sum xy = a_0 \sum x + a_1 \sum x^2 + a_2 \sum x^3 \quad (15)$$

$$\sum x^2 y = a_0 \sum x^2 + a_1 \sum x^3 + a_2 \sum x^4 \quad (16)$$

Other least square polynomial models were also adopted for the predictions.

RESULTS AND DISCUSSIONS

Table 1 shows the physiochemical properties of the cassava mill effluent. The pH value of the effluent placed the effluent on the acidic range. All other values were within the range of a typical cassava mill effluent as observed by Oviasogie and Ndiokwere (2008) [19]. Presented in table 2 are the physiochemical values of the CME polluted soil with respect to the sampling pedons and layers.

At different pedons, there is a glaring pH variation. pH values are lowest in pedon B (discharged point DP) but highest in pedon A (Background or control unit). This established the fact that the lowest values of pH observed at pedon B was due to the discharge of the effluent on the units [20]. The pH values in the pedon B (discharged point (DP) appeared to be lower than the value where the activities of micro-organisms responsible for increase in soil ecosystem pH, organic matter and nitrogen were observed to have been rendered inactive by cassava mill effluent. The soil pH seems to determine the level of abundance of many other selected heavy metals and soil properties. Heavy metal values at pedons A where the concentration of CME is low, are the lowest while the most heavy metal concentrations at pedon B except lead (pb) are highest in value, followed by those in pedon C.

Fig 2: Some physiochemical properties of the studied soil for various pedons; pedon A (Background or control unit)

Depth	Cn	Na	Cu	Zn	Cd	pb	Ph	%sand	%silt	Clay	CEC	OC	MC	%PO	BD
0-15	2.71	17.5	3.2	5.14	1.44	0.53	6.5	48	29	23	80	40	121	46	1.11
15-30	2.9	17.8	3.4	5.24	1.4	0.55	6.8	35	26	28	150	31	136	38	1.25
30-70	2.94	18.5	3.43	5.64	1.54	0.59	6.7	30	25	45	163	28	157	39	1.2
70-100	2.98	18.6	3.65	5.2	1.6	0.6	6.9	25	24	46	190	15	178	32	1.47
100-150	3.11	19.24	4.21	6.05	1.9	0.81	7.8	21	20	49	219	10	189	30	1.59
pedon B (DISCHARGE PIONT)															
0-15	21.8	#####	1.9	15.5	3.45	0.39	3.88	16	17	41	114	15	195	12	2.13
15-30	29.1	37.1	2.2	11.7	2.92	0.401	4.93	19	20	48	117	36	189	15	2.04
30-70	29.9	39.3	2.9	10.9	2.49	0.451	5.11	20	24	51	120	21	173	17	1.73
70-100	30.1	40.1	3.7	10.3	2.49	0.47	5.89	18	21	57	161	28	168	20	2.11
100-150	31.09	41.8	3.9	11.3	2.11	0.51	5.87	22	22	60	174	30	150	23	1.34
Pedon C Downstream)															
0-15	11.34	11.7	2.11	10.9	1.43	0.466	4.91	20	20	41	90	43	189	21	1.9
15-30	15.1	15.1	2.31	11.39	0.99	0.711	6.03	18	19	40	110	51	178	18	1.9
30-70	17.01	15.37	2.57	8.73	1.51	0.933	6.49	18	18	45	140	33	170	18	2
70-100	14.43	15.98	2.54	8.32	1.59	0.91	6.86	18	22	46	175	21	170	12	2.01
100-150	18.45	16.43	3.24	9.12	1.78	0.99	6.98	15	15	52	185	16	168	10	2.3

However, heavy metal distribution in all the pedons is represented in this order pedon B> pedon C>pedon A. This trend of distribution is attributed to high concentration presence of CME and its low pH values in pedon A which was equally observed by (Ogboghodo *et al.* 2001 [21], and soil natural attenuation of the effluent for the reduction of heavy metal values observed in Pedon C. The seemingly low pb values observed in pedon B is attributed to the pb mitigating ability of CEM effluent due to abundant complexes of the cassava effluent with pb occasioned by low pH [19]. The physiochemical characterization of wastewater polluted farm land by Onweremadu 2008 [22] is also similar to this observation. The excessively high values of Cd recorded at pedon B (discharge piont(DP) where appreciable amount of CME was received is not necessarily the Cd content of the CME but mainly due to the release of Cd from the soil occasioned by the low acidic medium of the effluent [23]

Low pH of the CME effluent equally has an influence on the other physiochemical properties of the soil. From table2,CEC increased with increase in pH. In Pedon B lowest value of CEC was recorded probably, due to its low acidic medium. This implies that the increased pH of CME polluted soil affected the CEC of the soil, as effective CEC increases with increased pH. This is similar to the observation of [24] where an increase in soil pH due to crude oil pollution resulted to a corresponding increase in its cation exchange capacity (CEC). Also in pedon B low concentration of organic matter is noticed,

Fig. 3: Model equations of ph with the selected soil properties in pedon A

Name	Model equation	R2
ph-pb	$Y = 0.2182x - 0.8982$	0.9574
ph-Cd	$Y = 0.3664x - 0.9668$	0.8691
ph-zn	$Y = 0.6385x + 1.0225$	0.6892
ph-cu	$Y = 0.7583x - 1.6846$	0.9681
ph-Na	$Y = 1.1565x + 10.302$	0.7125
ph-Cn	$Y = 0.2484x + 1.204$	0.741
ph-%sand	$Y = -15.968x + 142.62$	0.5881
ph-%silt	$Y = -6.087x + 67.043$	0.8761
ph-%clay	$Y = -28.738x^2 + 429.33x - 1551.4$	0.5724
ph-CEC	$Y = -170.97x^2 + 2547.7x - 9251.5$	0.9341
ph-OC	$Y = -20.415x + 166.48$	0.7091
ph-MC	$Y = -67.788x^2 + 1021.6x - 3655.1$	0.784
ph-PO	$Y = -10.277x + 108.32$	0.668
ph-BD	$Y = 0.353x - 1.1256$	0.7943

Fig. 4: Model equations of ph with the selected soil properties in pedon B

Name	Model equation	R2
ph-Cn	$y = 4.1653x + 7.0049$	0.8388
ph-Na	$y = 5.6642x + 8.4288$	0.9127
ph-Cu	$y = -0.5767x + 5.6541$	0.8667
ph-Zn	$y = -2.2366x + 23.427$	0.8069
ph-Cd	$y = 1.0035x - 2.2338$	0.8782
h-pb	$y = 0.053x + 0.1723$	0.878
ph-%sand	$y = -2.4618x^2 + 26.393x - 48.474$	0.9265
ph-%PO	$y = 1.418x^2 - 9.0632x + 25.769$	0.9243
ph-%silt	$y = -2.4618x^2 + 26.393x - 48.474$	0.7265
ph-%caly	$y = 8.9108x + 5.634$	0.9618
ph-CEC	$y = 26.434x^2 - 231.38x + 613.85$	0.9648
ph-MC	$y = -6.5816x^2 + 45.884x + 116.54$	0.8035
ph-OC	$y = -0.2075x + 2.9356$	0.8583

Fig. 5: Model equations of pH with the selected soil properties in pedon C

Name	Model equation	R2
ph-Cn	$Y = -0.8075x^2 + 12.259x - 29.367$	0.7097
ph-Na	$Y = 2.185x + 1.2512$	0.9566
ph-Cu	$Y = 0.9271e0.1604x$	0.7029
ph-Zn	$Y = -0.9166x^2 + 9.6307x - 14.175$	0.6802
ph-Cd	$Y = 0.0586x^2 - 0.2214x + 1.48$	0.6316
ph-Pb	$Y = -0.0445x^2 + 0.7667x - 2.2359$	0.9546
ph-%sand	$Y = -1.7338x + 28.643$	0.6591
ph-%silt	$Y = -0.4586x^2 + 4.4887x + 8.922$	0.1018
ph-%clay	$Y = 5.456x^2 - 60.364x + 205.85$	0.8857
ph-CEC	$Y = 29.903x^2 - 309.1x + 886.71$	0.9995
ph-OC	$Y = -19.353x^2 + 216.3x - 552.12$	0.9762
ph-MC	$Y = -10.255x + 239.13$	0.9708
ph-BD	$Y = 0.1745x^2 - 1.9333x + 7.1915$	0.7409

suggesting the possibility of the soil -organic matter-producing micro-organisms being affected by low pH as equally observed by (Oviasogie and Ndiokwere 2008) [19]

Down the soil profile, the pH values increased down the layer in all the pedons, implying that the pH range is highest at the deepest depth range of 100-150cm as displayed on table 2. The observed pH distribution pattern could explain why most heavy metal values increased with depth, further buttressing the abundance of heavy metals with increase in pH as studied by Igbozurike *et al* 2009 [25]. Values of other physiochemical properties of the soil also reacted to the pH variation down the soil profile. CEC values increases down the profile because of high pH value observed down the soil depth. Low concentration of the CEM effluent due to poor infiltration nature of the soil at the top layer, occasioned by the effluent is responsible for the high pH down the profile.

Regression Model: In this study, soil pH determined the fate of other physiochemical properties. In the light of this, regression analysis was carried, establishing the relationship between pH and other soil properties of

the pedosphere. Several models such as linear regression, polynomial, power function and exponential were used to test the data with the view of determining the model that best fit the data. In each case, pH was chosen as the independent variable (X) while other respective physiochemical property was taken as the dependent variable (Y). The choice of using pH as an independent variable was due to the easiness in determining its value in addition to being a function of other soil physiochemical properties. Confronting with similar problems in the same geology, other soil properties in the pedosphere can be evaluated by mere determining the pH value of the soil, saving financial resources and man hour. PH values, the predictor was regressed on individual bases with other physiochemical properties of the soil. Tables 3,4 and 5 represent the regression models in pedons A,B and C respectively, conducted on the predictability of other soil properties that correlate significantly with pH.

In table 3 and 4, pH relationship with all the heavy metals were linear with their linear coefficients (R2) ranging from 0.9-0.6 and 0.9-0.8 for metals in pedons A and B respectively. With other physiochemical properties, pH also varied linearly with %sand, %silt, OC, PO and BD in pedon A, having their coefficients of linearity as follows; 0.57, 0.87, 0.70 and 0.79 respectively while pH relationship with %clay, CEC and MC respectively showed quadratic regression model. Also, virtually all the other physiochemical properties of the soil in pedon B except %clay and OC showed quadratic relationship with high levels of relationship of between 0.73-0.97. There was what seemed to be a reverse trend in pedon C where the impacts of the CME were received in reduced concentrations. In this pedon, almost all the heavy metals except Cu and Na obeyed the quadratic regression model. Their strong quadratic relationships were evident on the high coefficient values (0.60-0.99). Various regression curves are represented in figs 1,2,3, 4 and 5.

Fig. 1: Shows regression curves of pH with selected soil physiochemical proerties in Pedon A

Fig. 2: Regression curves of ph in relation to the heavy metals in pedon B

Fig. 3: Shows the regression curves of ph-the selected soil physiochemical properties in pedon B

Fig. 4: Shows the regression curves of ph with the heavy metals in pedon C

Fig. 5: Shows the regression curves of ph with the selected soil chemical properties at pedon C

Linear relationship of pH with heavy metals showed positive coefficient of linearity in pedons A and B, implying more heavy metal abundance as the acidic range of the soil increases. But the reverse is true for Cu and Na where the distribution these metals are more with low acidic medium.

CONCLUSION

Cassava mill effluent has enormous influence on the soil acidity which on the other hand a function of the distribution of many soil properties on the study site. The influence is high on the pedons which have appreciable amount of the effluent. Linear regression models fitted the pH relationships with heavy metals while polynomials also fitted pH with other physiochemical properties especially in background pedon and the pedon which directly received the effluent. All the heavy metals and other physiochemical properties of the soil gave the highest linear and polynomial prediction respectively of soil pH in cassava- mill- effluent receiving pedon

ACKNOWLEDGMENT

Our profound gratitude goes to erosion unit of federal university of technology owerri for allowing us the use of the chemistry laboratory. The services of the laboratory attendant who was on hand to provide us with the needed materials are highly acknowledged.

REFERENCES

- Oti, E.E., 2002. Acute toxicity of cassava mill effluent to the African catfish fingerlings. *J. Aquatic Sci.*, 17(1): 31-34
- Food and Agricultural Organisation [FAO] 2004. Strategic Environmental assessment. [http:// www.fao.org/docrep](http://www.fao.org/docrep)
- Claude, F. and F. Denis, 1990. African cassava mosaic virus: Etiology, epidemiology and control. *Plant Dis.*, 74(6): 404-411.
- Idu, M. and O.M. Omoruyi, 2003. Effect of seed pre-treatments on germination of two surface types of *Dialium guineense*. *Seed. Technol.*, 25: 41-44.
- Francis O. Arimoro, Chukwujindu M.A. Iwegbue and Benedicta O. Enemudo Effects of cassava effluent on benthic macroinvertebrate assemblages in a tropical stream in southern Nigeria *Acta Zoologica Lituanica*, 2008, Volumen 18, Numerus 2 ISSN 1648-6919. DOI: 10.2478/v10043-008-0021-0
- Zualiya, R. and M.I. Muzondo, 1993. Protein enrichment of cassava by solid state fermentation. *Labensmitted*, 15: 171-174
- Alloway, B.J., 1995. Heavy metals in soils. 2nd. Ed. Chapman and Hall, Glasgow, pp: 34.
- Nabulo, G., Oryem H. Origa, G.W. N asinyama, D. Cole, 2008. Assessment of Zn, Cu, Pb and Ni contamination in wetland soils and plants in the lake basin. *Int. J. Environ. Sci. Tech.*, 5(1): 65-74. (10 pages)
- Okafor, E.C. and K. Opuene, 2007. Preliminary assessment of trace metals and polycyclic aromatic hydrocarbons in the sediments . *Int. J. Environ. Sci. Tech.*, 4(2): 233-240. (8 pages)
- Global positioning system(GPS) 1989. Handheld Global positioning system receiver collection for the year 1989. Imo state environmental issues. A yearly Published Magazine on Environmental Issues. pp: 45-48 .
- Orajaka, S.O., 1975. Geology. In: Nigeria in maps: Eastern States. in: Ofomata, G.E.K. (Ed.). Ethiope Publishers. Benin city Nigeria, pp: 5-7. (3 pages). Orajaka, S.O. 1975. Geology. In: Nigeria in maps: Eastern States. in: Ofomata, G. E. K. (Ed.). Ethiope Publishers. Benin city Nigeria, 5-7 (3 pages).
- FDALR, 1985. The reconnaissance soil survey of Imo state, Nigeria, Federal Department of Agricultural Land Resources (1:250,000), Soil report, pp: 133.
- Gee, G.W. and D. Or, 2002. Particle Size Distribution. in: Dane, J.H. and G.C. Topp, (Eds.). *Methods of soil analysis, Part 4. Physical Methods*, SSSA Book, Malison, WI. pp: 255-293.
- Barabara, F., K. Stephen and W. William, 2002. Speciation and character of heavy metals contaminate soil using computer controlled scan electron microscope. *Environ. Forensics*, 3(2): 131-143. (13 pages).
- Pardo, M.T., 2000. Sorption of lead, copper, zinc and cadmium by soils, effects of nitriloacetic acid on metal retention. *Soil Sci. Plant Anal.*, 31(1-2): 31-40 (10 pages).
- Grossman, R.B. and T.G. Reinsch, 2002. Bulk density and linear extensibility. in: Dane, J.H. and G.C. Topp, (Eds.). *Methods of the soil analysis part 4. Physical methods*. Soil Sci. Soc. Am. Book Series No.5, ASA and SSSA, Madison, WI., pp: 201-228.
- Obi, M.E., 1990. *Soil Physics: A Compendium of lectures*. Department of Soil Science, U.N.N. pp: 103.
- Nwagozie, I.F., 1998. *Probability and statistics for science and engineering practice*. ISBN; 976-074-675-7. University of Port Harcourt press

19. Oviasogie, O.P. and C.L. Ndiokwere, 2008. Fractionation of lead and cadmium in refuse dump soil treated with cassava mill effluent. *J. Agric. Environ.* 9, Jun. 2008. Technical Paper10 DOI: 10.3923/rjes.2007.166.172 URL: <http://scialert.net/abstract/?doi=rjes.2007.166.172>
20. Wang, D.C., Q.R. Y.S. Cui, X.M. Liu, Y.T. Dong and P. Christie, 2003. Soil contamination and plant uptake of heavy metals at polluted sites in China. *J. Envir, Sci. and Health. Part A - Toxic/Hazardous Substances and Environmental Engineering*, 38: 823-838.
21. Ogboghodo, I.A., I.O. Osemwota, S.O. Eke and A.E. Iribhogbe, 2001. Effect of cassava (*Manihot esculenta crantz*) mill grating effluent on the textural, chemical and biological properties of surrounding soils. *World J. Biotechnol.*, 2: 292-301.
22. Onweremadu, E.U., 2008. Physico-chemical charaterisation of a farmland affected by wastewater in relation to heavy metals. *J. Zhejiang U. Sci. A.*, 9(3): 366-372. (7 pages).
23. Bolan, N.S. and V.P. Duraisamy, 2003. Role of inorganic and organic soil amendments on immobilisation and phytoavailability of heavy metals: a review involving specific case studies. *Australian J. Soil Res.*, 41: 533-555.
24. Singer, M.J. and D.N. Munns, 1999. *Soils, an introduction*. 4th.Ed. Prentice-Hall, Inc. New Jersey.
25. Igbozurike, C.W.A., A.O. Opara-Nadi and I.K. Okorie, 2009. Concentrations of Heavy metals in soil and cassava plant on sewage sludge dump; UC Davis, the Proceedings of the International Plant Nutrition