

## UV Shielding Properties of Jarosite Vs. Gypsum: Astrobiological Implications for Mars

<sup>1</sup>G. Amaral, <sup>2</sup>J. Martinez-Frias and <sup>2,3</sup>L. Vázquez

<sup>1</sup>Departamento de Química Física I, Fac. Ciencias Químicas,  
Universidad Complutense, 28040 Madrid, Spain

<sup>2</sup>Centro de Astrobiología (CSIC-INTA), Associated to the NASA Astrobiology Institute,  
Ctra de Ajalvir km. 4, 28850 Torrejón de Ardoz, Spain

<sup>3</sup>Departamento de Matemática Aplicada, Facultad de Informática,  
Universidad Complutense de Madrid, 28040 Madrid, Spain

---

**Abstract:** The discovery of liquid water-related sulfates on Mars is of great astrobiological interest. UV radiation experiments, using natural Ca and Fe sulfates (gypsum, jarosite), demonstrate a large difference in the UV protection capabilities of both minerals and also confirm that the mineralogical composition of the Martian regolith is a crucial shielding factor. It is demonstrated that a thin (500  $\mu\text{m}$ -thick) sample of jarosite prevents UV transmission, providing suitable niches for life exploration.

**Key words:** UV shielding % Mars % jarosite % gypsum % astrobiology

---

### INTRODUCTION

Unlike Earth, there is a significant amount of UV flux on Mars, mainly due to the influence of the shorter wavelengths UVC (100-280 nm) and UVB (280-315nm). Various works on the biological effects of UV radiation [1-3] and the amount of Martian radiation [4] have established that even the present-day instantaneous Martian UV flux would not in itself prevent life. Nevertheless, it is a fact that this UV flux contributes, coupled with the lack of liquid water and extreme low temperatures, to the biologically inhospitable nature of the Martian surface. From the astrobiological point of view, these factors render a practical consequence for the exploration and detection of life on Mars: any living organism, as we know it, should have preferentially developed in a particular sub-surface microenvironment able to protect it from the harsh conditions on the surface. Terrestrial endolithic communities that live in the subsurface layers of rock that provide appropriate microenvironments against extreme external conditions have been proposed [5-8] as possible analogs to life on Mars. Extant Martian life would require strong UV shielding, which, in accordance with our experiments, could be perfectly accomplished by certain minerals already discovered on Mars.

### MARTIAN MINERALS: CA AND FE SULFATES

The search for life on Mars has been intimately linked to the identification of unequivocal ancient or modern geomarkers [9, 10] of water (e.g. some water-related minerals) on and inside the planet. Recently, some water-related sulfates (e.g. jarosite, gypsum) were discovered [11-13] on Mars surface. In particular, jarosite has proven to have a great astrobiological importance, not only for its relation with liquid water, but also because it can act as a sink and source of Fe ions for Fe-related chemolithoautotrophic microorganisms, such as those encountered in numerous extremophilic ecosystems (e.g. Tinto river [14-16]). In the present report, we present experimental results on the UV shielding capabilities of the above mentioned minerals: jarosite ( $\text{KFe}_3(\text{SO}_4)_2(\text{OH})_6$ ) and gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ).

The samples of jarosite and gypsum used in this work come from two selected areas of the SE Mediterranean region of Spain: Jaroso and Sorbas, which have been proposed as a relevant geodynamic and mineralogical model [17-21] to follow for the astrobiological exploration of Mars. The Jaroso Hydrothermal System is a volcanism-related multistage hydrothermal episode of Upper Miocene age, which includes oxy-hydroxides (e.g. hematite), base-and




Fig. 1: Typical jarosite pellet. Jarosite sample from El Jaroso ravine, Cuevas del Almanzora Natural Area, Almería province, SE Spain. Each division on the left scale represents 1 mm




Fig. 2: Light transmission spectrum (expressed as % transmitted light) of jarosite vs. gypsum samples

precious-metal sulfides and different types of sulphosalts. Hydrothermal sulfuric acid weathering of the ores has generated huge amounts of oxide and sulfate minerals [19, 22-24] of which jarosite is the most abundant. It is important to note that jarosite was first discovered on Earth at this area, in 1852, in the "Jaroso Ravine", which is the world type locality of jarosite [25, 26]. The Sorbas basin contains one of the most complete sedimentary successions of the Mediterranean (gypsum karst) reflecting the increasing salinity during the Messinian salinity crisis (desiccation of the Mediterranean Sea) [27-29] and showing a complex paleogeographical evolution, being a signature of its progressive restriction and isolation.

## MATERIALS AND METHODS

Jarosite samples were pulverized and shaped by pressure into flat round pellets of 1 cm diameter and 0.5 mm thickness (Fig. 1). Gypsum samples were scratched from a larger specimen. The samples were flattened to different thickness (between 0.1 mm to 1.6 mm) before exposed to UV light. All samples were positioned onto a sample holder which allowed the UV lamp light to go through the samples and be detected by a PMT through a monochromator. The UV light source is a Xe Lamp (SpectralProducts) with an integrated output from 220 nm to 500 nm of  $1.2 \text{ Wm}^{-2}$  (measured at a distance of 10 cm from the lamp exit and over a disc of 5 mm diameter). A monochromator (SpectralProducts, 1/8 m CM 110) is placed at the output of the lamp, but in our experiments it is set up so that it lets all wavelengths out without deformation of the Xe lamp output. Five (5) cm downstream the light beam irradiates the sample, the light which is transmitted through it is then collected by an optical fiber head (placed 10 cm away from the Xe lamp), which in turns directs the light through a spectrometer (Bentham, CM 150 double monochromator) which records the transmitted light spectrum. The light is detected by a peltier-cooled bi-alkali photomultiplier tube (PMT Bentham, DH-10-Te). Typically, each sample spectrum is measured from 220 nm to 500 nm every 1nm with a rate of approximately 1 nm/s. Usually each sample spectrum is the average of at least 5 spectra.

## CONCLUSIONS

It is well known that iron and iron-bearing compounds can provide an UV screen for life. For instance, Martian regolithic dust, along with palagonite-type compositions, have specifically been suggested as

a possible safe haven for life [1-3, 30-35]. Our experimental results demonstrate a large difference in the UV shielding capabilities of both minerals (Fig. 2). Whereas gypsum showed a much higher transmission percentage (with an error in the absorption coefficients of roughly 20%), jarosite samples, with a thickness of only 500  $\mu\text{m}$ , prevented transmission. Our mineralogical results complement other recent petrologic studies about shielding effects on Mars surface environment (e.g. basaltic dust, [36]) and have great astrobiological relevance as: a) jarosite typically occurs on Earth as alteration crusts and patinas and b) a very thin crust of jarosite on the surface of Mars would be sufficient to shield microorganisms from UV radiation.

## ACKNOWLEDGEMENTS

The authors express thanks for support from the Rover Environmental Monitoring Station project, associated with the Mars Science Laboratory-NASA mission to Mars and supported by the Centro de Astrobiología. Maite Fernandez Sampedro, Maria Paz Martín Redondo and Dr. Virginia Souza-Egipsy are acknowledged for their assistance with the analyses. Also, the authors acknowledge ArXiv database (Cornell University) for its support hosting an electronic draft version of the work. Special thanks to Dr. David Hochberg for the revision of the English version.

## REFERENCES

1. Cockell, C.S., 1998. The biological effects of high ultraviolet radiation on early Earth-A theoretical evaluation. *J. Theor. Biol.*, 193: 717-729.
2. Cockell, C.S., D. Catling, W.L. Davis, R.N. Kepner, P.C. Lee, K. Snook and C.P. McKay, 2000. The ultraviolet environment of Mars: Biological implications past, present and future. *Icarus*, 146: 343-359.
3. Gomez, F., A. Grau, L. Vazquez and R. Amils, 2003. UV radiation effects over microorganisms and study of protective agents. *Proc. of the III European Workshop on Exo-Astrobiology, Mars, The Search for Life, Madrid, Spain, 18-20 November 2003*, pp: 21-26.
4. Zorzano, M.P., A.M. Mancho and L. Vázquez, 2005. Numerical Integration of the Discrete-Ordinate Radiative Transfer Equation in Strongly Non-Homogeneous Media" *Applied Mathematics and Computation*, 164: 263-274.
5. Friedmann, E.I., 1982. Endolithic microorganisms in the antarctic cold desert. *Science*, 215: 1045-1053.

6. McKay, C.P., 1993. Relevance of Antarctic microbial ecosystems to exobiology. In Antarctic Microbiology, Wiley-Liss, New York, Friedmann E.I., (Ed.). pp: 593-601.
7. Wynn-Williams, D.D. and H.G.M. Edwards, 2000. Laser Raman microspectroscopy of surface microbial communities and protective biomolecules in situ: Overview of terrestrial Antarctic habitats and Mars analogs. *Icarus*, 144: 486-503.
8. Villar, S.E., H.G. Edwards and C.S. Cockell, 2005. Raman spectroscopy of endoliths from Antarctic cold desert environments. *Analyst.*, 130-132: 156-162.
9. Martinez-Frias, J., G. Amaral and L. Vázquez, 2006. Astrobiological significance of minerals on Mars surface environment. *Reviews in Environmental Science and Bio/technology*, 5: 219-231.
10. Martinez-Frias, J., E. Lázaro and A. Esteve, 2007. Geomarkers versus Biomarkers: Paleoenvironmental and astrobiological significance. *AMBIO: J. Human Environ.*, 36-4, (In Press).
11. Squyres, S.W., J.P. Grotzinger, R.E. Arvidson, J.F. Bell III, W. Calvin, P.R. Christensen, B.C. Clark, J.A. Crisp, W.H. Farrand, K.E. Herkenhoff, J.R. Johnson, G. Klingelhöfer, A.H. Knoll, S.M. McLennan, H.Y. McSween, R.V. Morris, Jr., J.W. Rice, R. Rieder and L.A. Soderblom, 2004. In situ evidence for an ancient aqueous environment at Meridiani Planum, Mars, *Science*, 306: 1709-1714.
12. Klingelhöfer, G., R.V. Morris, B. Bernhardt, C. Schröder, D.S.J.R. Rodionov, P.A. de Souza, A. Yen, R. Gellert, E.N. Evlanov, B. Zubkov, J. Foh, U. Bonnes, E. Kankeleit, P. Gülich, D.W. Ming, F. Renz, T. Wdowiak, S.W. Squyres and R.E. Arvidson, 2004. Jarosite and Hematite at Meridiani Planum from Opportunity's Mössbauer Spectrometer Science, 306-5702: 1740-1745.
13. Langevin, Y., F. Poulet, J.P. Bibring and B. Gondet, 2005. Sulfates in the North Polar Region of Mars Detected by OMEGA/Mars Express, *Science* 307-5715, 1584-1586.
14. López-Archilla, A.I., I. Marín and R. Amils, 2001. Microbial Community Composition and Ecology of an Acidic Aquatic Environment: The Tinto River, Spain. *Microb. Ecol.*, 41: 20-35.
15. Gonzalez-Toril, E., E. Llobet-Brossa, E.O. Casamayor, R. Amann and R. Amils, 2003. Microbial Ecology of an Extreme Acidic Environment, the Tinto River. *Appl. Environ. Microbiol.*, 69: 4853-4865.
16. Amaral Zettler, L.A., M.A. Messerli, A.D. Laatsch, P.J.S. Smith and M.L. Sogin, 2003 From Genes to Genomes: Beyond Biodiversity in Spain's Rio Tinto *Biol. Bull.*, 204: 205-209.
17. Martínez-Frías, J., R. Lunar and J.A. Rodríguez Losada, 2001. Hydrothermal mineralization in SE Spain as possible volcanics-related metallogenetic model for the early Mars, 2nd Astrobiology Minisymposium, CAB (CSIC/INTA), associated to NASA Astrobiology Institute, Madrid, Spain (abstract).
18. Martínez-Frías, J., R. Lunar, J. Mangas, A. Delgado, G. Barragan, E. Sanz-Rubio, E. Diaz, R. Benito and T. Boyd, 2001. Evaporitic and hydrothermal gypsum from SE iberia: Geology, geochemistry and implications for searching for life on Mars, Geological Society of America (GSA) Annual Meeting, Boston, Massachussets, November 5-8 (Abstract).
19. Martínez-Frías, J., R. Lunar, J.A. Rodríguez-Losada and A. Delgado, 2004. The volcanism-related multistage hydrothermal system of El Jaroso (SE Spain): Implications for the exploration of Mars. *Earth, Planets Space* 56: 5-8.
20. Rull, F., J. Martinez-Frias and J. Medina, 2005. Surface mineral analysis from two possible Martian analogs (Rio Tinto and Jaroso Ravine, Spain) using micro, macro-and remote laser Raman spectroscopy. *European Geosciences Union, Geophysical Research Abstracts*, Vol. 7, 09114.
21. Martinez-Frias, J., A. Delgado-Huertas, F. García-Moreno, E. Reyes, R. Lunar and F. Rull, 2006. Oxygen and carbon isotopic signatures of extinct low-temperature hydrothermal chimneys in the Jaroso Mars analog. *Planetary and Space Science* (In Press).
22. Frost, R.L., M.L. Weier, J.T. Klopogge, F. Rull and J. Martinez-Frias, 2005. Raman spectroscopy of halotrichite from Jaroso, Spain *Spectrochimica Acta Part A Molecular and Biomolecular Spectroscopy*, 62: 1-3: 166-180.
23. Frost, R.L., D.L. Wain, B.J. Reddy, W. Martens, J. Martinez-Frias and F. Rull, 2006. Sulphate efflorescent minerals from the El Jaroso ravine, Sierra Almagrera, Spain-a scanning electron microscopic and infrared spectroscopic study. *J. Near Infrared Spectroscopy*, 14: 167-178.

24. Frost, R.L., M. Weiera, J. Martinez-Frias, F. Rull and B. Jagannadha Reddy, 2007. Sulphate efflorescent minerals from El Jaroso Ravine, Sierra Almagrera - An SEM and Raman spectroscopic study, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 66: 177-183.
25. Amar de la Torre, R., 1852. Descripción de los minerales, algunos de ellos nuevos, que constituyen el filón del Barranco Jaroso de Sierra Almagrera, por el caballero profesor el doctor Augusto Breithaupt, de Freiberg, *Revista Minera*, 3: 745-754.
26. Martinez-Frias, J., 1999. Mining vs. Geological Heritage: The Cuevas del Almanzora Natural Area (SE Spain), *AMBIO: A Journal of the Human Environment*, 28-2: 204-207.
27. Martin, J.M. and J.C. Braga, 1994. Messinian events in the Sorbas Basin in southeastern Spain and their implications in the recent history of the Mediterranean. *Sedimentary Geol.*, 90: 257-268.
28. Ridig R., J.C. Braga, H.M. Martin and I.M. Sánchez-Almazo, 1998. Mediterranean Messinian salinity crisis: Constraints from a coeval marginal basin, Sorbas, Southeastern Spain, *Marine Geology*, 146: 1-20.
29. Krijgsman, W., H.J. Hilgen, I. Raffi, F.J. Sierro and D.S. Wilson, 1999. Chronology, causes and progression of the Messinian salinity crisis. *Nature*, London, 400: 652-655.
30. Olsen, J.M. and B.K. Pierson, 1986. Photosynthesis 3.5 thousand million years ago. *Photosynthesis Res.*, 9: 251-259.
31. Pierson, B.K., H.K. Mitchell and A.L. Ruff-Roberts, 1993. Chloroflexus aurantiacus and ultraviolet radiation: Implications for archean shallow-water stromatolites. *Origins Life Evol. Biosphere*, 23: 243-260.
32. Kumar, A., M.B. Tyagi, G. Srinivas, N. Singh, H.D. Kumar, R.P. Sinha and D.-P. Haeder, 1996. UVB shielding and the role of FeCl<sub>3</sub> and certain cyanobacterial pigments. *Photochem. Photobiol.*, 63: 321-325.
33. Sagan, C. and J.B. Pollack, 1974. Differential transmission of sunlight on Mars: Biological implications. *Icarus* 21: 490-495.
34. Allen, C.C., K.M. Jager, R.V. Morris, D.J. Lindstrom, M.M. Lindstrom and J.P. Lockwood, 1998. Martian Soil Simulant Available for Scientific, Educational Study. *EOS*, 79: 405.
35. Phoenix, V.R., K.O. Konhauser, D.G. Adams and S.H. Bottrell, 2001. The role of biomineralization as an ultraviolet shield: Implications for the Archean. *Geol.*, 29: 823-826.
36. Muñoz-Caro, G.M., E. Mateo-Martí and J. Martínez-Frías, 2006. Near-UV transmittance of basalt dust as an analog of the Martian regolith: Implications for sensor calibration and astrobiology. *Sensors*, 6: 688-696.