

Study on Seed Germination and Seedlings Growth of Bamboo (*Dendrocalamus hamiltonii*)

¹Tooba Abedi and ²Firouzeh Herfeh doost

¹Member of Environmental Research Institute of Academic Centre for Education,
Culture and Research (ACECR), Rasht, Guilan, Iran

²Graduate student of Department of Forestry,
Faculty of Natural Resources, University of Guilan, Someesara

Submitted: Aug 14, 2013; **Accepted:** Sep 21, 2013; **Published:** Sep 24, 2013

Abstract: Bamboos are naturally propagate both sexually and asexually from seeds and rhizomes. In this study two experiments were carried out in a greenhouse to evaluate the influence of seed orientation (embryo-end-up, lay-flat and embryo-end-down) and sowing depth (3, 6 and 9 mm) of the bamboo *Dendrocalamus hamiltonii* on seed germination, seedling survival and growth after 60 days. Factorial design under two blocks containing 90 samples were used. Plastic pots filled with a mixture of sand, agricultural soil and peat (rate 3:1:1 and 2:2:1, respectively). Seeds were sown in two soil treatments. Seed germination and seedling survival rates in first soil (3:1:1) were 24 and 66.6 percent and in second soil treatment (2:2:1) were 31.1 and 85.7 percent, respectively. Mean seedlings height was higher at 6 mm depths and embryo-end-down in both soil treatments (11.5 and 27.5), it had significant difference in two soil treatments ($P < 0.05$). Mean seedlings height didn't show significant difference among seed orientation and sowing depth.

Key words: Bamboo • Seed germination • Seedling growth • *Dendrocalamus hamiltonii*

INTRODUCTION

Bamboos are tall perennial grasses with tree stature that grow up to about 30 m in height and 35 cm in diameter. They belong to the Poaceae (Gramineae) family and Bambuseae subfamily [1]. The main stem of the aboveground part of the plant is the culm, while the underground part constitutes the rhizome and root system. Most bamboo plants flower only once in their lifetime (14 to 50 years in most species) and then die soon after. They emerge again from germinating seeds if the site is not severely.

Disturbed by detrimental factors such as rodents, fire, etc. These phenomena were actually observed in bamboo forests [2]. Bamboo naturally propagates both sexually and asexually from seeds and rhizomes. Artificial propagation by vegetative methods includes planting of off-sets, culm cuttings, layering and grafting of rhizome [3]. Most bamboos are found on sandy loam to loamy clay soils, derived from river alluvium or frequently from

the underlying rock [4]. *Dendrocalamus* species, after germination, produce a grass-like seedling the first year. The plumule, which appears as a conical bud covered by sheathing scaly leaves, develops rapidly into a thin wiry stem bearing single leaves, alternate at the nodes, the leaf bases covering the stem [4]. *Dendrocalamus hamiltonii* Nees et Arn. ex Munro is a large caespitose bamboo, culms 10-20 m high, 10-16 cm in diameter, thin walled with 0.75-1.25 cm thickness [5], internodes 30-50 cm, sheaths stiff and persistent [5]. Its seeds are like those of wheat, seed year happen in intervals of 30 years and flowering nature is Gregarious and sporadic [6].

Aim of this study is finding out the optimum sowing depth and seed orientation that could be recommended for successful seedling production and growth of the *Dendrocalamus hamiltonii* in greenhouse condition.

Pittman (1965) records enhanced germination and early growth for corn seeds oriented with respect to magnetic lines of force [7].

Patten and Van-Doren (1970) found substantially higher emergence and seedling growth of corn with embryo end-up than with embryo-end-down orientation [8].

Maun and Riach (1981) studied sand deposition effects on seedling emergence of *Calamovilfa longifolia* at field sites and greenhouse plantings. They found that seedling emergence was negatively correlated with planting depth [9].

Sanchez and King (1994) identify sowing depth as one of the most important factors that affect seedling emergence, survival and subsequent onward growth of acacia species from Ethiopia [10].

Chen and Maun (1999) sorted seeds of *Cirsium pitcheri* into three groups (small, medium and large) and buried at 2, 4, 6, 8, 10 and 12 cm depths in plastic pots filled with unsterilized sand. Data showed that percent seed germination and emergence of seedlings were not related to seed size. However, both variables were negatively correlated with depth. Seedling emergence occurred from a maximum depth of 6 cm with most seedlings emerging from 2 cm depth [11].

Embaye and *et.al* (2003) used A randomized complete block design to evaluate seedling emergence, subsequent survival and growth of bamboo (*Oxytenanthera abyssinica*). Seeds were sown in plastic pots that filled with mixture of sand and peat (rate 3 sand: 1peat). Top of the soil surface and at 2.5 mm depth achieved faster and higher seedling emergence than those sown at 5 and 10 mm depths. However, mean seedling height and number of leaves per seedling were higher in 5 and 2.5 mm depths than surface and 10 mm depths. There were significant quadratic relationships between sowing depth and seedling height ($p=0.034$) as well as number of leaves per seedling ($p=0.032$), both peaking around 5 mm soil depth. Lay-flat orientation, which was the most frequent position in broadcast sowing, was recommended at 5 mm sowing depth for the lowland bamboo based on overall performance in seedling emergence, survival and growth [3].

Damizade (2004) planted seeds of *Capparis deciduas* in nursery for evaluating germination and survival rates of seedlings [12].

Zhao and *et.al* (2006) sorted seeds of *Nitraria sphaerocarpa* into three size-classes and buried at 2 cm depths in plastic pots filled with sand under controlled greenhouse condition. Two weeks after seedling emergence, seedlings were buried in sand to various depths of 0, 33, 67, 100 and 133% of their mean height.

Seedling height, mass and absolute height growth rate in partial burial treatments were higher than those of the unburied and completely buried treatments. In each seed size class, with increasing burial depth, or in each burial depth, with decreasing seed size, there was a tendency that both biomass allocation to root and biomass allocation to belowground stem increased, while biomass allocation to aboveground stem decreased [13].

Tabari and Tabandeh (2007) investigated germination rate of *Tilia platyphyllos* at 1.5 and 3 cm of sowing depth and watering was done with two regulations: everyday and every other day. Results showed treatment composition of 3 cm sowing depth and everyday watering was the most germination rate than 1.5 cm sowing depth with every other day watering [14].

Zhang and Maun (2007) examine the effects of sand burial on seed germination, seedling emergence and establishment of *Panicum virgatum* L. on the foredunes of Lake Erie. In the field, all the seedlings established in one growing season had originally emerged from sand burial depths of less than 12 cm. Within this burial range, seedlings from shallower burial depths had lower chances of establishment than expected, whereas those from deeper burial depths had higher probabilities of establishment than expected [15].

Azmy (1998) planted *Gigantochloa ligulata* at FRIM and results showed the seedlings developed in the second week after sowing. With 1:3 soil-sand rate and the germination was 76% [16].

MATERIALS AND METHODS

This research is composed of literature review work and greenhouse experiment. Seeds were collected from Indian bamboo stands. Three experiments including soil, sowing depth and seed orientation were implemented. The experiments period was 60 days. Factorial design was used in the experiments. Plastic pots filled with a mixture of sand, agricultural soil and peat (rate 3:1:1 and 2:2:1 sand, agricultural soil and peat, respectively). There were two blocks that each of them consisted of 45 pots. Sowing depth experiment was determined based on average size of the seeds and consisted of one, two and three times of seed size *i.e.* tree treatments as 3, 6 and 9 mm at soil depth, respectively. It tried to select uniform size of seeds. The seeds were soaked with clean water for 12 hours and were sown in embryo-end-up, lay flat and embryo-end-down orientations at different soil depths.

The amount of watering was adequate moisture to maintain in the soil to avoid seed desiccation while keeping it below field capacity to ensure adequate oxygen supply. The treatments were done through the greenhouse at 27 °C temperature and 80% relative humidity. Seedlings heights were measured every 20 days.

RESULTS AND DISCUSSION

First seed germination emerged 8 days after sowing and continued 8-14 days. Germination period of the species is expressed 8-28 days [6]. The period for some species of *Dendrocalamus* is 2-3 weeks [17] and about *Oxytenanthera abyssinica* is reported 14 days after sowing [3].

Germination and survival rates in different soils are showed in table 1.

Seed germination rate in soil 1 based on sowing depth after 60 days was in the order 9mm>6mm>3mm and survival rate followed 6mm>9mm>3mm and in soil 2 the amounts were 6mm>9mm>3mm and 3mm>6mm>9mm respectively. In both of soils the most survival rates were related to shallow sowing depth and medium surface. Chen and Maun (1999) and Zhang and Maun (2007) approved the fact about other species [11, 15], it maybe happen because of access to light quickly [7] and water maintenance around the seeds [11]. Differing results are documented on seedling growth in shallow sowing depth and significant difference between germination and survival of seedlings [9-11, 13]. Seeds of *T. platyphyllos* were germinated in high deep soil better than shallow depth; the reason is probably for seed dormancy and physiological characteristics of seeds (e.g. seed size) [14]. Seed treatment with water showed significant difference in seedling survival rate of *C. deciduas* [12]; we used the treatment in this study. Although, changing soil rates in different treatments were illustrated significant difference of survival rates about the species [12].

Seed germination rate in soil 1 based on seed orientation after 60 days was 60% consisted of embryo-end-down at the 6mm sowing depth and its survival rate was 100%.

Table 1: Germination and survival rates of two soil treatments

Soil treatment	Seed Germination rate(%)	Seedling survival rate (%)
3:1:1 (soil 1)	24.4	66.6
2:2:1(Soil 2)	31.1	85.7




Fig. 1: Germination rate in 3mm sowing depth




Fig. 2: Germination rate in 6mm sowing depth




Fig. 3: Germination rate in 9mm sowing depth




Fig. 4: Survival rate in 3mm sowing depth




Fig. 5: Survival rate in 6mm sowing depth




Fig. 6: Survival rate in 9mm sowing depth

Table 2: Relationship of soil, sowing depth and seed orientation with seedling height 20 days after sowing

Average seedling height									
Soil	3 mm Sowing depth			6 mm Sowing depth			9 mm Sowing depth		
	Embryo-end-up	Lay flat	Embryo-end-down	Embryo-end-up	Lay flat	Embryo-end-down	Embryo-end-up	Lay flat	Embryo-end-down
Soil 1	0 ^a	0 ^a	1.4 ^a	3 ^a	0 ^a	3.43 ^a	2.45 ^a	2 ^a	0 ^a
Soil 2	0 ^a	0 ^a	4.77 ^a	1 ^a	8.88 ^a	10 ^a	2.5 ^a	7.5 ^a	0 ^a

Table 3: Relationship of soil, sowing depth and seed orientation with seedling height 40 days after sowing

Average seedling height									
Soil	3 mm Sowing depth			6 mm Sowing depth			9 mm Sowing depth		
	Embryo-end-up	Lay flat	Embryo-end-down	Embryo-end-up	Lay flat	Embryo-end-down	Embryo-end-up	Lay flat	Embryo-end-down
Soil 1	0 ^a	0 ^a	2 ^a	0 ^a	0 ^a	6.67 ^a	4.5 ^a	0 ^a	2.25 ^a
Soil 2	0 ^a	0 ^a	12 ^a	0 ^a	13 ^a	17.75 ^a	0 ^a	10.5 ^a	0 ^a

Table 4: Relationship of soil, sowing depth and seed orientation with seedling height 60 days after sowing

Average seedling height									
Soil	3 mm Sowing depth			6 mm Sowing depth			9 mm Sowing depth		
	Embryo-end-up	Lay flat	Embryo-end-down	Embryo-end-up	Lay flat	Embryo-end-down	Embryo-end-up	Lay flat	Embryo-end-down
Soil 1	0 ^a	0 ^a	10 ^a	0 ^a	0 ^a	11.5 ^a	9.5 ^a	0 ^a	5.5 ^a
Soil 2	0 ^a	0 ^a	10.17 ^a	0 ^a	20.13 ^a	27.5 ^a	0 ^a	15.75 ^a	5 ^a

^a Data bearing same letters in a column do not differ significantly from each other at $P < 0.05$.

Seed germination rate in soil 2 based on seed orientation after 60 days was 80% consisted of lay flat at the 6mm sowing depth and its survival rate was 100%.

The most survival rate about *O. abyssinica* was explained in lay flat orientation [3]. Growth rate is influenced by soil type [16].

Germination and survival rates at different sowing depth and both soils are shown in Figures 1-6.

The most seedling height was 11.5 cm at age 60 days in embryo-end-down orientation and 6mm sowing depth in soil 1 (Table 4), highest seedling in soil 2 was 27.5 cm at the same orientation and sowing depth (Table 4).

The most seedling height was reported at lay flat orientation for *O. abyssinica* [3] and embryo-end-up for corn [8].

The SPSS statistical software package was used to analysis the sets of data. Block means of the different treatments were applied for 3-WAY ANOVA calculations. Seedlings height between two soils were statistically significant differences until 40 days ($P < 0.05$). After that the seedlings height growth were independent on soil type. Seed orientations and sowing depths didn't have significant differences on seedlings height ($P < 0.05$) (Tables 2-4).

REFRECES

1. Ohrnberger, D., 1999. The Bamboos of the World: Annotated nomenclature and literature of the species and the higher and lower taxa. Elsevier, Amsterdam, pp: 585.
2. Liese, W., 1985. Bamboos-Biology, Silvics, Properties, Utilization. Eschborn, GTZ.
3. Embaye, K., 2003. Ecological aspects and resource management of bamboo forests in Ethiopia. Doctoral thesis, Department of Short Rotation Forestry Uppsala, Swedish University of Agricultural Sciences, pp: 25.
4. Huberman, M.A., 1956. Bamboo silviculture. Unasylva, 13: 1.
5. Rawat, J.K. and D.C. Khanduri, 2007. The status of bamboo and rattan in india. INBAR, pp: 31.
6. Ahlawat, S.P., K. Haridasan and S.N. Hegde, 2002. Field Manual for Propagation of bamboo in North East India. State Forest Research Institute, Department of Environment & Forests, Government of Arunachal Pradesh. Itanagar. SFRI Information Bulletin, 14: 1-18.
7. Pittman, U.J., 1965. Magnetism and plant growth III Effect on germination and early growth of corn and beans. Canadian Journal of Plant Science, 45(6): 549-555.
8. Patten, G.P., D.M. and Van-Doren, 1970. Effect of seed orientation on emergence and growth of corn. Agronomy Journal, 62(5): 592-595.
9. Maun, M.A. and S. Riach, 1981. Morphology of caryopses, seedlings and seedling emergence of the grass Calamovilfa longifolia from various depths in sand. Oecologia, 49: 137-142.
10. Sanchez, B.F. and G.W. King, 1994. Ecological factors affecting the early development of seedlings of three acacia species from Ethiopia. South African Journal of Plant & Soil, 11(3): 147-148.
11. Chen, H. and M.A. Maun, 1999. Effects of sand burial depth on seed germination and seedling emergence of Cirsium pitcheri. Plant Ecology, 140(1): 53-60.
12. Damizade, G.H., 2004. Effect of environmental condition on seedlings survival rate of Capparis decidua (Forssk.) Edgew. Poplar and Forest Research of Iran, 12 (4): 509-531. (In Persian, English summary).
13. Zhao, W.Z., Q.Y. LI and H.Y. FANG, 2006. Effects of sand burial disturbance on seedling growth of Nitraria sphaerocarpa. Plant and soil, 295(1-2): 95-102.
14. Tabari, M. and A. Tabandeh, 2007. Reaction of stratified seed germination of Tilia platyphyllos Scop on watering and sowing depth. Poplar and Forest Research of Iran, (In Persian, English summary), 15: 144-151.
15. Zhang, J. and M.A. Maun, 2007. Sand burial effects on seed germination, seedling emergence and establishment of Panicum virgatum. Ecography, 13(1): 56-61.
16. Azmy, H.J.M. and S. Appanah, 1998. Bamboo resources conservation and utilization in Malaysia. FRIM, Kepong, Kuala Lumpur, Malaysia. In: Rao A.N., V. Ramanatha Rao (editors): The work on Bamboo and Rattan Genetic Resources is supported by Japanese Government. Proceedings of training course cum workshop, 10-17 May 1998, Kunming and Xishuanbanna, Yunnan, China.
17. www.Gator-Ventures.com