

Liver Histopathological Changes in Milkfish (*Chanoschanos*) Exposed to Petroleum Hydrocarbon Exposure

¹Narges Amrollahi Biuki, ¹Ahmad Savari, ²Mohammad Sedigh Mortazavi,
¹Hossein Zolgharnein and ¹Negin Salamat

¹Department of Marine Biology, Faculty of Marine Science,
Marine Science and Technology University, Khorramshahr, Iran
²Persian Gulf and Oman Sea Ecology Research Center, Bandarabbas, Iran

Abstract: Oil pollution is a significant hazard for the marine environment. There is an increasing risk of a major pollution spill in the Persian Gulf waters because of heavy transport of crude oil, petrol and its derivations. The aim of this study was to investigate the effect of exposure petroleum hydrocarbons on the milkfish liver at different concentration and different time points. After determination of lethal concentration of total petroleum hydrocarbons (Lc₅₀-96h), collected specimens of *Chanoschanos* were exposed to different concentrations of petroleum hydrocarbons. The main histological alternation observed in this study include: sinusoid dilation, vacuolation, lipidiosis, necrosis, hepatocytic nuclear pleomorphism and bile stagnation. Our result showed that, the intensity of these histological changes was influenced by the extent of the exposure period and concentration of pollution.

Key words: Petroleum Hydrocarbons • Pollution • Sublethal • LC₅₀ • Histology

INTRODUCTION

In recent years, petroleum products pollution is one of the global environmental issues in aquatic ecosystems. The assessing and prediction of the effects of petroleum pollution on aquatic environment are very urgent and important issue.

Petroleum hydrocarbons and its products can enter the aquatic environment via different sources such as petroleum extraction, transportation, urban runoff, biogenic hydrocarbons produced naturally, accidents involving and oil spills [1]. Petroleum hydrocarbons are lipophilic in nature and can be extremely uptake by a wide spectrum of marine animals and accumulate in lipid compartments like cellular membrane [2, 3], disturbed the physiological and physiochemical membrane properties and can be integrated into biological system [4]. Many physiological processes such as enzyme function, muscle contraction and osmoregulation are directly dependent on the unique properties of biological membranes [2]. Some petroleum compounds are potential to be biomagnified through the food chain [2].

Petroleum pollutants tend to accumulate more in the organism than the environment, therefore fish are largely being used for the assessment of the quality of aquatic environment and can be used as bioindicator to evaluate the environmental contamination levels [5-7].

Histological biomarkers were used for assessing the effect of petroleum hydrocarbons in aquatic organisms [8]. Histological study is a rapid method for detection of pollutants effects on various tissues of fish and it has been extensively used to determine the deleterious effects of hydrocarbons [8-10].

Nowadays, it is accepted that biomarkers are now useful tools in monitoring programs for the assessment of the impact of pollutants on marine organisms and ecological health [11-13]. Instead of measurement of accumulated pollutants in tissues, biomarkers can determine a more complete and more ecologically relevant information about the potential impact of toxic pollutants on the living organisms and more important, allow the early detection of pollutants existence in ecosystems [14, 15].

Exposure to pollutants may cause histological changes in fish organ, therefore histological investigations of exposed specimens may produce meaningful results [16]. The liver is a main detoxification organ and has two essential physiological role for both the basic metabolism and the accumulation, biotransformation and excretion of toxic substances in fish [17]. The harmful effect of pollutants on fish liver histology may depend upon the duration of the exposure (chronic or acute) and the concentration level of the specific pollutants as well as other factors such as temperature, age of fish, interaction with other pollutants, water chemistry and metabolic activity of the fish [18].

The toxicity of petroleum and derivation has been studied for a few species of fish [10, 19, 20] but no information is available about the toxicity of petroleum and derivatives with milkfish.

The first aim of present study was to determine the four days lethal concentration of petroleum hydrocarbons (PHCs) to *Chanoschanos* (LC_{50} -96h).

Then this species exposed to sublethal concentration of petroleum hydrocarbons and the liver histological alteration was studied after different exposure time. This histological alteration can be used as early biological markers for evaluation of polluted aquatic ecosystems.

MATERIALS AND METHODS

Test Species and Acclimation Conditions: Milkfish, *Chanoschanos* (mean standard length: 16.7 ± 0.4 cm; mean weight: 64 ± 1.2 g) were collected from Tiyab estuary and Shoor&Shirin river estuary (relatively unpolluted site) near the city of Bandarabbas (Iran). Fish were caught with cast nets and transferred to the experimental laboratory of the Persian Gulf and Oman Sea Ecology Research Center. Specimens were washed with 0.1% $KMnO_4$ solution to remove dermal infection, if any and were acclimatized to laboratory condition for a period of 45 days before experimentation in well aerated concrete stock tank, containing natural filtered sea water. During the acclimatization process, the fish were fed twice daily with commercial dry pellets (Havoor-Rash, Iran) containing 38% crude protein, 6% crude lipid, 13% ash and 10% moisture. The water was changed daily to discard the metabolic wastes. Fitness of the test animals is of prime importance in toxicity bioassay studies. Therefore diseased fish or fish showing any abnormal behavior were removed from the tank as soon as possible. Unnecessary handling of fish was also strictly avoided.

Determination of 96-h LC_{50} of PHCs and Selection of Exposure Concentrations: After acclimatization period, only active and healthy milkfish (mean standard length: 17.82 ± 0.6 cm; mean weight: 75.3 ± 2.1 g) with no sign of distress and injury were used for this study. Specimens were randomly transferred from stock tank to polyethylene experimental tanks (280 l) which had continuous aeration. Four fish were randomly placed in each tank filled with seawater and allowed to acclimate to these conditions for two days. Fish were exposed to petroleum hydrocarbons (PHCs) concentrations of 2.4, 2.9, 3.4, 3.9, 4.4, 5.5 and 7 ml/l plus one control treatment. The tested form of PHCs was the Gasoline which was obtained from a petroleum station in Bandar Abbas, Hormozgan province, Iran. The gasoline is the main fuel for marine vehicles. Three replications were carried out for each concentration (therefore 12 fishes per treatment). The tanks were randomly arranged in a laboratory experiment were run for 96h. The day before and during the experiment the fish were not fed. Throughout the experiment fish mortality was regularly monitored every 4h. Dead fish were immediately removed from the tank to prevent contamination of the test solution. To estimate 96-h LC_{50} value (LC_{50} : the concentration of the toxicant that caused 50% mortality in fish after a specific exposure time), the total mortality of milkfish in each concentration of the toxicant (each treatment) was recorded during 96h exposure. The data was analyzed by Finney's Probit Analysis Method Program to determine 96-h LC_{50} with $\pm 95\%$ confidence [21].

Experimental Design: In the present study, young juvenile milkfish were exposed to PHCs, at concentrations of 0.26 ml/l ($1/20 LC_{50}$), 0.51 ml/l ($1/10 LC_{50}$) and 1.02 ml/l ($1/5 LC_{50}$). The 30% of test water were renewed every other day and at the same time to remove the metabolic waste and fresh pollutant added to maintain the concentration of PHCs at a constant level. 21 day exposures to these doses of PHCs were conducted and six samples were randomly collected from each treatment on acute time (12, 24 and 96h) and subchronic time (7, 14 and 21 days) at a same sampling time. The fish were quickly captured and being anesthetized by immersion, until sedate, in a clove essence solution. Standard length (L) and body weight (W) of each specimen were recorded.

Histology of the Liver: After each of the respective exposure periods, liver tissue was immediately removed. Liver samples of each specimen at the same morphological

region of each liver, to allow for comparative histological analysis, ($5 \times 5 \times 5 \text{mm}^3$). The samples were fixed in Bouin's solution for 24 h and they were transferred to 70% ethanol until routine processing for histopathological analysis using standard technique. The fixed tissues were dehydrated in ascending concentrations of ethanol series, cleared in xylene and embedded in paraffin blocks. The blocks were sectioned with a rotary microtome (Leica RM2245, Germany) at $4.5 \mu\text{m}$ thickness. Sectioned tissues were stained with Hematoxylin and Eosin [22] and observed under light microscope (Olympus-CH₄O, Japan). A total of 10 sections were randomly observed from each treatment. Histopathological alteration was assessed using ascore ranging from - to ++++ depending on the degree and extent of the alteration: (-) none, (+) mild occurrence, (++) moderate occurrence, (+++) severe occurrence and (++++) very severe occurrence. Histopathological changes induced by treatments in the tissues were photographed using Dino-lite photomicroscope.

RESULTS AND DISCUSSION

Water Analysis: The physical-chemical characteristics of the test seawater for all the experimental periods remained stable. The mean obtained values were: temperature, $23.4 \pm 0.6^\circ\text{C}$; pH, 8.1 ± 0.2 ; dissolved oxygen concentration, $7.8 \pm 1 \text{ mg/l}$ and salinity, $38.3 \pm 0.5 \text{ p.s.u}$ (practical salinity unit) [23].

Lc₅₀ Estimation: Fig. 1 shows the relation between the PHCs concentration and the mortality rate of *Cahnoschanos*. No mortality was observed during the 96h at PHCs concentrations of 2.4, 2.9 and 3.4 ml/l; and mortality were 8.33, 33.33, 58.33 and 91.67% at 3.9, 4.4, 5.5 and 7 ml/l respectively. The results obtained from acute static toxicity experiments of PHCs upon milkfish were evaluated by using Finney's Probit Analysis and 96-h LC₅₀ value of PHCs with 95% confidence for milk fish was found to be 5.12 ml/l.

Histopathology of the Liver: The liver histology of control specimens appeared relatively normal.

Compared to the control specimens, various histological changes were identified in the livers of juvenile milkfish exposed to the petroleum hydrocarbons. PHCs induced alterations in the histoarchitecture of the liver.




Fig. 1: The mortality percentage of milkfish (*Chanoschanos*) at different PHCs concentrations in 96h




Fig. 2: Sinusoid dilatation (arrow).




Fig. 3: Lipidosis, liver section showing high level of lipid accumulation (circle).

The relevant histological alterations observed in the liver of milkfish were: Sinusoid dilatation (Fig. 2), lipidosis (Fig. 3), Necrosis (Fig. 4), Vacuolations (Fig. 5), Nucleus pleomorphism (Fig. 6), Bile stagnation (Fig. 7). The histological changes in the liver tissues of exposed fish at the different sampling time and concentration are summarized in Table 1-3.

Table 1: Semiquantitative scoring of liver lesion in *Chanoschanos* during acute and subchronic exposure of 1/20 LC₅₀ PHCs

Lesion	Acute exposure (h)			Subchronic exposure (days)		
	12h	24h	96h	7d	14d	21d
Vacuolations	-	+	+	++	++	+++
Sinusoid dilatation	-	+	+	++	+++	+++
lipidiosis	-	-	+	++	++	+++
Nucleus pleomorphism	-	-	+	+	++	++
Bile stagnation	-	-	-	+	++	++
Necrosis	-	-	-	+	+	++

Score value: None (-), mild (+), moderate (++), severe (+++), very severe (++++).

Table 2: Semiquantitative scoring of liver lesion in *Chanoschanos* during acute and subchronic exposure of 1/10 LC₅₀ PHCs

Lesion	Acute exposure (h)			Subchronic exposure (days)		
	12h	24h	96h	7d	14d	21d
Vacuolations	-	+	++	++	+++	++++
Sinusoid dilatation	-	+	++	+++	+++	+++
lipidiosis	-	+	+	++	++++	++++
Nucleus pleomorphism	-	-	+	++	++	+++
Bile stagnation	-	-	+	++	++	+++
Necrosis	-	-	+	+	++	+++

Score value: None (-), mild (+), moderate (++), severe (+++), very severe (++++).

Table 3: Semiquantitative scoring of liver lesion in *Chanoschanos* during acute and subchronic exposure of 1/5 LC₅₀ PHCs

Lesion	Acute exposure (h)			Subchronic exposure (days)		
	12h	24h	96h	7d	14d	21d
Vacuolations	-	+	++	+++	++++	++++
Sinusoid dilatation	-	+	++	+++	++++	++++
lipidiosis	-	+	++	+++	++++	++++
Nucleus pleomorphism	-	-	+	++	+++	++++
Bile stagnation	-	-	+	++	+++	++++
Necrosis	-	-	+	++	+++	+++

Score value: None (-), mild (+), moderate (++), severe (+++), very severe (++++).




Fig. 4: Showing necrobiotic changes of hepatocytes (N).




Fig. 6: Nucleus pleomorphism: Nuclear megalocity (□), pyknosis (○), Nuclear deformation (Arrow).




Fig. 5: Showing vacuolation (arrow) of hepatocytes.




Fig. 7: Bile stagnation-brown granules (arrow).

The blood sinusoids are dilated between the cords of hepatocytes. According to [16, 25] Vacuolations of hepatocytes are associated with the inhibition of protein synthesis, energy depletion, disaggregation of microtubules, or shifts in substrate utilization. In this study, all treatments led to hepatocytic necrosis compared to control groups. Hepatocytic necrosis changes were observed in fish exposed over one day periods [26]. Lipidiosis (the excessive accumulation of fat in the cytoplasm) is characteristic of many exposed livers. The large vacuole in the cell forces the nuclei to the periphery of the hepatocytic and this condition is usually accompanied by nuclear atrophy [8, 16].

Nuclear pleomorphism is characterized by marked nuclear enlargement, nuclear pyknosis and deformation of spherical structure of nucleus in the present study. Hepatocellular pleomorphism is thought to be sublethally injured cells resulting from exposure to several types of toxicants [26].

Bile stagnation is characterized by the remains of the bile in the form of brownish-yellow granules in the cytoplasm of the hepatocytes [27], indicates that the bile is not being released from the liver. This accumulation of bile indicates possible damage to the hepatic metabolism [28].

As a conclusion, the findings of the present histological investigations demonstrate a direct correlation between petroleum hydrocarbons (PHCs) nonlethal concentrations and exposure time with histopathological disorders and lesions observed in milkfish liver tissue. Therefore, histopathology is a useful biomarker for environmental contamination, since it discriminated between control and test groups. The liver showed to be the organ affected by the type of stressors to which the fish were subjected. The *Chanochanos* was shown to be appropriate for environmental monitoring and *in situ* tests because of its hardly nature to survive under laboratory conditions.

ACKNOWLEDGMENT

The authors would like to gratefully acknowledge the staff of aquaculture lab at *Persian Gulf and Oman Sea Ecology Research Center* for their assistance with this project. Thanks go to the staff at the department of marine biology at Khorramshahr University of marine science and technology. Special thanks for *Dr. Movahedinia* for his guidance and for *AzadehAtabati* for histological help.

REFERENCES

1. Botello, A.V., S.F. Villanueva and G.G. Diaz, 1997. Petroleum pollution in the Gulf of Mexico and Caribbean Sea. *Reviews of environment contamination and toxicology*, 153: 91-118.
2. Neff, J.M., 1979. Polycyclic Aromatic Hydrocarbons in the Aquatic Environment Sources, Fates and Biological Effects. *Applied Science*, Barking, Essex.
3. De la Torre, F.R., A. Salibian and L. Ferrari, 2000. Biomarkers assessment in juvenile *Cyprinus carpio* exposed to waterborne cadmium. *Environmental Pollution*, 109: 277-282.
4. Sikkema J., J.A. De Bont and B. Poolman, 1994. Interactions of cyclic hydrocarbons with biological membranes, *Journal of Biological Chemistry*, 269: 8022-8.
5. Lopes, P.A., T. Pinheiro, M.C. Santos, M. da Luz Mathias, M.J. Collares-Pereira and A.M. Viegas-Crespo, 2000. Response of antioxidant enzymes in freshwater fish populations (*Leuciscus alburnoides complex*) to inorganic pollutants exposure. *Science of the Total Environment*, 280: 153-163.
6. Gravato, C. and M.A. Santos, 2002. Juvenile sea bass liver P450, Erod induction and erythrocytic genotoxic responses to PAH and PAH-like compounds, *Ecotoxicology and Environment Safety*, 51: 115-27.
7. Ankyakora, C., A. Ogbeche, P. Palme and H. Coker, 2005. Determination of polynuclear aromatic hydrocarbons in marine samples of Siokolo Fishing Settlement. *Journal of Chromatography*, 1073(1-2): 323-30.
8. Simonato, J.D., C.L.B. Guede and C.B.R. Martinez, 2008. Biochemical, physiological and histological changes in the neotropical fish *Prochilodus lineatus* exposed to diesel oil. *Ecotoxicology and Environment Safety*, 69(1): 112-20.
9. Brand, D.G., R. Fink, W. Bengueyfield, L.K. Birtwell and C.D. Mcallister, 2001. Salt water acclimated pink salmon fry (*oncorhynchus gorbuscha*) develop stress-related visceral lesion after 10 days exposure to sublethal concentrations of the water-soluble fraction of north slope crude oil. *Toxicological Pathology*, 29(5): 574-84.

10. Alkaishi, F.M., H.C.S. Assis, S.C.G. Jakobi, D.R. Eiras-Stoffla, S.D. St-Jean, S.C. Courtenay, E.F. Lima, A.L.R. Wagener, A.L. Scofield and C.A. Oliveira Ribeiro, 2004. Morphological and neurotoxicological findings in tropical freshwater fish (*Astyanax sp.*) after waterborne and acute exposure to water soluble fraction (WSF) of crude oil. Archives of Environmental Contamination and Toxicology, 46: 244-253.
11. Abou El-naga, E.H., K.M. El-moselhy and M.A. Hamed, 2005. Toxicity of cadmium and copper and their effect on some biochemical parameters of marine fish *Mugilseheli*. Egyptian Journal of Aquatic Research, 1687-4285. 31(2): 60-71.
12. Wang, M.E. and Q.X. Zhou, 2006b. Effects of herbicide chlorimuron-ethyl on physiological mechanisms in wheat (*Triticumaestivum*). Ecotoxicology and Environmental Safety, 64: 190-197.
13. Zhou, Q.X., 2006. Advance in the effects of climate change on environment and health. Chinese Journal of Meteorology and Environment, 22: 38-44.
14. Ferrat, L., C. Pergent-Martini and M. Romeo, 2003. Assessment of the use of biomarkers in aquatic plants for the evaluation of environmental quality: application to seagrasses. Aquatic Toxicology, 65: 187-204.
15. Almedia, J.A., Y.S. Diniz, S.F.G. Marques, L.A. Faine, B.O. Ribas, R.C. Burneiko and E.L.B. Novelli, 2002. The use of the oxidative stress responses as biomarkers in Nile tilapia (*Oreochromis niloticus*) exposed to *in vivo* cadmium contamination. Environment International, 27: 673-679.
16. Hinton, D.E. and D.J. Lauren, 1990. Integrative histopathological effects of environmental stressors on fishes. American Fish Society Symposium, 8: 51-66.
17. Fernandes, C., A. Fontainhas-Fernandes, E. Rocha and M.A. Salgado, 2008. Monitoring pollution in Esmoriz-Paramos lagoon, Portugal: Liver histological and biochemical effects in *Liza saliens*. Environment Monitoring and Assessment, 145: 315-322.
18. Heath, A.G., 1995. Water Pollution and Fish Physiology. CRC Press, Boca Raton, pp: 245.
19. Khan, R.A., 2003. Health of flatfish from localities in Placentia Bay, Newfoundland, contaminated with petroleum and PCBs. Archives of Environmental Contamination and Toxicology, 44: 485-492.
20. Neff, J.M., S. Ostazeski, W. Gardiner and I. Stejskal, 2000. Effects of weathering on the toxicity of three offshore Australian crude oils and a diesel fuel to marine animals. Environmental Toxicology and Chemistry, 19: 1809-1821.
21. Finney, D.J., 1971. Probit Analysis, third ed. Cambridge University Press, London, pp: 337.
22. Woods, A.E. and R.C. Ellis, 1994. Laboratory Histopathology: A Complete Reference. Churchill Livingstone, New York.
23. APHA, American Public Health Association, American Water Works Association and Water Pollution Control Federation, 1998. Standard methods for the examination of water and wastewater. 20th ed. APHA, New York.
24. Marigomez, I., M. Soto, I. Cancio, A. Orbea, L. Garmendia and M.P. Cajaraville, 2006. Cell and tissue biomarkers in mussel and histopathology in hake and anchovy from Bay of Biscay after the Prestige oil spill (Monitoring Campaign 2003). Marine Pollution Bulletin, 53: 287-304.
25. Van Dyk, J.C., G.M. Pieterse and J.H.J. van Vuren, 2007. Histological changes in the liver of *Oreochromis mossambicus* (Cichlidae) after exposure to cadmium and zinc. Ecotoxicology and Environmental Safety, 66: 432-440.
26. Agamy, E., 2012. Histopathological liver alterations in juvenile rabbit fish (*Siganus canaliculatus*) exposed to light Arabian crude oil, dispersed oil and dispersant. Ecotoxicology and Environmental Safety, 75(1): 171-179.
27. Pacheco, M. and M.A. Santos, 2002. Biotransformation, genotoxic and histopathological effects of environmental contaminants in European eel (*Anguilla anguilla L.*). Ecotoxicology and Environmental Safety, 53: 331-347.
28. Fanta, E., F.S. Rios, S. Romao, A.C.C. Vianna and S. Freiberger, 2003. Histopathology of the fish *Corydoras paleatus* contaminated with sublethal levels of organophosphorus in water and food. Ecotoxicology and Environmental Safety, 54: 119-130.