

Stability and Adaptability of Promising Lowland Rice Genotypes for Yield and Cold Tolerance in High Elevation Rice Growing Areas of Ethiopia

¹Tadesse Lakew, ¹Abebaw Dessie, ¹Asaye Berie, ¹Betelehem Asrat and ²Dejene Kebede

¹Ethiopian Institute of Agricultural Research, Fogera Rice Research and Training Center, Ethiopia

²Ethiopian Institute of Agricultural Research, Jimma Agricultural Research Centre, Jimma, Ethiopia

Abstract: Multi-environmental yield trial is imperative in assessing rice genotypes for adaptability and stability across environments having different stresses prior to release of the newly developed variety for commercial cultivation. The performance and stability variation in genotypes are the combination of environment, genotype and genotype by environment (GE) interaction. With the objective of assessing and identifying high yielding, stable and cold tolerant lowland rice varieties, a total of sixteen rice genotypes were arranged in a randomized complete block design of four replications. Analysis of variance revealed highly significant effects of genotype and environment for all traits studied while the interaction effect was significant for six of eleven traits. The AMMI analysis in grain yield showed that genotype, environment and their interaction were highly significant and the environment explained the highest variation, followed by the interaction. The first two multiplicative interaction principal component axes were highly significant and explained 83.8% of the interaction sum of squares. Spikelet fertility ranged from 89.9% (G1) to 97.8% (G9 and G11). Genotypes G4, G9, G11, G10, G12, G13 and G14 exhibited nearly complete panicle exertion and high spikelet fertility indicating their tolerance to cold stress. AMMI 1 and GGE ranking biplots identified G4, G9, G11 and G12 as high yielding genotypes. While G9 was the best genotype in terms of mean yield, G4 was both high yielding and most stable genotype. Thus, genotypes that combine cold tolerance, high yield and farmers' preference (G4, G9 and G11) were verified and consequently, G9 and G11 were recommended for release in Fogera and similar areas.

Key words: Rice Genotype • Cold Tolerance • AMMI Biplots • GGE Biplots • Grain Yield

INTRODUCTION

Rice (*Oryza sativa* L.) is a tropical and subtropical, diploid ($2n=24$) and self-pollinating crop that belongs to the Gramineae family [1- 3]. The crop is the most important cereal in the world and one of the main staple foods for millions of people in developing countries including Sub-Saharan Africa. It is the principal source of energy, protein, iron, calcium, thiamine, riboflavin and niacin in many developing countries where rice is the major food source. Rice being a tropical and sub-tropical plant requires a fairly high temperature, ranging from 25° to 35°C. The optimum temperature of 30°C during day time and 20°C during night time is considered favorable for its growth and development [4]. Temperatures below 25°C can cause growth abnormalities in temperate and high-elevated tropical areas [5]. Cold stress shows different effects on germination, seedling, vegetative, reproductive

and grain maturity and weakens photosynthetic ability by inducing leaf discoloration, reduces plant height, produces degenerated spikes, delays days to heading, reduces spikelet fertility and can cause yield losses of 26% to 80% [6-8]. Although the main symptom of damage from cold is the high spikelet sterility [9], incomplete panicle exertion has also been cited as a symptom of cold injury in many rice growing countries [10] and this trait has been suggested as an indicator of genotype adaptability to cold affected areas [11].

In Ethiopia, rain fed rice is predominantly cultivated in altitudes above 1000 m.a.s.l. [12, 13] such as Fogera, Dera, Libokemkem, Dembia, North Achefer, Tis-Abay, Pawe, Jimma, Bako-Chewaka, Guraferda, Assosa-Kamashi and Tepi for both home consumption and local market. Districts such as Fogera, Dera, Libokemkem, Dembia and Jimma, with minimum daily temperature of always < 15°C, most often experience damage to rice crop due to low

temperature effect. Ethiopia is also reported to have several water lodged areas where other crop plants do not grow well and are assumed to be potential for rice cultivation. Expanding rice cultivation to such localities in the country could contribute towards meeting the food demand of the rapidly growing human population and help curb the huge annual rice import. However, such areas often experiences low temperature effect. Except few, most improved rice varieties in Ethiopia are sensitive to cold stress and hence cannot be grown in such cold prone areas which implicate the critical demand for cold tolerant, high yielding and stable rice varieties. It is therefore important to assess the performance of rice genotypes in areas that are most often cold affected with a view of growing the genotypes in the high elevation areas of Ethiopia.

However, multi-location evaluation of rice genotypes for identifying high yielding, stable and cold tolerant rice varieties is not a straight forward. Plant breeders most frequently encounter Genotype by Environment Interaction (GEI) when testing genotypes across a number of environments influencing the selection process [14]. Several methods are employed to estimate stability and adaptability performance in genotypes tested across environments by determining GEI effects. The two frequently used analysis methods are the Additive Main effects and Multiplicative Interaction (AMMI) model and the genotype main effects and GEI effects (GGE) model [5]. AMMI is preferred when both main effects and interaction effects are important and it integrates analysis of variance and principal component analysis [16]. The GGE provides visual evaluation of the data by creating a biplot that simultaneously represents mean performance and stability as well as identifying winner genotypes with respective mega-environments and relationships among testing environments [17, 18]. Therefore, the objectives of this investigation were (1) to evaluate introduced lowland rice genotypes for cold tolerance under field conditions and (2) to assess adaptability and stability performance of rice genotypes for grain yield and related traits and identify genotypes that showed higher grain yield performance with wider or specific adaptation.

MATERIALS AND METHODS

Plant Materials: Through Africa Rice breeding task force network and in connection with STRASA (Stress-Tolerant rice for Africa and South Asia) project, a total of 99 lowland rice genotypes were obtained for cold tolerance

evaluation and for overall adaptation in Ethiopia. Of these, thirty three rice genotypes which showed better performance and cold tolerance were selected for preliminary variety trial in 2013 main cropping season, out of which only 14 genotypes were selected and promoted to multi-location variety trial in 2014 and 2015. Including two control varieties (one improved and cold tolerant i.e *Ediget* and another cold tolerant local variety i.e *X-Jigna*), a total of 16 rice genotypes were evaluated as national variety trial for their adaptability, stability and performance in grain yield and related traits in addition to their tolerance to cold stress under field conditions (Table 1).

Testing Environments: The experiment was conducted under rain fed lowland conditions at research fields of Fogera (Woreta and Kokit), Jimma (Jemma and Gojeb) and Assosa (Assosa) research centers during the main cropping seasons (May to November) of 2014 and 2015 (Table 2). At these locations, the rice crop often experiences low temperature effect at least once during the growing periods. The locations varied in rainfall amount and distribution, elevation, temperature (Table 2) and soil types (data not shown). The location-year combination indicated seven environments such as E1 (Woreta2014), E2 (Jimma2014), E3 (Assosa2014), E4 (Woreta2015), E5 (Kokit2015), E6 (Jimma2015) and E7 (Gojeb2015) in this study.

Experimental Design and Trial Management: At each location, the experiment was laid out using a randomized complete block design of four replications. Seeds of each genotype were planted at the rate of 60 kg/ha by hand drilling in a plot size of 7.5m² and with a spacing of 25cm between rows. Each experimental plot was with six rows of 5m long each. Fertilizers (Urea and DAP) were applied as per to local recommendations. All DAP was applied at planting while Urea was used in three splits; planting, tillering and panicle initiation. Other crop management and protection practices were applied to the entire experimental area uniformly.

Data Collection: Data on yield, yield related and other morphological traits were collected both on plot and individual plant bases. Five randomly selected plants from the middle four rows of each experimental plot were used in plant base data collection for panicle length (PL, cm), plant height (PH, cm), number of filled grains per panicle (FGP) and fertility rate (FR, %) was calculated from the ratio of filled grains per panicle to total grains per panicle.

Table 1: Description of 16 rice genotypes used in this study

No.	Genotypes	Code	Source
1	scrid079-1-5-4-2	G1	Africa Rice
2	exp304	G2	Africa Rice
3	FOFIFA 161	G3	Africa Rice
4	FOFIFA 171	G4	Africa Rice
5	FOFIFA 172	G5	Africa Rice
6	FOFIFA 167	G6	Africa Rice
7	HR 17512-11-2-3-1-4-2-3	G7	Africa Rice
8	scrid006-2-4-3-4-5	G8	Africa Rice
9	scrid006-3-2-3-2	G9	Africa Rice
10	scrid14-1-1-1-1	G10	Africa Rice
11	scrid017-1-4-4-4-1	G11	Africa Rice
12	scrid019-1-1-1-1-2	G12	Africa Rice
13	scrid037-4-2-2-5-2	G13	Africa Rice
14	scrid113-3-5-3-5-4	G14	Africa Rice
15	Edget (standard check)	G15	Fogera NRRTC, Ethiopia
16	X-JIGNA(local check)	G16	Fogera NRRTC, Ethiopia

Table 2: Description of testing locations used for the evaluation of 16 rice genotypes

Location	Coordinates			Rain fall (mm)	Temperature (°C)	
	Latitude	Longitude	Elevation (m asl)		Maximum	Minimum
Woreta	11° 58' N	37° 41' E	1810	1292.6	26.7	13.0
Jimma	7° 46' N	36° 00' E	1753	1091.0	26.9	11.1
Kokit	11° 92' N	37° 7' E	1780	1292.6	26.7	13.0
Assosa	10° 03' N	34° 59' E	1590	1020.0	26.7	15.2
Gojeb	7°15' N	36°0' E	1235	1710.0	24.0	16.7

masl: meter above sea level, mm: millimeter, Note: Rainfall amount and temperature conditions refers to the average values during rice growing periods at each location. These data were provided by the Ethiopian Meteorology Agency.

Data for days to 50% heading (DHT, days), days to 85% maturity (DMT, days), panicle exertion (Panex-scale: 1-5; 1 completely exerted, 5 remain closed), phenotypic acceptability (Phac-scale: 1-9; 1 fully accepted, 9 completely poor and not accepted) were collected on plot basis following standard evaluation system of IRRI [19]. Grain yield (Gy) and thousand seed weight (TSW) were also taken on plot basis from the four harvestable rows in gram. Grain yield harvested from each plot was converted into kg/ha at 14% standard grain moisture content.

Statistical Analysis: Data collected on grain yield, agronomic and morphological traits were subjected to analysis of variance using the General Linear Model (PROC GLM) of the SAS procedure version 9.0 of the SAS software [20] to determine significant variation among genotypes and environments and their interaction. Mean performance of different traits were separated using Least Significant Difference (LSD) method at 0.05 level of probability. Additive main effects and multiplicative interaction (AMMI) model was applied to assess the effect of genotype by environment interaction,

adaptability and stability of rice genotypes [16] using GenStat 16th edition statistical package. Moreover, GGE analysis, according to [21], was employed to visualize grain yield stability and performance of 16 rice genotypes at seven environments.

RESULTS AND DISCUSSION

Genotypes Performance for Yield and Cold Tolerance

Traits: Combined analysis of variance and overall mean performance of 16 rice genotypes at seven environments for all agronomic traits as well as cold tolerance indicator traits (spikelet fertility and panicle exertion) are presented in Table 3. Highly significant genotype and environment effects were observed for spikelet fertility and panicle exertion and for all other traits. The genotype by environment interaction (GEI) was highly significant for spikelet fertility but not for panicle exertion. Similarly, GEI was also highly significant for five traits including grain yield but not for days to heading, panicle length, fertile tillers/plant and thousand seed weight (Table 3). The presence of highly significant

Table 3: Mean performances of lowland rice genotypes for grain yield (kg/ha) and yield related traits across seven environments during 2014 and 2015 main cropping season

Genotype	DH	DM	PL	PH	FTP	FGP	FR	TSW	Gy	Panex	Phac
G1	93.6	132.1	19.8	103.1	7.5	90.5	89.9	27.4	3952.2	2.9	4.9
G2	86.7	130.1	18.7	100.2	8.0	91.5	93.6	28.1	4184.0	3.6	4.5
G3	89.8	128.8	16.6	83.8	6.3	86.4	94.8	28.1	4070.8	3.9	4.5
G4	89.9	130.0	19.2	97.7	7.1	84.6	97.3	29.8	4826.3	2.0	3.4
G5	78.0	126.0	15.2	80.4	8.3	55.8	93.0	28.9	3196.7	5.1	7.2
G6	91.4	127.0	17.7	105.2	7.8	85.2	92.5	24.1	4108.5	2.3	3.9
G7	95.7	134.4	16.2	72.0	8.2	89.7	92.1	20.1	3934.8	4.3	5.6
G8	86.4	126.1	19.0	102.8	7.8	76.2	94.0	34.6	4385.7	2.0	3.4
G9	98.4	125.8	19.0	97.6	8.2	79.4	97.8	34.7	5166.4	1.7	2.8
G10	89.0	129.2	19.5	106.6	7.1	108.4	94.6	21.8	4449.0	1.7	2.6
G11	90.1	129.0	18.4	105.3	7.2	106.1	97.8	24.3	4841.2	1.6	1.7
G12	92.4	130.5	18.5	101.1	6.8	103.0	94.2	25.4	4903.9	1.6	2.4
G13	93.0	128.8	18.5	102.5	7.1	104.0	94.5	25.1	4588.0	1.8	2.7
G14	90.1	127.8	17.3	100.4	7.1	87.7	94.3	24.4	4146.6	1.8	2.8
G15	87.4	130.8	17.4	88.1	6.1	80.2	92.0	29.7	3671.4	4.0	4.5
G16	95.4	132.3	19.1	97.6	6.9	94.0	91.4	25.5	3637.3	2.1	4.0
Mean	89.8	96.5	18.1	96.5	7.34	88.9	94.0	27	4254.0	2.67	3.8
CV (%)	10.23	6.11	8.1	6.1	23.8	16.65	3.73	10.51	23.1	32.8	31.34
LSD (0.05)	4.84	2.08	0.77	3.1	0.92	7.78	1.84	1.49	516.62	0.46	0.63
Genotype (G)	***	***	***	***	***	***	***	***	***	***	***
Environment (E)	***	***	***	***	***	***	***	***	***	***	***
G x E	NS	**	NS	**	NS	***	***	NS	***	NS	***

*, ** and ***, significant at 0.05, 0.01 and 0.001 levels, NS: not significant at 0.05 level, DH: days to heading, DM: days to maturity, PL: panicle length (cm), PH: plant height (cm), FTP: fertile tillers per panicle, FGP: filled grains per panicle, FR(%): fertility rate of spikelets per panicle, TSW: 1000 seed weight (g), Gy: grain yield (kg ha⁻¹), Panex: panicle exertion (scale; 1-5, 1-fully exserted, 5-remain closed), Phac: phenotypic acceptability (scale; 1-9, 1-fully accepted and 9-not accepted at all).

Table 4: Mean grain yield (kg ha⁻¹) performance of 16 rice genotype at seven environments

Genotype	E1	E2	E3	E4	E5	E6	E7	Mean
G1	2734	4216	3235	5867	4949	1855	4807	3952.2
G2	4254	5002	2404	4922	4938	6321	3466	4184.0
G3	3145	4811	3833	5998	3939	5575	3701	4070.8
G4	4327	5068	4796	4945	5102	5780	3807	4826.3
G5	2520	4568	2823	5354	3534	4670	3024	3196.7
G6	3583	4744	2966	4056	3895	5750	4118	4108.5
G7	3977	3731	2405	4750	4415	4156	4939	3934.8
G8	3387	3587	2785	4288	4607	7251	4553	4385.7
G9	3969	3797	2881	4108	5033	9580	4733	5166.4
G10	4083	3790	2719	4512	4342	5343	4580	4449.0
G11	4309	5044	4322	4928	4656	5212	4479	4841.2
G12	4183	2821	4699	2985	4910	6044	4547	4903.9
G13	3739	4556	3762	4043	4979	5167	3801	4588.0
G14	3810	3877	3135	4595	4077	5404	3652	4146.6
G15	3031	3654	2433	4076	3777	4325	4378	3671.4
G16	3225	4895	2624	5253	4016	3344	3966	3637.3
Mean	3642.3	4260.1	3238.9	4667.5	4448.1	5361.1	4159.4	4254.0
CV (%)	9.5	15.6	33.7	19.1	17.1	26.3	21.2	23.1
LSD (0.05)	491.2	946.6	1553.4	1268.1	1080.5	2005.3	1253.8	516.6

E1: Woreta2014, E2: Jimma 2014, E3: Assosa2014, E4: Woreta2015, E5: Kokit2015, E6: Jimma2015, E7: Gojeb2015

G x E interactions for grain yield indicates that genotypes tended to rank differently in grain yield at different locations and over years. Genotypes (G9, G10, G11 and G12) with high performance in cold tolerance indicator

traits (spikelet fertility and panicle exertion) also showed high performance in grain yield indicating their linear relationship and it suggested that higher yielding genotypes showed strong tolerance to cold stress under

field condition across locations (Table 3). The overall mean grain yield of 16 rice genotypes ranged from 3196.7 kg ha⁻¹ to 5166.4 kg ha⁻¹ with a grand mean yield of 4254.0 kg ha⁻¹ and the highest grain yields were obtained from Genotypes G9, G12, G4 and G11 (Table 3). As indicated in Table 3, seven genotypes scored highest grain yield over grand mean (4254.0 kg ha⁻¹) and thirteen genotypes scored highest grain yield over standard (*Ediget*) and local checks (*X-Jigna*). The wide variation in grain yield explained by the environments revealed the presence of different mega-environments where specific genotypes performed best within each mega-environment.

The average environmental grain yield across genotypes ranged from 238.9 kg ha⁻¹ at Assosa2014 to 5361.1 kg ha⁻¹ at Jimma2015, with a grand mean yield of 4254.0 kg ha⁻¹ (Table 4). Environments E4 (4667.5 kg ha⁻¹) and E6 (5361.1 kg ha⁻¹) were the highest yielding environments while E1 and E3 were the lowest whereas E2, E5 and E7 scored intermediate mean grain yield (Table 4). Average grain yield of genotypes across environments ranged from the lowest of 3196.7 kg ha⁻¹ for G5 to the highest of 5166.4 kg ha⁻¹ for G9. The genotypes G4, G9, G11 and G12 had higher average grain yield across environments with average yield of 4826.3, 5166.4, 4841.2 and 4903.9 kg ha⁻¹, respectively (Table 4). Genotype G4 ranked first at four environments (E1, E2, E3 and E5). The other better performing genotypes at different environments include G1 (E4, E5, E6 and E7), G2 (E1, E2 and E6), G3 (E3 and E4), G5 (E4), G7 (E7), G9 (E5, E6 and E7), G10 (E7), G11 (E1, E2, E3), G12 (E1, E3 and E6) and G13 (E5) (Table 4). The local check (G16 i.e. *X-Jigna*) showed better performance for average grain yield at two environments (E2 and E4). Genotypes G3 and G9 recorded the best yield of 5998 kg ha⁻¹ and 9580 kg ha⁻¹ at the highest yielding environments (E4 and E6), respectively, while G4 had the best yield of 4327 kg ha⁻¹ and 4796 kg ha⁻¹ at the lowest yielding environments, E1 and E3 (Table 4).

AMMI Analysis: Significant analysis of variance for the main effects and their interaction in grain yield indicated the importance of the additive main effects and multiplicative interaction (AMMI) model for further interpreting the grain yield data to identify stable genotypes. AMMI model calculates genotypes and environment additive (main) effects using analysis of variance and then analyze the residual from this model using principal component analysis (PCA) [16]. AMMI analysis of variance for grain yield revealed highly significant effect for genotypes, environments and genotype by environment interaction (Table 5). The effect

of the environment was higher than the genotype but lower than the interaction effect (data not shown) indicating wider variation among environments and differential response of genotypes across environments. The AMMI analysis also showed that the first interaction principal component axis (IPCA1) and second interaction principal component axis (IPCA2) explained 70.57% and 13.26% of the interaction sum squares, respectively (Table 5). Significant interaction indicated that the genotypes respond differently across the environments. The report in the present study are in line with the previous findings [22] and AMMI procedure has been widely applied by various authors in selection of stable rice genotypes [23, 24].

AMMI recommendation of genotypes for different environments is shown in Table 6. AMMI selection identified first four high yielding genotypes per environment. The result indicated that some high yielding genotypes are suitable for specific environment (G10, G3) and some other genotypes (G4, G11, G2, G12, G8, G9, G13, G1) can be recommended for two or more environments. The illustration in Table 6 suggested that we can advise the right genotype for all environments or specific genotype for specific environments through AMMI evaluation. As reported by Bantayehu *et al.* [22] and Tekdal and Kendal [25] the AMMI analysis indicates recommendations of correct genotype for one or more environments. In our report, therefore, the selection of best candidates based on overall performance above coincides with that of AMMI recommendation as G4, G9 and G11 work for more than one environment. Table 7 also showed mean grain yield performance and stability coefficient of 16 lowland rice genotypes across seven environments through AMMI analysis. The mean grain yield value of genotypes averaged over environments indicated that genotype G9 had the highest (5166 kgha⁻¹) and genotype G5 (3197 kgha⁻¹) the lowest grain yield, respectively. Genotype superiority with the smallest and the highest measured values indicate the more stable and less stable genotypes, respectively. Therefore, from the present study, genotype G9 was the most stable, followed by genotypes G12, G8, G4 and G11 while genotype G1 was the least stable genotype, followed by genotypes G5, G16, G15, G7 and G14 (Table 7).

AMMI Biplots: In the AMMI biplot analysis, the IPCA scores of environments and genotypes are plotted against main effect means and the plot helps to visualize the average productivity of the genotypes, environments and their interaction. The nature and magnitude of interaction can be visualized for each genotype and each

Table 5: AMMI analysis of variance for grain yield (kg ha^{-1}) of 16 lowland rice genotypes grown at seven environments

Source	d.f.	s.s.	m.s.	v.r.	F pr
Total	447	828169840	1852729		
Genotypes	15	117483339	7832223	9.33	<0.001
Environments	6	182255791	30375965	11.15	<0.001
Interactions	90	206788573	2297651	2.74	<0.001
IPCA 1	20	145929536	7296477	8.69	<0.001
IPCA 2	18	27420057	1523336	1.81	0.0229
IPCA3	16	13356291	834768ns	0.99	0.4625
Error	315	264430076	839461		

d.f: degree of freedom, s.s: Sum square, m.s: mean square, v.r: variance ratio, Fpr: probability level.

Table 6: AMMI selections of genotypes per environment

Environment	IPCAe[1]	IPCAe[2]	Mean (kg ha^{-1})	Score	First four AMMI selections			
					1	2	3	4
E1	0.9587	-0.8826	3642	0.96	G4	G11	G2	G12
E2	7.5632	-4.2214	4260	7.56	G12	G4	G10	G9
E3	15.2395	-41.7081	3239	15.24	G4	G12	G11	G3
E4	24.0049	13.2793	4668	24.00	G11	G1	G13	G9
E5	8.3692	7.6541	4448	8.37	G4	G9	G13	G1
E6	-70.0667	0.8925	5361	-70.07	G9	G8	G2	G12
E7	13.9313	24.9864	4160	13.93	G8	G1	G10	G11

IPCAe1 and IPCAe2: the first and second interaction principal component axis for the environments.

Table 7: Mean grain yield (kg ha^{-1}) and stability coefficient 16 lowland rice genotypes

Genotypes	Grain yield (kg ha^{-1})	Genotype superiority
G1	3952	4672945 (16)
G2	4184	1695345 (7)
G3	4071	1792192 (10)
G4	4826	1204886 (4)
G5	3197	3679873 (15)
G6	4109	1789834 (9)
G7	3935	2841980 (12)
G8	4386	1163475 (3)
G9	5166	323865 (1)
G10	4449	1722031 (8)
G11	4841	1406870 (5)
G12	4904	992304 (2)
G13	4588	1551579 (6)
G14	4147	1906506 (11)
G15	3671	2973606 (13)
G16	3637	3673647 (14)

Numbers in brackets give the position of each genotype, ranked according to the stability coefficient (running downwards from 1 = best)

environment using IPCA1 vs. IPCA2 biplot of AMMI 2 [22, 23]. In AMMI1 biplot (Fig. 1), genotype and environment main effects are drawn against their corresponding IPCA1 score. In this biplot, the genotypes with IPCA1 scores close to zero tend to express general adaptation and the larger scores showed more specific adaptation to one or more environments. Rice genotypes plotted on the right-hand side of the vertical midline have

higher grain yield compared to those on the left-hand side (Fig. 1). Genotypes G9, G12, G4, G11, G13, G10 and G8 had high average yield with G9, G12, G11 and G4 being the highest yielding genotypes. Genotypes G11 and G13 adapted to E4, E5 and E2 while genotypes G8 and G9 adapted to the most favorable environment (E9). On the other hand, genotypes G4, G12, G10, G14 and G3 had small IPCA1 scores, suggesting their general adaptation. Genotypes G1, G16, G15, G7 and G5 had low average yield and G1 had the largest positive IPCA1 score making it the most unstable genotype. Of the environments, E1, E2 and E5 exhibited relatively small IPCA1 scores, suggesting that they had little interaction with genotypes indicating stable environments, of which E1 being the most stable environment. On the other hand, two favorable environments (E4 and E6), one intermediate environment (E7) and the poorest environment (E3) had relatively high IPCA1 scores producing large interactions with genotypes, making them unstable (Fig. 1).

As demonstrated by AMMI 2 biplot in Fig 2, environments with short vectors exert weak interaction and those with long vectors exert strong interaction [26]. Environments E3, E4, E6 and E7 with long vectors were very interactive and discriminated differences among genotypes more than other environments with short vectors (E1, E2 and E5) which were less interactive and provide little information about differences among genotypes for grain yield performance [23].




Fig. 1: AMMI1 biplot showing the main effect vs. stability (IPCA1) of both genotypes and environments on grain yield (kg ha^{-1}). E1: Woreta2014, E2: Jimma 2014, E3: Assosa2014, E4: Woreta2015, E5: Kokit2015, E6: Jimma2015, E7: Gojeb2015.




Fig. 2: AMMI 2 biplot graph showing interaction of PC1 and PC2 scores of 16 rice genotypes at seven environments. E1: Woreta2014, E2: Jimma 2014, E3: Assosa2014, E4: Woreta2015, E5: Kokit2015, E6: Jimma2015, E7: Gojeb2015.

The genotypes near the origin are not sensitive to environmental interaction and those away from the origins are sensitive and have large interaction [17]. In this result, G1, G8, G9 and G16 are sensitive to the environments since they were away from the origin whereas most other genotypes were relatively close to the origin and hence they were less sensitive to the environments.

GGE Biplots: Fig. 3 shows the polygon view of which-when-where pattern in GGE biplot analysis of mean grain yield in 16 lowland rice genotypes evaluated at seven

environments. The GGE biplot accounted for 70.02% of the total variation consisting of 51.19% and 19.01% of the variation attributed to the first and second principal components (PC1 and PC2), respectively. The polygon in the biplot was divided in to six sectors in which environments are grouped to different sectors. The genotype at the vertexes of each sector is the best genotype to the environment(s) contained in the particular sector and all other genotypes within the sector are recommended for all environments contained in that sector [26]. In this study, GGE biplot analysis identified




Fig. 3: GGE biplot showing winner genotypes at specific environments. E1: Woreta2014, E2: Jimma 2014, E3: Assosa2014, E4: Woreta2015, E5: Kokit2015, E6: Jimma2015, E7: Gojeb2015




Fig. 4: GGE biplot showing grain yield rank and stability of 16 lowland rice genotypes across seven environments. E1: Woreta2014, E2: Jimma 2014, E3: Assosa2014, E4: Woreta2015, E5: Kokit2015, E6: Jimma2015, E7: Gojeb2015.

G1 as the best genotype at two environments (E4 and E7), which represented one mega-environment. The winner genotypes at two environments (E3 and E5) were G10, G11 and G13; with G11 being the best fit genotype. Similarly, genotypes G4 and G9 performed best at environments E1 and E6. On the other hand, the remaining two sectors with G2 and G5 as vertex genotypes contained no environment within them suggesting that grain yield performance of all genotypes, including checks (G15 and G16), in the two sectors was low in any of the environments.

Fig. 4 shows ranking of 16 lowland rice genotypes based on mean grain yield performance and stability using genotype focused scaling of GGE analysis. The average-environment coordination (AEC) line (abscissa) which passes through the biplot origin points towards the direction of high mean yield across environments and the vector line of each genotype which is perpendicular to the AEC indicates the stability of the varieties. Regardless of the direction of vectors, varieties with longer vector are less stable compared to varieties with shorter vectors [27]. Genotypes placed below the

AEC ordinate line (red line in this case) performed below the average and those placed above ordinate line performed above average [17, 27]. Thus, genotypes G12, G4, G9 and G11 were the best performing genotypes for mean grain yield, followed by G13 and G10 while G1, G2, G3, G6, G7, G8, G14, G16 and G15 were low yielding genotypes, with G5 being the least in grain yield (Fig. 4). Genotypes G3, G7, G10 and G15 were relatively stable genotypes while genotypes G1, G2, G5 and G16 were less stable, with G1 being the most unstable genotype. Of the top four high yielding genotypes (G4, G9, G11 and G12), G12 was the most stable genotype, followed by G4, while G9 and G11 were less stable. The latter two high yielding genotypes could be recommended to specific environments. With regard to environments, four environments (E1, E4, E6 and E7) were relatively unstable while the remaining three (E2, E3 and E5) were stable environments (Fig. 4) though E3 was illustrated contrary to this in AMMI 2 biplot.

CONCLUSION AND RECOMMENDATION

The current study assessed the performance, stability and cold tolerance in 16 lowland rice genotypes under rain fed field condition. All the traits considered were significantly affected by genotype, environment and genotype by environment interaction. The significant interaction effect for grain yield indicated the inconsistent performance of genotypes across test locations and over seasons. Application of different statistical techniques including ANOVA, AMMI and GGE biplots enabled us to detect variation and identify broadly and/or specifically adapted and high yielding rice genotypes. As result, desirable lowland rice genotypes in terms of cold tolerance, yield performance and stability had been identified that can be recommended to farmers for cultivation. Although no genotype showed superior performance uniformly across all the test environments, some genotypes were identified with better mean grain yield performance. Among the tested genotypes, genotype G4 ranked the first at three of the seven environments, followed by G8, G9, G11 and G12 which ranked first at least in one of the environments as recommended by AMMI selection. Vertex genotypes including G1, G9, G11 and G12 in GGE polygon biplot were identified as winner genotypes for different sectors. Similarly, GGE ranking biplot identified four genotypes (G9, G4, G12 and G11) as high yielding, out of which G4 was the most stable, followed by G9 and G12 while G11 was less stable. Considering wider and /or specific

adaptation, farmers' positive feedback and response to cold stress, three genotypes (G4, G9 and G11) were identified as candidate varieties and verified for final evaluation. Consequently, genotypes G9 and G11 were officially approved for release by the national variety releasing committee for cultivation in cold prone potential rice growing agro ecologies of Ethiopia, predominantly in Fogera and similar areas. Genotypes G4 and G12 can be used as potential parental lines in our crossbreeding programs to make use of their merits.

ACKNOWLEDGMENTS

Rice research team at Fogera, Jimma and Assosa research centers are highly appreciated for the critical support during trial implementation. Authors also acknowledged Africa Rice Centre for providing rice germplasms and Ethiopian Institute of Agricultural Research for financial support.

REFERENCES

1. De Data, K.S., 1981. Principles and practices of rice production. John Wiler & Sons, Inc. New York, pp: 642.
2. De, G.M., J. Thomas, R. Liyanage, J.O. Lay and S. Basu, 2019. Cold tolerance response mechanisms revealed through comparative analysis of gene and protein expression in multiple rice genotypes. PLoS ONE 14: e0218019.
3. Islam, M.D., M.D. Rasel, S. Khanam and L. Hassan, 2020. Screaming of Rice (*Oryza sativa* L.) Genotypes for cold Tolerance at seedling and reproductive Stages base on Morpho- Physiological Markers and Genetic Diversity Analysis. International Journal of Natural Science, 10: 58.
4. Ranawake, A.L. and C. Nakamura, 2011. Cold tolerance of an inbred line population of rice (*Oryza sativa* L.) at different growth stages. Tropical Agricultural Research & Extension 14(2): 25-33.
5. Fujino, K., H. Sekiguchi, T. Sato, H. Kiuchi, Y. Nonoue, Y. Takeuchi, T. Ando, S.Y. Lin and M. Yano, 2004. Mapping of quantitative trait loci controlling low-temperature germinability in rice (*Oryza sativa* L.). Theor. Appl. Genet., 108: 794-799.
6. Lee, M., 2001. Low temperature tolerance in rice?: The Korean experience, In: Shu Fukai, Basnayake, J. (Eds.), Increased lowland rice production in the Mekong region ACIAR Proceedings 101. IRRI, Makati, Philippines.

7. Shimono, H., M. Okada, E. Kanda and I. Arakawa, 2007. Low temperature-induced sterility in rice: Evidence for the effects of temperature before panicle initiation. *Field Crops Research*, 101: 221-231. doi:10.1016/j.fcr.2006.11.010.
8. Sarita, P.K., R. Jhansi and R.C. Damodar, 2017. Cold tolerance in rice at seedling and reproductive stage. *Journal of Pharmacognosy and Phytochemistry*, 6(6): 984-988.
9. Jacobs, B.C. and C.J. Pearson, 1994. Cold damage and development of rice: a conceptual model. *Australian Journal of Experimental Agriculture*, 34: 917-919.
10. Renata, P.C., C.K.M. Sandra and C.F. Luiz, 2008. Inheritance of panicle exertion in rice. *Science Agriculture*, 65(5): 502-507.
11. Takahashi, N., 1984. Differentiation of ecotypes in *Oryza sativa* L. In: Takahashi N and Tsunoda S. (Ed.) *Biology of rice*. Tokyo: Japanese Scientific Society, pp: 31-67.
12. Dessie, A., T. Lakew, Z. Zewdu, B. Asrat, M. Atnaf, A. Berier, F.Woreda, H. Solomon, H. Kinfe and M. Bitew, 2020. Green Super Rice (*Oryza sativa* L.) Variety Evaluation Under Rain Fed Conditions in Ethiopia. *Journal of Rice Research*, 8: 229.
13. Zelalem, Z., D. Abebaw and A. Mulugeta, 2020. Evaluation of yield and yield components of lowland rice (*Oryza sativa* L.) varieties in the rainfed-lowland rice producing areas of Ethiopia. *Academic Research Journal Agricultural Research*, 7(7): 569-576.
14. Ogunbayo, S.A., M. Sie, D.K. Ojo, A.R. Popoola, O.A. Oduwaye, I.O. Daniel, K.A. Sanni, M.G. Akinwale, B. Toulou, A. Shittu, G.B. Gregorio and E.F. Mercado, 2014. Comparative performance of forty-eight rice genotypes in diverse environments using AMMI and GGE biplot analysis. *International Journal of Plant Breeding and Genetics*, 8(3): 139-152.
15. Gauch, H.G., 2006. *Statistical Analysis of Yield Trials by AMMI and GGE*. Crop and Soil Sciences, Cornell University, 519 Bradfield Hall, Ithaca, NY, 14853.
16. Zobel, R.W., M.J. Wright and H.G. Gauch, 1988. Statistical analysis of a yield trial. *Agronomy Journal*, 80: 388-393.
17. Yan, W., 2001. GGE biplot-A windows application for graphical analysis of multi-environment trial data and other types of two-way data. *Agronomy Journal*, 93: 1111-1118.
18. Yan, W. and M.S. Kang, 2003. *GGE-biplot Analysis: A graphical tool for breeders, geneticists and agronomists*. CRS press, Boca Raton, FL.
19. IRRI, 1996. *Standard Evaluation System for Rice*. Rice knowledge Bank. IRRI, Philippines.
20. SAS Institute Inc, 2002. Version 9.0. SAS Institute Inc., Cary, NC.
21. Yan, W., L.A. Hunt, Q. Sheng and Z. Szlavnic, 2000. Cultivar evaluation and mega-environment investigation based on the GGE biplot. *Crop Science*, 40: 597-605.
22. Bantayehu, M., J. Esmael and Y. Awoke, 2013. Additive Main Effect and Multiplicative Interaction Analysis and Clustering of Environments and Genotypes in Malting Barley. *African Journal of Agricultural Research*, 8(18): 1896-1904.
23. Yan, W. and L.A. Hunt, 2001. Interpretation of genotype x environment interactions: An Empirical Comparison. *Agronomy Journal*, 87: 121-126.
24. Bharat, T.J., A.K. Serial, R.P. Saharan and H.A. Harikesh, 2018. AMMI biplot analysis for stability in basmati rice (*Oryza sativa* L.) in different production systems. *Electronic Journal of Plant Breeding*, 9(2): 502-510.
25. Tekdal, S. and E. Kendal, 2018. AMMI Model to Assess Durum Wheat Genotypes in Multi-Environment Trials. *Journal of Agricultural Science Technology*, 20: 153-166.
26. Lingaiah, N., B.S. Chandra, V. Venkanna, D.K. Rukmini and Y. Hari, 2020. AMMI Bi plot analysis for genotype x environment interaction on yield in rice (*Oryza sativa* L.) genotypes. *Journal of Pharmacognosy and Phytochemistry*, 9(3): 1384-1388.
27. Yan, W. and N.A. Tinker, 2006. *Biplot analysis of multi-environment trial data: Principles and applications*. Canadian Journal of Plant Science, 86: 626-645.