

Effects of Drying Methods on Proximate Compositions of Catfish (*Clarias gariepinus*)

Ogbonnaya Chukwu and Ibrahim Mohammed Shaba

Agricultural and Bioresources Engineering Department,
Federal University of Technology, PMB 65, Minna, Nigeria

Abstract: The effects of two different drying methods (smoking kiln and electric oven) on proximate compositions of catfish (*Clarias gariepinus*) were determined. Purchased quantities of catfish were shared into two parts: one part was used to determine the proximate compositions of the raw fish and the other part was sub-divided into two; a part was dried using smoking kiln at a temperature range of 60°-70°C for 24 hours and the remaining one was dried using electric oven at a temperature range of 120° C for 30 minutes. The proximate compositions of the dried samples were determined. Mean moisture, protein, lipid, ash, fibre, carbohydrate, vitamin A, Potassium and Phosphorus contents of raw fish were 71.85±0.07, 19.51±0.18, 14.28±0.19, 33.06±0.04, 0.98±0.01, 5.48±0.02, $3.6 \times 10^{-4} \pm 10^{-5}$, $1.2 \times 10^{-4} \pm 10^{-5}$ and $1.2 \times 10^{-4} \pm 10^{-5}$ %, respectively, while the energy value was 4.89±0.06 J/100g and vitamin C content was $1.1 \times 10^{-3} \pm 10^{-4}$ mg/ml. The changes in moisture, protein, lipids, energy value, vitamin A and phosphorus contents were found to be significant for the two drying methods. Ash, fibre, carbohydrate, vitamin C and potassium contents showed no significant differences for the two drying methods used in this study. The results indicate that drying methods have effects on the proximate compositions of catfish. Electric oven drying is recommended for healthy eating and for longer shelf-life of dried fish.

Key words: Catfish • Drying methods • Electric oven • Proximate compositions • Smoking kiln

INTRODUCTION

Fish is a very important source of animal protein in the diets of man. Smoked or dried fish is a traditional part of the diet of a large section of the world's population. However, the gap between the demand and supply of fish is widening due to increase in population, poor postharvest handling, lack of processing and storage facilities and utilization of unconventional fish species. For instance, the estimated fish demand in Nigeria in 1994 was put at 1,139,833 tonnes based on the population figure of 94,986,044 and per capita consumption of 12.0kg which was considered globally adequate for normal and healthy growth. However, only 280,307 tonnes were produced, indicating a deficit of 94,705,737 [1].

Methods of drying and smoking fish vary between different countries and within the same country depending on the species of fish used and the type of product desired. The processor may use unsalted fish products ranging from less than 2% to over 20%. The fish

may be dehydrated to various degrees with moisture levels in the final product ranging from about 10% to 60%. Processing temperatures may range from less than 5°C to up to 120°C and processing times from half an hour to several months. The fish may be dried only or smoked only or there may be a combination of smoking and drying. In some countries the fish is boiled before being smoked and/or dried. Adding to this complexity, the fish species used as raw material may be fresh water or marine species and may range from very lean to fatty fishes and its condition from fresh to stale. This variation makes it difficult to arrive at general conclusions regarding processing effects of smoking and drying on protein quality and the proximate compositions of the final products.

The effects of different processing and cooking methods on nutritional compositions of different species of fish have been studied. Nutritive and organoleptic changes of Nigerian traditionally-processed freshwater fish species were studied by Afolabi *et al.* [2]. The effect

of traditional drying processes on the nutritional values of fish was studied by Eves and Brown [3]. Changes in chemical composition and nutritional quality of fried sardine (*Clupea pilchardus*) produced by frozen storage and microwave reheating were reported by Castrillon *et al.* [4]. Proximate and mineral compositions of dried salted roes of hake (*Merluccius merluccius*, L.) and ling (*Molva molva*, L.) were reported by Rodrigo *et al.* [5]. Proximate compositions of raw and cooked Thai freshwater and marine fish were studied by Puwastien *et al.* [6]. Experimental investigation on solar drying of fish using solar tunnel dryer was carried out by Bala and Mondol [7]. Effects of cooking methods on the proximate compositions and mineral contents of rainbow trout (*Oncorhynchus mykiss*) were studied by Gokoglu *et al.* [8]. Tao and Linchun [9] reported influences of hot air drying and microwave drying on nutritional and odorous properties of grass carp (*Ctenopharyngodon idellus*), while Turkkan [10] studied the effects of cooking methods on the proximate and fatty acid compositions of seabass (*Dicentrarchus labrax*, L. 1758).

It has been observed that different processing and drying methods have different effects on nutritional compositions of fish. This is because heating, freezing and exposure to high concentration of salt lead to chemical and physical changes and therefore digestibility is increased, due to protein denaturation protein, but the content of thermolabile compounds and polyunsaturated fatty acids is often reduced [3, 9, 11]. Therefore the qualities of fish dried using different methods cannot be the same. Also the shelf life of fish dried in an electrically-operated oven varies from that of fish dried using a smoking kiln. The objective of this study is to determine the proximate compositions of raw catfish and catfish dried using smoking kiln and electric oven so as to ascertain the effects of drying methods on proximate compositions of the fish.

MATERIALS AND METHODS

Twenty freshly harvested Catfish (*Clarias gariepinus*) were obtained from Fish Market in Minna, Nigeria. The mean weight and length of the fish were respectively 278.34±2.21g and 31.24±0.98 cm. The twenty fish were shared into two equal parts: ten were used to determine the proximate compositions of the raw fish and the other ten were divided into two, each part being five fish. Five fish were prepared (eviscerated, beheaded and washed) and dried using smoking kiln at a temperature

range of 60°-70°C for 24 hours and the remaining five were prepared and dried using electric oven at a temperature of 120°C for 30 minutes. After drying using the two methods, all fish for each drying method were homogenized using a kitchen blender.

Proximate Analysis: The proximate compositions were assayed as described by AOAC [12]. All chemicals used were of analytical grade and supplied by Sigma Co. (St. Louis, USA). Each analysis was carried out in triplicates.

Statistical Analysis: The design was completely randomized. Proximate composition analysis was replicated three times (n = 3). Results presented are mean values of each determination ± standard deviation (SD). Analysis of variance was performed by one-way ANOVA procedures (SPSS 11.0 for Windows). Differences between the mean values of the treatments were determined by the least significant difference (LSD) test and the significance was defined at p<0.05.

RESULTS AND DISCUSSION

The proximate compositions of raw and dried catfish are presented in Table 1. Each value is the mean±standard deviation of triplicate determinations. Raw samples presented low protein, potassium and phosphorus; intermediate lipids, fibre and vitamin A; high moisture and ash contents, similar to previous reported by Eyo [13]. Decrease of moisture content and increase of protein, vitamin A, potassium and phosphorus contents were the most prominent changes in catfish after drying. Moisture, protein, lipids, energy value, vitamin A and phosphorus contents increased significantly in processed samples ($P<0.05$). This trend is in agreement with the results by [9]. The significant increase in protein levels ($P<0.05$) in dried catfish, when compared with the raw fish, suggests that protein nitrogen was not lost during drying. This is in accordance with the findings of Puwastien *et al.* [6] Gokoglu *et al.* [8]. Tao and Linchun [9]. There was also a significant difference between smoking kiln drying and electric oven drying for the protein content.

After drying, there was a significant increase in lipids content. Electric oven-dried samples retained higher lipids content than smoking kiln-dried samples ($P<0.05$). This result indicates that the fat loss phenomenon was more intensive in the smoking kiln-dried fish than in electric oven-dried samples. Fat may exude with the moisture evaporation and extended heat treatment during smoking kiln drying seems to enhance this phenomenon.

Table 1: Proximate Compositions of Raw and Dried Catfish (*Clarias gariepinus*)

Proximate Compositions	Raw	Kiln-dried	Electric-dried
Moisture (%)	71.85±0.07 ^a	28.92±0.03 ^b	15.62±0.03 ^c
Protein (%)	19.51±0.18 ^a	53.10±0.14 ^b	67.21±0.16 ^c
Lipids (%)	14.28±0.19 ^a	21.20±0.14 ^b	29.60±0.14 ^c
Ash (%)	3.06±0.04 ^a	3.92±0.03 ^b	3.62±0.03 ^b
Fibre (%)	0.98±0.01 ^a	1.71±0.01 ^b	1.96±0.03 ^b
Carbohydrate (%)	5.48±0.02 ^a	2.78±0.03 ^b	3.84±0.06 ^b
Energy Value J/kg	4.89±0.06 ^a	4.14±0.03 ^b	6.24±0.03 ^c
Vitamin A (%)	0.00036±0.00001 ^a	0.00053±0.00003 ^b	0.00073±0.00003 ^c
Vitamin C (mg/ml)	0.0011±0.0001 ^a	0.0010±0.0001 ^a	0.0014±0.0004 ^a
Potassium (%)	0.00012±0.00001 ^a	0.00045±0.00001 ^b	0.00048±0.00001 ^b
Phosphorus (%)	0.00012±0.00001 ^a	0.00026±0.00003 ^b	0.00040±0.00003 ^c

Values in the same row followed by different letters are significantly different (P<0.05)

CONCLUSIONS

In general, there were significant influences of drying on proximate compositions of catfish. Lack of negative influence of the drying processes on the protein, lipids, ash, fibre, vitamins and mineral contents of catfish is of great practical importance, although drying resulted in a significant loss of energy value. These results show that different nutritional components of fish undergo different changes at elevated temperatures. However, electric oven drying could improve the protein quality and prevent lipid oxidation in catfish, as compared with the conventional smoking kiln drying. The electric oven-dried samples showed lower lipids loss and higher protein content than smoking kiln-dried samples. This research provides basic nutritional information on freshwater catfish, both raw and dried. The present study also provides a possible application of electric oven drying as an efficient drying process for fish in Nigeria and elsewhere.

REFERENCES

- Food and Agriculture Organization (FAO), 1999. World Production of Fish, Crustaceans and Molluscs by Major Fishing Areas. Fisheries Information and Statistics Unit (FIDI), Fisheries Department, FAO., Rome.
- Afolabi O.A., O.A. Arawomo and O.L. Oke, 1984. Quantity Changes of Nigeria Traditional Processed Freshwater Species I: Nutritive and Organoleptic Changes. J. Food Technol., 19: 333-340.
- Eves, A. and R. Brown, 1993. The Effect of Traditional Drying Processes on the Nutritional Values of Fish. Tropical Sci., 33: 183-189.
- Castrillon, A.M., P. Navarro and E. Álvarez-Pontes, 1997. Changes in Chemical Composition and Nutritional Quality of Fried Sardine (*Clupea pilchardus*) Produced by Frozen Storage and Microwave Reheating. J. Sci. Food and Agric., 75: 125-132.
- Rodrigo, J., G. Ros, J. Priago, C. Lopez and J. Ortuno, 1998. Proximate and Mineral Compositions of Dried Salted Roes of Hake (*Merluccius merluccius*, L.) and Ling (*Molva molva*, L.). Food Chem., 63 (2): 221-225.
- Puwastien, P., K. Judprasong, E. Kettwan, K. Vasanachitt, Y. Nakngamanong and L. Bhattacharjee, 1999. Proximate Composition of Raw and Cooked Thai Freshwater and Marine Fish. J. Food Composition and Analysis, 12: 9-16.
- Bala, B.K. and M.R.A. Mondol, 2001. Experimental Investigation on Solar Drying of Fish Using Solar Tunnel Dryer. Drying Technol., 19: 427-436.
- Gokoglu, N., P. Yerlikaya and E. Cengiz, 2004. Effects of Cooking Methods on the Proximate Composition and Mineral Contents of Rainbow Trout (*Oncorhynchus mykiss*). Food Chem., 84: 19-22.
- Tao, W. and M. Linchun, 2008. Influences of Hot Air Drying and Microwave Drying on Nutritional and Odorous Properties of Grass Carp (*Ctenopharyngodon idellus*) Fillets. Food Chem., 110 (3): 647-653.
- Turkkan, A.U., S. Cakli and B. Kilinc, 2008. Effects of Cooking Methods on the Proximate Composition and Fatty Acid Composition of Seabass (*Dicentrarchus labrax*, L. 1758). Food and Bioproducts Processing, 86: 163-166.
- Eyo, A.A., 2001. Fish Processing Technology in the Tropics. National Institute for Freshwater Fisheries Research (NIFFR), New Bussa, Nigeria, pp: 10-170.
- AOAC., 2005. Official Methods of Analysis (18th Edn.). Association of Official Analytical Chemists International, Maryland, USA.
- Eyo, A.A. 1998. Shelf-life of Moonfish (*Citharus citharus*) and Tunk Fish (*Mormyrus rume*) During Storage at Ambient Temperature and on Ice. FAO Fisheries Report No. 574, pp: 35-37.