

Improvement of Propagation by Hardwood Cuttings with or Without Using Plastic Tunnel in (*Hamelia patens*)

Mohy Eldeen Nour Eldaim Elgimabi

Biology Division, Faculty of Science,
Taif University, Sudia Arabia, P.O. Box 888, Zip Cod 21974

Abstract: The propagation of *Hamelia* plant which was known as one of types of difficult to root cuttings of ornamental plants, was used to study the effect of time of take the cuttings and humidity on rooting and growth of this cuttings. Stem cuttings of the plant were subjected to different treatments, comprising season of cutting (winter, summer or karif," the rainy season") and level of humidity (using plastic tunnels). The experiment was carried out under the normal nursery conditions to determine the best conditions for growth and development of stem cuttings of plant. The range of time for this experiment extended from 4/12/1999-16/11/2000 and A split plot design was used with three replicates. Time of cuttings (irrespective of the treatment covered or not covered, wit or without mist) had a significant effect on growth(1%).The winter season gave good growth compared to other seasons. Plastic tunnels with water mist gave the best rooting and vegetative growth followed by the cutting under plastic tunnels without mist, irrespective of season.The control treatment (not under plastic cover) gave the poorest results.

Key words: Cutting • Season • Plastic Tunnel • *Hamelia* plant

INTRODUCTION

It is generally accepted that most of perennial ornamental plants are multiplied and propagated by the use of vegetative propagation that comprise cuttings, layering or grafting. The use of cutting from stems, leaves, roots or terminal buds is considerable as the most commonly applied technique, due to its practicability and simplicity, specially in devolving countries.

It is well known that the success of rooting of woody stem cuttings, in the majority of ornamental plants and fruit trees depends mainly on the physiological stage of the mother plant [1], the time of planting of the cutting [2,3] and the type of growth regulators used [4].

The prevailing environmental conditions in the nursery i.e. light, temperature and humidity play an important role in rooting and succeseeding growth and development stages of cuttings [5-7].This may be related to changes in the indigenous plant growth regulators or carbohydrate conditions[1]. Several researchers have investigated the influence of planting time on the rooting of stem cuttings and plant growth such as Darwesh [3] on *Ficus retusa* "Hawaii"; Deng

Xiong *et al.* [8] on *Quisqualis indica* and Rowezak [4] on *Ficus benjamina* var. *exotica*.In general, thy showed that the plants gave the best results concerning the rooting and the vegetative growth when the cuttings were planted during Marsh and April. However, other species gave the best results when the cuttings were planted in July and August as reported by Yong Kweon and KiSun [9] on *Abeliophyllum distichum*.

On the other hand Pipattanawong *et al.* [10] study the effect of the plastic pavilions on the propagation of (*Ficus carica*) hardwood cuttings. The result indicate that using plastic pavilion increased the temperature which promotes early callus formation of buds and roots in the propagation of *Ficus* cuttings in cool area.

The *Hamelia* ornamental plants (*Hamelia patens*) related to the family Rubiaceae is considered as an important plant used in outdoor gardens in Khartoum, for its flower colors and its potential for other uses such as hedge plant or single plant in different shape.

The problem with this plant in that is difficult to propagate in the nursery under the dry conditions of Khartoum State, the render it rare and difficult to fined in most of the nurseries.

MATERIALS AND METHODS

The experiment was carried out in the nursery of the Mogran Family Park, in Khartoum-Sudan. The nursery is made of wood stands and covered with Bamboos in 10 x 10 cm spaces.

Stem cuttings (20 cm long each) from one year old stems of *Hamelia* plants growing in the garden of Mogran Family Park were taken during winter (4/12/1999), summer (7/3/2000) and Kharif, the rainy season (17/8/2000). The cuttings were planted in rectangular plastic containers filled with sand. Fifteen cuttings were planted in each of nine plastic in each treatment, as follows

1st Treatment: Cuttings planted in containers and without plastic cover and no water mist (control).

2nd Treatment: Cuttings planted in the containers under plastic tunnels (70 x 100 x 150 cm) and without water mist.

3rd Treatment: Cuttings planted in the containers, under plastic tunnels (70 x 100 x 150 cm) and with water mist.(once in the morning).

All the treatments were irrigated in the mid day, during the course of the experiment in the three seasons.

A split plot design was used, with three replications. The seasons were considered as the main plots and the plastic tunnel with the mist, the plastic tunnel without mist and the uncovered treatments were the subplots.

Growth parameters, that comprise; number and length of roots, number and length of leaves of the cuttings, in each treatment were measured every fifteen days using gentle digging, to uproot the cuttings. After washing the required readings were taken and the cuttings were replanted in their respective container.

The average temperature during the day was recorded during the course of the experiment.

RESULTS AND DISCUSSION

The maximum number of roots from the *Hamelia* cuttings, in the experiment was obtained from the treatments under the plastic tunnels, both with and without water mist in three seasons as compared with the control (Table 1). The winter season gave the maximum number of roots compared to the other two seasons (summer and Kharif). These results suggest the importance of planting of the *Hamelia* stem cuttings

Table 1: Effect of Time of Cuttings Taking and Humidity on the Average Number of Root in the Stem Cutting of *Hamelia* Plant

Treatments	Time of cuttings taking			Mean for seasons
	Winter	Summer	Karif	
1-without plastic tunnel (control)	2.2	2.1	1.1	1.8
2-plastic tunnel no mist	3.1	2.4	1.6	2.4
3- plastic tunnel +mist	3.8	3.1	1.6	2.8
Mean for treatments	3.0	2.5	1.4	2.3

Table 2: Effect of Time of Cuttings Taking and Humidity on the Average of Root length in the Stem Cutting of *Hamelia* Plant

Treatments	Time of cuttings taking			Mean of seasons
	Winter	Summer	Karif	
1-without plastic tunnel(control)	0.6	0.6	0.2	0.5
2-plastic tunnel no mist	2.5	2.1	0.2	1.6
3- plastic tunnel +mist	4.0	3.4	0.2	2.5
Mean of treatments	2.4	2.0	0.2	1.5

Table 3: Effect of Time of Cuttings Taking and Humidity on the Average Number of Leaves in the Stem Cutting of *Hamelia* Plant

Treatments	Time of cuttings taking			Mean of seasons
	Winter	Summer	Karif	
1-without plastic tunnel(control)	1.6	1.4	1.8	1.6
2-plastic tunnel no mist	2.9	2.5	2.2	2.4
3- plastic tunnel +mist	2.8	2.7	2.3	2.6
Mean of treatments	2.3	2.2	2.1	2.2

Table 4: Effect of Time of Cuttings Taking and Humidity on the Average of Leaves length in the Stem Cutting of *Hamelia* Plant

Treatments	Time of cuttings taking			Mean of seasons
	Winter	Summer	Karif	
1-without plastic tunnel(control)	0.2	0.3	0.3	0.3
2-plastic tunnel no mist	2.8	2.3	0.2	1.8
3- plastic tunnel +mist	2.9	2.2	2.0	1.8
Mean of treatments	2.0	1.6	0.2	1.3

under plastic tunnels in the nursery under the dry arid conditions of Khartoum State, since it will insure continuous high humidity and temperature microenvironment needed for the initiation for the roots [10-12] and thus maintain the humidity balance between the outside and inside of cutting [13]. These stimulant effects on root growth may be attributed to the low temperature and to increase in photosynthetic rate during winter months, which leads to produce more

carbohydrate content[14]. Similar results were obtained by Rowezak [4] on *Ficus retusa* who recorded an increase in root growth during January, which help the initiation of the mechanism needed for the complicated physiological activities between the time of cuttings taking and the roots initiation [15].

The length of roots of cuttings in the treatments under the plastic tunnels was significantly longer than that in the treatment without plastic cover (control), regardless of season (Table 2). This effect on root elongation under the plastic tunnels might be attributed to the temperature (about 25°C during the day), light and humidity which were prevailing under the plastic tunnels during winter and summer seasons in the nursery [16,1], whereas during Kharif season the light was not intense enough during the intermitant cloud cover prevailing during this growing season.

The same trend was observed for number of leaves and the length of the leaf, where the cuttings under the plastic cover gave the largest number of leaves and the longest leaves (Table 3 and 4) compared to the control treatment. The plastic tunnel with mist treatment gave the best number of leaves in winter compared with the other two seasons. These results emphasized, the positive relationship between the growth of leaves and humidity, which is similar to that obtained by Darwesh[3] on *Ficus retusa*, Whitecomb,[17] and Sutter and Hutze[18].

The above obtained results, emphasized the importance of humidity and time for cuttings taking to obtain successful plants from the hard to root cuttings, like *Hamelia* plant.

REFERENCES

1. Day, J.S. and B.R. Loveys, 1998. Propagation from cuttings of two woody ornamental Australian shrubs, *Boronia megastigma* and *Hypocalymma angustifolium*, Endl.(white myrtle).Austral J. Exper. Agric., 38: 201-206.
2. Hartmann, H.T. and F. Loreti, 1965. Seasonal variation in rooting of leafy olive cuttings under mist. Proc. American Soc. Hort. Sc., 87: 194 & 98.
3. Darwesh, R.S.S., 2000. Studies on propagation of *Ficus retusa* cv. Hawaii. M. Sc. Thesis, Faculty. Agric, Cairo Unvi, Egypt.
4. Rowezak, M.M.A., 2001. Response of some Ornamental plants to Treatment with Growth substances.M. Sc. Thesis, Fac. Agric., Cairo Univ., Egypt.
5. Hofferma, G.J., 1979. Humidity in controlled environment. Guide Lines for plant research.
6. Bunce, J.A., 1984. Effect of humidity on photosynthesis. Jour. Of Exp. Botany, 65: 1245-51.
7. Karaguzel, O., 1997. Studies on the propagation of bougainvilleas from cuttings. Ziraat Fakultesi Dergisi. Akeniz Universitesi., 10: 109-118.
8. Deng Xiong, C., C. BagPing, D. JianWen and C. MuLin, 2000. Cutting propagation techniques for *Quisqualis indica*. Journal of Zhejiang Forestry College, 17: 384-388.
9. Yong Kweon, Y. and K. KiSum, 1996. Seasonal variation in rooting ability, plant hormones, carbohydrate, nitrogen starch and soluble sugar contents in cutting of white forsythia (*Abeliophyllum distichum*, Nakai). J. Kor. Soc. Hort. Sci., 37: 554-560.
10. Pipattanawong, N., S. Tiwong, B. Thongyeon, R. Darak, P. Thamin and W. Techa, 2008. Improvement of propagation by hardwood cuttings with and without using plastic pavilions in fig (*Ficus carica* L). Kasetsart J(Nat. Sci) 42: 207-214.
11. Bose, J.K., D.P. M-Mondal and D.K. Prumink, 1973. Propagation of Ixora, Hibiscus and Jasminum cutting under mist. Progressive Hort., 5-43-50.
12. Al-Musoum, S.S., 1982. Plant and root length of pepper (*Capcicum annum* L) under various mulches at high temperature. M.Sc. thesis abs. Univ. Arizona.
13. Dahab, M.A., 1992. Production of ornamental plants.Cairo Press, pp: 6-8.
14. Devlin, R.M. and F.H. Witham, 1986. Plant Physiology. CBS Publishers and Distributors. Delhi. India.. 4th Ed. pp: 57, 292 & 381.
15. Rahman, H.M.A., K. Khokar and M.H. Laghari, 1991. Effect of season of rooting on the ability of cuttings of Guava (*Psidium guajava*). Indian Jour. Of Agric. Sci., 16: 404-406.
16. Kitani, U.M., 1984. Flowers and Ornamental plants (Book in Arabic) Egyptian Univ. Hose (6,7,117).
17. Whitcomb, C.E., 1983. Rooting of cuttings under wet tent. Hort. Proc., 32: 450-455.
18. Sutter, E.G. and M. Hutzel, 1984. Use of humidity tent and antitranspirants in acclimatization of tissue cultured plants in the green house. Hort. Sci., 23(4): 450-455.