

## Effect of Different Levels of Bioneema and Nitrogenous Fertilizer on Bio Chemical Constituents in Tomato (*Lycopersicon esculentum* Mill.) C.V. Money Maker

<sup>1</sup>F.M. Prasad, <sup>2</sup>A.M. Lal, <sup>1</sup>A.A. Jilani, <sup>1</sup>J.P. Singh, <sup>3</sup>R. Arnold, <sup>2</sup>V.M. Prasad and <sup>1</sup>R.R. Singh

<sup>1</sup>Department of Chemistry, School of Chemical Sciences, St. John's College, Agra, India

<sup>2</sup>Faculty of Agriculture, Chemistry Allahabad Agricultural Institute, Allahabad, India

<sup>3</sup>Department of Environmental Sciences, Govt. T.R.S. College, Rewa, M.P., India

**Abstract:** A field experiment was conducted at AAI Allahabad during the winter season to study the effect of different levels of Bioneema and nitrogenous fertilizers on chlorophyll, phenol, ascorbic acid, acidity, total soluble sugar, reducing sugar lycopene and carotene, contents in tomato (*Lycopersicon esculentum* Mill.). It is evident from the results that there is no particular treatment, which has increased all the parameters simultaneously. The highest chlorophyll content recorded at B<sub>2</sub>N<sub>1</sub> (225.76 mg LG<sup>1</sup>) while total phenol found to be 280.0 mg/100 g when bioneema was not used. The ascorbic acids acidity total soluble sugar enhanced at B<sub>1</sub>N<sub>2</sub> to the extent of 32.78 mg/100 g, 1.96, 6.2 and 7.00% respectively. The lycopene and carotene content remarkably increased at B<sub>2</sub>N<sub>2</sub> over control.

**Key words:** Bioneema % urea % lycopersicon esculentum % constituents

### INTRODUCTION

Phenolic compounds have been shown to play a key in the resistance of plant to diseases causing microorganisms. In addition Phenolic may also serve as protectants against leaves eating insect, increases in phenolic content in host plant and can also be brought about some organic amendment in the soil of seed treatments. The alkaloid nimbodin and thionemone in neem tree is reported to have antinematic action against nematode. These several reports on the use of neem and its products against various diseases alarmed me to do research on tomato crop along with urea as the nitrogen source.

### MATERIALS AND METHODS

The experiment was conducted at the research plot of agricultural biochemistry department during the winter season. The experimental soil was of alluvial in nature having available N 172 kg haG<sup>1</sup>, P 14 kg haG<sup>1</sup>, available K 218 kg haG<sup>1</sup>, OC 0.44% and soil pH 7.6. The factorial RBD designed was adopted. The levels of Bioneema were applied at the rate of 2 g potG<sup>1</sup>, 4 g potG<sup>1</sup> and 6 g potG<sup>1</sup> while N through urea as incorporated @ of 10.86 g mG<sup>2</sup> potG<sup>1</sup> and 22.174 g mG<sup>2</sup> potG<sup>1</sup>. The chlorophyll

content was extracted from the leaves with cold 80% acetone v/v and estimated spectrophotometrically as suggested by Arnon [1]. Total phenol was estimated by following Folin Ciocalteu Reagent (FCR) method given by Bray *et al.* [2], TSS% by hand refractometer, carotene and lycopene by HPLC method. Ascorbic acid and reducing sugar were analyzed by titrimetric method.

### RESULTS AND DISCUSSION

It is evident from the results presented in (Table 1) indicated that the highest chlorophyll content appreciably increased at B<sub>2</sub>N<sub>1</sub> level whereas the lowest chlorophyll content declined to the extent of 188.522 mg LG<sup>1</sup> at B<sub>2</sub>N<sub>0</sub> where no fertilizers were incorporated. The phenol content recorded maximum at B<sub>0</sub>N<sub>1</sub> (255.83 mg/100 g) but lowest phenol content declined in control because of the non-availability of N and bioneema to the plants. It is interesting to note that B<sub>1</sub>N<sub>2</sub> of bioneema and urea remarkably promoted the ascorbic acid content (32.78 mg/100 g) while it narrowed down at B<sub>0</sub>N<sub>1</sub> to the extent of 15.78 mg/100 g. Similarly the percent acidity in tomato fruits observed highest at B<sub>2</sub>N<sub>2</sub> level (1.96%) while B<sub>2</sub>N<sub>1</sub> recorded lowest % acidity (0.38). No particular trend in increasing and decreasing manner could be

**Corresponding Author:** Dr. F.M. Prasad, Department of Chemistry, School of Chemical Sciences, St. John's College, Agra, India

Table 1: Efficacy of different levels of bioneema and nitrogenous fertilizers on biochemical constituents in tomato

Treatments	Chlorophyll content (mg/100 g)	Total phenol (mg/100 g)	Ascorbic acid (mg/100 g)	Acidity (%)	Total soluble sugar (%)	Reducing sugar (%)	Carotene (mg/100 g)	Lycopene (mg/100 g)
B <sub>0</sub> N <sub>0</sub>	174.96	94.16	21.04	1.06	4.2	9.05	7.46	17.20
B <sub>0</sub> N <sub>1</sub>	182.48	255.83	15.78	0.95	5.2	10.90	10.32	19.52
B <sub>0</sub> N <sub>2</sub>	176.84	280.00	31.56	1.02	4.1	17.40	10.74	20.41
B <sub>1</sub> N <sub>0</sub>	129.81	231.66	23.67	1.20	3.3	18.50	11.31	20.67
B <sub>1</sub> N <sub>1</sub>	146.74	241.66	28.93	0.95	4.6	19.10	12.70	22.89
B <sub>1</sub> N <sub>2</sub>	191.89	170.83	32.78	1.42	6.2	7.00	15.40	23.77
B <sub>2</sub> N <sub>0</sub>	118.52	254.16	23.67	1.22	3.3	21.50	15.23	22.58
B <sub>2</sub> N <sub>1</sub>	225.76	109.16	26.30	0.38	4.1	12.00	17.20	24.27
B <sub>2</sub> N <sub>2</sub>	146.73	165.0	31.65	1.96	4.2	16.15	20.37	24.40
B <sub>3</sub> N <sub>0</sub>	141.09	142.50	28.93	1.19	4.7	24.50	13.18	23.70
B <sub>3</sub> N <sub>1</sub>	129.81	117.50	26.50	1.48	4.1	20.20	15.90	22.60
B <sub>3</sub> N <sub>2</sub>	122.28	145.83	18.44	0.78	5.4	25.50	12.60	17.84
CD at 5 %	7.45	8.54	4.827	23.74	50.40	18.96	1.897	0.3385

recorded. Like wise B<sub>1</sub>N<sub>2</sub> level of bioneema and urea appreciably increased the total soluble sugar (6.2%) in fruit juices but it declined at B<sub>1</sub>N<sub>0</sub> due to non-availability of N. In case of reducing sugar it tremendously increased with the rise in Bioneema and nitrogen at B<sub>3</sub>N<sub>2</sub> level (25.50%) where as B<sub>1</sub>N<sub>2</sub> level did not show any appreciable effect on reducing sugar. The lycopene and carotene contents recorded to be 24.40 and 20.37 mg/100 g, which is highest among other treatments over control. The increased in lycopene and carotene contents may be due to the biosynthesis of chlorophyll in to lycopene, which is affected by varying intensity of light and temperature. It may concluded from the results that bioneema (Neem product) @ 4 g plotG<sup>1</sup> and nitrogenous fertilizers (Urea) @ 21.74 g plotG<sup>1</sup> was found to be more economical and appropriate dose to enhances the biochemical parameter and reduced the ill effect of pathogens and nematodes in tomato plant which adversely influence these parameters. The above results are in conformity with Alam [3-5].

## REFERENCES

1. Arnon, D.I., 1949. Copper enzyme in isolated chloroplast, polyphenoloxidase in *Beta vulgaris*. Plant Physiol., 24: 1-15.
2. Bray, H.G. and W.Y. Thorpe, 1954. Analysis of phenolic compound of interest in metabolism. In Meth in Bioch. Ann., Glick, D. (Ed.). Interscience Pub. Inc., New York, pp: 27-52.
3. Alam, M.M., 1991. Control of plant parasitic nematodes with oil seed cakes on some vegetables in fruits. Bibliographic Citation, Pak. J. Nematol., 19: 89-92.
4. Bhandari, A.L., K.N. Sharma and R.S. Brar, 1993. Relative agronomic efficacy of neem extract coated urea V/S prilled urea for rise. Proceeding Worlds Neem Conference, pp: 91.
5. Boomathi, N., P. Sivasubramanian and S. Raguraman, 2006. Biological activity of cow excreta with neem seed kernel extract against *helicoverpa armigera* (Hubner), 14: 11-16.