

Important Cattle Ticks and Tick Born Haemoparasitic Disease in Ethiopia: A Review

Nejash Abdela

Jimma University College of Agriculture and Veterinary,
Medicine School of Veterinary Medicine, Jimma, Ethiopia

Abstract: Ethiopia has the largest livestock population in Africa, but the contribution for the economic aspect of the country is still lowest and disease can be considered as major constrain. Ticks are the most important ectoparasites of livestock in tropical and sub-tropical areas. Ethiopia is not exceptional and ticks are responsible for severe economic losses both through the direct effects of blood sucking and indirectly as vectors of pathogens and toxins. Feeding by large numbers of ticks causes reduction in live weight gain and anaemia among domestic animals, while tick bites also reduce the quality of hides. However, the major losses caused by ticks are due to the ability to transmit protozoan, rickettsial and viral diseases of livestock, which are of great economic importance world-wide. This review concerns with general aspects of tick biology, the taxonomy, pathogenic effects and methods for the control of ticks. Further, the paper highlight the status of ticks and tick borne haemoparasitic diseases in Ethiopia. Ticks belong to the suborder Ixodida, which contains a single super family, the Ixodoidea, which is divided into two major families, Argasidae (soft ticks) and Ixodidae (hard ticks) and the rare family Nuttalliellidae, with a single African species. The main tick genera found in domestic animals of Ethiopia are *Amblyomma*, *Hyalomma*, *Rhipicephalus*, *Haemaphysalis* and *Rhipicephalus* (*Boophilus*). Various breeds of cattle differ in their response to tick infestations. *Bos indicus* pure breeds and crossbreeds were reported to be more innately resistant than *Bos Taurus* breeds. The conventional method of controlling tick infestations in Ethiopia is application of acaricide, either by hand spraying, by hand dressing. Therefore, to minimize tick adverse effect appropriate and timely strategic control measures are crucial.

Key words: Acaricide • Argasidae • Ectoparasites • Ixodidae • Livestock

INTRODUCTION

Ethiopia is believed to have the largest livestock population in Africa. This livestock sector has been contributing considerable portion to the economy of the country and still promising to rally round the economic development of the country [1]. In Ethiopia, livestock production remains crucial and represents a major asset among resource-poor small holder farmers by providing milk, meat, skin, manure and traction force [2]. The contribution of livestock to the national economy particularly with regard to foreign currency earnings is through exploration of live animal, meat and skin and hides [3].

According to Onu and Shiferaw [4] Poor health and productivity of animal due to disease has considerably become the major stumbling block to the potential of

livestock industry. As Regasa *et al.* [5] indicated currently parasitism represents a major obstacle to development and utilization of animal resource. In Ethiopia ectoparasites in ruminant causes serious economic losses to small holder farmers, the tanning industry and country as a while through mortality of animals, decreased production, downgrading and rejection of skin and hide. From the ectoparasites, ticks are ranked as the most economically important of livestock in tropics including sub-Saharan Africa [6].

Eyo *et al.*, (2014) indicated that Ticks cause substantial losses in cattle production, in terms of diseases, reduced productivity and fertility and often death and are economically the most important ecto-parasites of cattle. Huruma *et al.* [8] showed that different ticks have different predilection sites on the host's body. Ticks suck blood; damage hides and skins

introduce toxins and predispose cattle to myiasis and dermatophilosis [9, 10]. Moreover, tick reduces body weight gains and milk yield, in addition to creating sites for secondary invasion by pathogenic organisms [9]. More significantly, ticks transmit diseases from infected cattle to healthy ones. Ticks transmit a greater variety of pathogenic micro-organisms than any other arthropod vector group and are among the most important vectors of diseases affecting animals [10].

Walker *et al.* [11] indicated ticks which are considered to be most important to health of domestic animal in Africa comprise about seven genera. Among these genera the main tick genera found in Ethiopia includes *Ambylomma*, sub genus *Rhipicephalus* (*Boophilus*), *Haemaphysalis*, *Hyalomma* and *Rhipicephalus*. According to Gebre *et al.* [12] the genus *Ambylomma* and *Rhipicephalus* are predominating in many parts of country, *Hyalomma* and sub genus *Rhipicephalus* (*Boophilus*) also have significant role. Due to economic and veterinary importance of ticks, their control and transmission of tick born diseases remain challenge for the cattle industry of the world and it is a priority for many countries in tropical and subtropical regions [13, 14]. As Tadesse and Sultan, [15] indicated In Ethiopia there are about 47 species of ticks found on livestock and most of them have importance as vector and disease causing agent and also have damaging effect on skin and hide production. Therefore the objective of this paper is to review available literature on tick biology, the taxonomy, pathogenic effects and methods for the control of ticks and highlighting status of ticks and tick borne haemoparasitic diseases in Ethiopia.

Literature Review

Classification of Ticks: Ticks are within a member called the phylum (Arthropoda), class (Arachnida), sub class (Acari) and Order (Parasitiformes) [16]. Within the Parasitiformes, ticks belong to the suborder Ixodida, which contains a single super family, the Ixodoidea, which is divided into two major families, Argasidae (soft ticks) and Ixodidae (hard ticks) and the rare family Nuttalliellidae, with a single African species [17]. Accordant to Jongejan and Uilenberg, [18], the family Ixodidae, or hard ticks, contains some 683 species. As adults, Ixodids exhibit prominent sexual dimorphism: the scutum covers the entire dorsum in males, but in females (and immatures) the scutum is reduced to a small podonotal shield behind the capitulum, thereby permitting great distention of the idiosomal integument during feeding [19]. Ixodidae ticks are relatively large and

comprise thirteen genera. Seven of these genera contain species of veterinary and medical importance: *Amblyomma*, sub genus *Rhi.* (*Boophilus*), *Rhipicephalus*, *Haemaphysalis*, *Hyalomma*, *Dermacentor* and *Ixodes* [17]. Adult argasids lack a dorsal sclerotized plate or scutum, their integument is leathery and wrinkled, their mouthparts are not visible from above and they show no obvious sexual dimorphism. Argasidae are wandering ticks, which only remain on their host while feeding [20]. According to Latif and Walker, [21]. The family Argasidae, or soft ticks, consists of about 185 species worldwide and have one important genus that infests cattle, *Ornithodoros*. Epidemiology of ticks

Host Relationship: Some ticks live in open environments and crawl onto vegetation to wait for their hosts to pass by. This is a type of ambush and the behavior of waiting on vegetation of is called questing. Thus in genera such as *Rhipicephalus*, *Haemaphysalis* and *Ixodes* the larvae, nymphs and adults will quest on vegetation. The tick grabs onto the host using their front legs and crawl over the skin to find a suitable place to attach and feed. Adult tick of genera *Ambylomma* and *Hyalomma* are active hunters, they run across the ground after nearby hosts [11]

Attachment Site: Tick attachment site specificity is one of the populations limiting system that operate through the restriction of tick species to certain parts of the host body. They seek out places on the hosts where they are protected and have favorable conditions for their development [22]. Huruma *et al.* [8] indicated that different ticks have different predilection sites on the host's body.

The favorable predilection sites for *B. decoloratus* was the lateral and ventral side of the animal; *A. variegatum*, teat and scrotum; *A. coherence* udder and *H. truncatum*, scrotum and brisket and *H. marginatum rufipes* udder and scrotum, *R. evertsi evertsi* under tail and anus and *R. preaxatus* anus and under tail [8]. Depending on the tick, site preference on the host depends on the accessibility for attachment, to get blood and protection to overcome the environment damage that inhibits its existence and grooming activity of the host. Tick location on the host is lined to the possibility of penetration by hypostome. Genera with short hypostome for example *Rhipicephalus*, *Dermacentor* and *Haemaphysalis* species usually attach to hairless area such as undertail and anovulval area [8].

Life Cycle: In the hard ticks mating takes place on the host, except with *Ixodes* where it may also occur when the ticks are still on the vegetation. Male ticks remain on the host and will attempt to mate with many females whilst they are feeding. They transfer a sac of sperm (spermatheca) to the female. The females mate only once, before they are ready to engorge fully with blood. When they finally engorge they detach from the host and have enough sperm stored to fertilize all their eggs. Female hard ticks lay many eggs (2,000 to 20,000) in a single batch. Female argasid ticks lay repeated small batches of eggs. Eggs of all ticks are laid in the physical environment, never on the host [23].

Members of the family Ixodidae undergo either one-host, two-host or three-host life cycles. During the one-host life cycle, ticks remain on the same host for the larval, nymphal and adult stages, only leaving the host prior to laying eggs. During the two-host life cycle, the tick molts from larva to nymph on the first host, but will leave the host between the nymphal and adult stages. The second host may be the same individual as the first host, the same species, or even a second species. Most ticks of public health importance undergo the three-host life cycle. The three hosts are not always the same species, but may be the same species, or even the same individual, depending on host availability for the tick. Argasid ticks have two or more nymphal stages, each requiring a blood meal from a host. Unlike the Ixodidae ticks, which stay attached to their hosts for up to several days while feeding, argasid ticks are adapted to feeding rapidly (about an hour) and then promptly leaving the host [11].

All feedings of ticks at each stage of the life cycle are parasitic. For feeding, they use a combination of cutting mouthparts for penetrating the skin and often an adhesive (cement) secreted from the saliva for attachment. The ticks feed on the blood and lymph released into this lesion. All ticks orient to potential hosts in response to products of respiration [16].

Pathogenic Role of Ticks: Direct effects of ticks on cattle are tick worry, blood loss, damage to hides and skins of animals and introduction of toxins [9]. The ecology and physiology of ticks have made them second most important vectors after mosquitoes. Ticks transmit a large variety of intercellular bacteria in the *Rickettsia* group like *Rickettsia*, *Ehrlichia* and *Anaplasma*. Similarly several piroplasm protozoa like *T. annulata*, *T. parva* and *Babesia bigemina* are also transmitted specifically by

ticks [24, 25]. Hard ticks (Acari: Ixodidae) are obligate hematophagous ectoparasites and important vectors of viruses, bacteria and protozoa. They are considered second only to mosquitoes as the most medically important group of arthropods [26]. Tick worry is a generalized state of unease and irritability of cattle severely infested with ticks, often leading to serious loss of energy and weight. This negative effect on the growth of animals and their production is thought to be due to the effects of a toxin in the saliva of ticks [9].

Anaemia is another inevitable consequence of heavy infestation by any blood-feeding parasite and cattle deaths attributable to anaemia as a result of tick infestation are common. Engorging Ixodidae females will increase their weight by 100–200 times but the actual amount of blood ingested is much greater than this, as blood meal is concentrated and fluid excreted in saliva. Estimates of the amount of blood removed vary according to the species under consideration [9]. Tick saliva contains toxins which have a specific pathogenic effect. The toxins affect not only the attachment site but also the entire organs of the host. Some ticks produce neurotropic toxins which induce tick paralysis that is characterized by an acute ascending flaccid motor paralysis. Females of the species *Hyalomma truncatum* produce a dermatropic (epitheliotropic) toxin which causes sweating sickness in calves and some adult cattle [26].

Ticks Control Methods: The aim of tick control campaign is not to control all ticks simultaneously, but a definite species because of its particular role [28]. The successful implementation of rational and sustainable tick control programmes in grazing animals is dependent upon a sound knowledge of the ecology or epidemiology of the tick as it interacts with the host in specific climatic, management and production environments. In most situations, however, efficient and reliable methods for the control of cattle ticks and TBD are based on the use of a chemical treatment (acaricide application), often without a local understanding of appropriate ecology or epidemiology [29].

The availability of each of these options, their advantages and disadvantages and the cost benefit of each alternative strategy should be assessed before deciding on a control programme [30]. Ideally, strategies should target the parasitic and free-living phases of the life cycle and the role of the ticks in the transmission of Tick-borne diseases should not be neglected [12]. It is now generally understood that tick control should not

only be based on acaricide use, despite the fact that this remains the most efficient and reliable single method. Complementary approaches have been developed and are being researched to enable integrated control strategies against the tick and its haemoparasites [29, 30]. The following are the most commonly used tick control methods.

Ecological Tick Control: Ecological control method is used for habitat and host linked treatment. Tick control in the habitat and vegetation requires modification of the plant cover by removal of vegetation that shelters ticks [30]. Pasture management, including spelling and seasonal changes in cattle grazing areas in Australia and in Zambia respectively has been used as a tick control strategy and are believed to be responsible for a decrease its burden [11].

Biological Tick Control: A first attempt at tick biocontrol was made with the introduction of tick parasitic wasps from France to the USA and Russia. During the past decades, interest in developing antitick biocontrol agents such as birds, parasitoides, entomopathogenic nematodes, entomopathogenic fungi and bacteria have gained momentum [31]. In biological tick control the activities of the hyperparasites chalcid flies *Hunterellus* are probably important in nature, but they are difficult to evaluate and it is still more difficult to manipulate or reproduce them for practical use. The biological agents, which potentially include predators like rodents, birds, ants, spiders, lizards and beetles as well as Prasinid wasps (destroy the host: the wasp lay the eggs in the engorged ticks and larvae eats the tick and emerges as adult to attack another tick) and parasites (Nematodes and fungus) attack soil living stages of the ticks are effective and depending on the conditions, these predators can consume a large number of ticks. Yet, having such effective importance the development of a biological tick control methods has been neglected as compared to the control of plant pests or dipterous insects harmful to men and animals [30].

Chemical Tick Control: Acaricide treatments are commonly used in a suppressive approach, applying multiple treatments at regular intervals during the height of infestation. Suppressive treatments are the most effective in the short term; keeping animals almost tick free, thereby reducing the direct effect of the ticks and the risk of disease transmission. This procedure will, however,

select heavily for acaricide resistance in the ticks. An ideal acaricide would be cheap, easily applied, with a strong knock down effect and sufficient residual effect on female ticks to prevent egg laying and to protect cattle from reinfestation by larvae. It should not select for resistance through a prolonged, gradual decay on the animal (i.e. it should have a sharp cut off in efficacy with time). In addition, it should be non-toxic to livestock and humans and have no detectable residues in meat and milk. Unfortunately, such an ideal acaricide has not yet been produced. Generally, although the use of acaricides for the control of ticks has limitations and tick resistance to acaricides is an increasing problem and real economic threat to the livestock worldwide, most livestock holders depend completely on acaricides to control ticks, but do not have access to guidelines on how to make a profit from their tick control program or how to detect and resolve problems with resistance to acaricides [31].

Genetic Tick Control: The application of acaricides is the most common method used to control cattle ticks. However, the improper use of these chemicals compounds has been causing the development of tick resistance to various pesticides available in the market, reducing these products' useful lifetimes. Besides this, problems generated by the presence of chemical residues in meat, milk and the environment have prompted reflection on the need for better monitoring of their application [32]. Therefore, the study of the genetic resistance to ticks among different breeds of cattle can contribute to the development of alternative control methods [33]. It is widely known that *Bos indicus* cattle are more resistant to ectoparasites than are *Bos taurus* animals. There are great differences between these two breeds of cattle in regard to their susceptibility to parasitism by cattle ticks [34]. Studies are intensifying the crossing of these two groups, aiming to obtain animals that are more resistant to the conditions found in tropical countries and are also good meat producers [35].

The Distribution of Ticks in Ethiopia: The distribution and abundance of tick species infesting domestic ruminants in Ethiopia vary greatly from one area to another area [36]. The main tick genera found in domestic animals of Ethiopia are *Amblyomma*, *Hyalomma*, *Rhipicephalus*, *Haemaphysalis* and *Rhipicephalus* [36]. Among the genera *Rhipicephalus*, *Rhipicephalus lunulatus* species were observed in Central Ethiopia [37] and *Rhipicephalus muhasmae* in Borena [38], in wetter

western areas of the country [39]. Seyoum, [40] has recorded *Rh. humoralis*, *Rh. cliffordi*, *Rh. compositus* and *Rh. distinctus* in Wollo and Northeast areas. *Rhipicephalus evertsi evertsi*, “Red-legged tick” [11], is the most widespread species of *Rhipicephalus* [41]. Regassa [38] in Borena zone showed that *A. variegatum*, *A. gemma* and *A. lepidum* distributed in wider area of southern Ethiopia. *Amblyomma variegatum* and *A. cohaerens* are the two most prevalent *Amblyomma* species in Awassa areas in decreasing order [42] In eastern Ethiopia, *A. variegatum* and *A. gemma* are the two most widely spread species [43]. *Amblyomma gemma*, is also recorded in eastern and southern Ethiopia [39]. *Amblyomma variegatum* and *Amblyomma coherence* in was also recorded in Haramaya [44]. It is clearly associated with dry types of vegetation or semi-arid rangelands. *Amblyomma lepidum*, is most commonly inhabits arid habitats and in open bushed shrub or wooded grassland and its distributions overlap with *Amblyomma gemma* and that of *Amblyomma variegatum* [11] In Ethiopia, about eight species of *Hyalomma* that affect cattle are identified, which includes *Hyalomma marginatum rufipes*, *Hy. dromedarii*, *Hy. tuncatum*, *Hy.m. marginatum*, *Hy. impelatum*, *Hy. anatolicum excavatum*, *Hy. anatolicum anatolicum* and *Hy. albiparmatum* [44].

Two species of *Rhipicephalus* (*Boophilus*) sub genus are known to exist in Ethiopia, which include *Rhipicephalus* (*Boophilus*) *decoloratus* and *Rhipicephalus* (*Boophilus*) *annulatus*. The study done by Regassa [38] in Borena zone; Mekonnen *et al.*, [38] in central Ethiopia; Assefa [45] in Asella; Berhane [42] in Awassa; Dessie [46] in Asella; Seyoum [40] in Wollo and Asosa area Fantahun and Mohammed, [47] indicated the distribution of *Rhipicephalus* (*Boophilus*) *decoloratus*. *Rhipicephalus* (*Boophilus*) *annulatus* is known to present in Gambella region and recorded [39].

Tick Borne Diseases and Status of Tick Borne Haemoparasitic Diseases in Ethiopia

Tick Borne Diseases: The term vector-borne disease refers to any of a broad array of infectious diseases caused by pathogens that are transmitted by arthropods or other biologic intermediaries [48] Ticks and tick-borne diseases (TBDs) affect the productivity of bovines and leads to a significant adverse impact on the livelihoods of resource-poor farming communities [49]. Four main TBDs, namely anaplasmosis, babesiosis, theileriosis and cowdriosis (heartwater) are considered to be the most

important tick-borne diseases (TBDs) of livestock in sub-Saharan Africa, resulting in extensive economic losses to farmers in endemic areas [50]. They are responsible for high morbidity and mortality resulting in decreased production of meat, milk and other livestock by-products [51]. These diseases generally affect the blood and/or lymphatic system and cause fever, anaemia, jaundice, anorexia, weight loss, milk drop, malaise, swelling of lymph node, dyspnoea, diarrhoea, nervous disorders and even death. Major cattle tick-borne diseases in Ethiopia are anaplasmosis, babesiosis, theileriosis [52] and Dermatophilosis [53]. Besides to disease transmission ticks inflict a huge economic loss. Production losses due to ticks and tick-borne diseases around the globe have been estimated at US\$ 13.9 to US\$ 18.7 billion annually leaving world’s 80% cattle at risk [54]. Bekele [43] estimated an annual loss of US\$ 500,000 from hide and skin downgrading from ticks and approximately 65.5% of major defects of hides in eastern Ethiopia are from ticks. Furthermore, the costs associated with maintaining chemical control of ticks in tropical and subtropical regions of the world have been estimated at US\$ 25.00 per head of cattle per year [55].

Status of Tick Borne Haemoparasitic Diseases in Ethiopia:

Similar to other countries, there are a considerable number of economically important livestock diseases occurring in Ethiopia. Among others, tick borne haemoparasitic diseases are of the major constraints to the livestock industry of the country. Sileshi [56] indicated the existence of *Anaplasmosis*, *Babesiosis*, *Cowdriosis* and *Theileriosis* (*T. mutans*), but their significance in terms of mortality and productive losses and the degree of enzootic stability are not yet very well known. There are no clinical or serological reports of the presence of either *T. parva* in Ethiopia. But, there is relatively uncontrolled movement of livestock from Sudan and Kenya, where these diseases and their vectors are found [56]. In previous studies, conducted by Mekonin *et al.*, (1992) *B. bovis*, *T. orientalis* and *T. velifera* were reported from Gambella region, western Ethiopia. Gebrekidan *et al.* [57] reported a widespread distribution of *Theileria* spp. among domestic ruminants in northern Ethiopia. Furthermore the study by Gebrekidan *et al.* [57] reported the presence *T. annulata*, the cause of tropical theileriosis, in Ethiopia for the first time. Table 1 indicates existing TB haemoparasitic diseases in four regions of Ethiopia.

Table 1: existing Tick-borne haemoparasitic diseases in four regions of Ethiopia

Region	TBDs	Samples	Diagnostic Tests	Sources
Amhara	<i>Babesia species</i>	Blood smear	Microscopic	[40]
	<i>B. bigemina</i>	Lymph node impression		
	<i>Theileria mutans</i>			[58]
SNNPRS	<i>Babesia species</i>	Blood smear	Microscopic	[59]
	<i>Theileria species</i>	Lymph node impression		
Central Ethiopia	<i>abesia species</i>	Blood smear	Microscopic	
Oromia	<i>Babesia species</i>	Serum	Microscopic	[60]
	<i>B. bigemina</i>	Blood smear	Serology (ELISA)	
	<i>Theileria mutans</i>			

CONCLUSIONS

Ticks are obligate blood feeding ectoparasites of vertebrates and induce huge production loss in livestock industry and creating serious public health problems in the world. The main tick genera found in Ethiopia are *Amblyomma*, *Haemaphysalis*, *Hyalomma* and *Rhipicephalus*. Tick-borne diseases of cattle such as anaplasmosis, babesiosis, ehrlichiosis and theileriosis (*T. mutans*) are present in Ethiopia. Heavy infestations by different tick species suppress the immunity of cattle and also damage teats and reduce productivity of animals and there are direct effects associated with tick infestation that leads to tick worry, anorexia and anemia. These all are the impacts of tick infestation so, to minimize tick impact appropriate and timely strategic control measures are crucial. The conventional method of controlling tick infestations in Ethiopia is application of acaricide, either by hand spraying, by hand dressing. The availability of vaccine is very small. The ability to induce an effective, sustained immunological response is crucial but needs improvement. Problems of acaricide resistance, chemical residues in food and the environment and the unsuitability of tick resistant cattle for all production systems make the current situation unsatisfactory and require the development of absolute control through effective vaccine. Therefore, in line with the above conclusions; the following recommendations were forwarded:

- The government should monitor the use of potentially dangerous chemicals and conserve foreign exchange.
- Intensive acaricide application to control ticks has a number of limitations, Therefore, immunization together with strategic tick control are recommended for exotic and crossbred cattle
- Research should be conducted on tick species and their epidemiology for the continuous understanding of improved control strategies

- Awareness should be given to animal breeder on problem of tick and TBD and different control method.

ACKNOWLEDGEMENTS

Above all, I would like to praise my Almighty God, Allah, for supporting me health, wisdom and strength in my work and for his perfect protection and guidance of my life. I would like to express my sincere thanks and best regards to my beloved and respected family for their invaluable help and encouragement during my journey for their moral and financial support throughout my entire academic career. Finally I would like to thanks all my friends those who helped me for my successfulness.

REFERENCES

1. CSA., 2013. Federal Democratic Republic of Ethiopia, Central Statistical Authority, Agricultural sample survey (2012/2013), Report on livestock and livestock characteristics (Privet and Peasant Holdings), Addis Ababa, pp: 9-20.
2. Mesfin, T. and M. Lemma, 2001. The role of traditional veterinary herbal medicine and its constraints in the animal health care system in Ethiopia. In: Conservation and Sustainable Use of Medicinal Plants in Ethiopia. Z. Medhin and D. Abebe, (Eds): Institute of Biodiversity conservation and Research. Addis Ababa, Ethiopia, pp: 22-28.
3. MoARD, 2008. The effect of skin and hide quality on domestic and export market and evaluation of the campaign against ectoparasites of sheep and goat in Amhara, Tigray and Afar region, official report to Region and other sectors, Addis Ababa, Ethiopia.
4. Onu, S.H. and T.Z. Shiferaw, 2013. Prevalence of ectoparasite infestations of cattle in Bench Maji zone, southwest Ethiopia. *Veterinary World*, 6(6): 291-294.

5. Regasa, T.D., A. Kebede Tsegay and H. Waktole, 2015. Prevalence of major ectoparasites of calves and associated risk factors in and around Bishoftu town. African Journal of Agricultural Research, 10(10): 1127-1135.
6. Lorusso, V., K. Picozzi, B.M. de Bronsvort, A. Majekodunmi, C. Dongkum, G. Balak and S.C. Welburn, 2013. Ixodid ticks of traditionally managed cattle in central Nigeria: where *Rhipicephalus* (*Boophilus*) *microplus*. Parasit Vectors, 6(6): 171.
7. Eyo, J.E., F.N. Ekeh, N. Ivoke, C.I. Atama, I.E. Onah, N.E. Ezenwaji and C.B. Ikele, 2014. Survey of Tick Infestation of Cattle at Four Selected Grazing Sites in the Tropics.
8. Huruma, G., M. Abdurhaman, S. Gebre and B. Deresa, 2015. Identification of tick species and their prevalence in and around Sebeta town, Ethiopia. J. Parasitol. Vect. Biol., 7(1): 1-8.
9. Marufu, M.C., 2008. Prevalence of Ticks and Tick-borne Diseases in Cattle on Communal Rangelands in the Highland Areas of the Eastern Cape Province, South Africa. Master of Science in Agriculture (Animal Science) in the Department of Livestock and Pasture Science Faculty of Science and Agriculture, pp: 1-134.
10. Yiwombe, K., 2013. An investigation to detect *Boophilus* Tick (Blue Tick) T (Doctoral dissertation, Midlands State University).
11. Walker, A.R., A. Bouattour, J.L. Camicas, A. Estrada-Pena, I.G. Horak, A.A. Latif, R.G. Pegram and P.M. Preston, 2003. Ticks of domestic animals in Africa: a guide to identification of species, Bioscience report, pp: 1-221.
12. Gebre, S., M. Nigist and B. Kassa, 2001. Seasonal variation of ticks on calves at Sebeta in western Shewa Zone. Ethiop. Vet. J., 7(1-2): 17-30.
13. Rajput, Z.I., S.H. Hu, W.J. Chen, A.G. Arijo and C.W. Xiao, 2006. Importance of ticks and their chemical and immunological control in livestock. Journal of Zhejiang University Science B, 7(11): 912-921.
14. Salih, D.A., Hussein A.M. El and L.D. Singla, 2015. Diagnostic approaches for tick-borne haemoparasitic diseases in livestock. Journal of Veterinary Medicine and Animal Health, 7(2): 45-56.
15. Tadesse, B. and A. Sultan, 2014. Prevalence and distribution of tick infestation on cattle at Fitchelale, North Shewa, Ethiopia. Livest. Res. Rural Dev, 26(8).
16. Torres, D.F., 2008. The brown dog tick *Rhipicephalus sanguinus* (Acari: Ixodidae): From taxonomy to control. Veterinary Parasitology, 152(9): 173-185.
17. Rodriguez-Vivas, R.I., M.Y. Mata, G.E. Pérez and W. Wagner, 2004. The effect of management factors on the seroprevalence of *Anaplasma marginale* in *Bos indicus* cattle in the Mexican tropics. Trop Anim Health Prod., 36: 135-143.
18. Jongejan, F. and G. Uilenberg, 2004. The global importance of ticks. Parasitology, 129(S1): S3-S14.
19. Lora, R.B., 2001. Veterinary Parasitology. The Practical Veterinaria, Arthropods. Butterworth-Heinemann, A member of the Reed Elsevier group, Library of Congress Cataloging, United State of America, pp: 16-21.
20. Barker, S.C. and A. Murrell, 2004. Systematic and evaluation of ticks with a list of valid genus and species names. J. Parasitol., 129(7): 15-36.
21. Latif, A.A. and A.R. Walker, 2004. An introduction to the biology and control of ticks in Africa. ICTTD-2 project, pp: 1-29.
22. Sathaporn, J.A., J.A. Weeraphol, O. Omar, B. Barriga and W. Roger, 2004. Reduced Incidence of *Babesia bigemina* Infection in Cattle Immunized against the Cattle Tick, *Boophilus microplus*. Department of Parasitology. FVM, Kasetsart University, Bangkok, Thailand, pp: 36-47.
23. Charles, M.H. and E.D. Robinson, 2006. Diagnostic Parasitology for Veterinary technicians 3rd edition, pp: 192-195.
24. Sharma, A., L.D. Singla, A. Tuli, P. Kaur, B.K. Batth, M. Javed and P.D. Juyal, 2013. Molecular prevalence of *Babesia bigemina* and *Trypanosoma evansi* in dairy animals from Punjab, India by duplex PCR: A step forward to detection and management of concurrent latent infections. Biomed Research International Volume 2013, Article ID 893862, pp: 8.
25. Tuli, A., L.D. Singla, A. Sharma, M.S. Bal, G. Filia and P. Kaur, 2015. Molecular epidemiology, risk factors and haematochemical alterations induced by *Theileria annulata* in dairy animals of Punjab (India). Acta Parasitologica, 60(3): 378-90.
26. Ali, Z., 2010. Immunoprophylaxis Of Tick Infestation In Bovine (Doctoral dissertation, University of Veterinary and Animal Sciences, Lahore).
27. Kahn, C.M., 2006. The Merck Veterinary Manual, 9th Edition, Merck and Co. Inc, Whitehouse Station, USA.

28. FAO., 2004. "Acaricide resistance: diagnosis, management and prevention," in Guidelines Resistance Management and Integrated Parasite Control in Ruminants, Animal Production and Health Division, Agriculture Department, Food and Agriculture Organization of the United Nations, Rome, Italy, pp: 25-77.
29. Alanr, W., 2011. Eradication and control of livestock ticks: biological, economic and social perspectives. Royal (Dick) School of Veterinary Studies, University of Edinburgh, Summerhall Place, Edinburgh EH9 1QH, UK, pp: 236-253.
30. Kirby, C., 2010. Tick Management Hand book. Biological tick Control, 2nd Edit. TheConnecticut Agricultural Experimentation Station, New Haven, pp: 70-71.
31. George, J.E., J.M. Pound and R.B. Davey, 2008. Acaricides for controlling ticks on cattle and the problem of acaricide resistance. In Ticks: Biology, Disease and Contro. Cambridge University Press, UK, pp: 408-423.
32. Castro-Janer, E., J.R. Martins, M.C. Mendes, A. Namindome, G.M. Klafke and T.T.S. Schumaker, 2010). Diagnoses of fipronil resistance in Brazilian cattle ticks *Rhipicephalus (Boophilus) microplus* using *in vitro* larval bioassays. Vet. Parasitol., 173: 300-306.
33. Ibelli, A.M.G., A.R.B. Ribeiro, R. Giglioti, L.C.A. Regitanod, M.M. Alencard, A.C.S. Chagas, A.L. Pac, H.N. Oliveirac, J.M.S. Duarte and M.C.S. Oliveirad, 2011. Resistance of cattle of various genetic groups to the tick *Rhi. (Boophilus) microplus* and the relationship with coat traits. Vet.Parasitol.,doi:10.1016/j.vetpar.2011.11.019.
34. Bianchin, I., J.B. Catto, A.N. Kichel, R.A.A. Torres and M.R. Honer, 2007. The effect of the control of endo- and ectoparasites on weight gains in crossbred cattle (*Bos Taurus taurus* ×*Bos taurus indicus*) in the central region of Brazil. Trop. Anim. Health. Prod., 39: 287-296.
35. Graf, J.F., R. Gogolewski, N. Leach-Bing, G.A. Sabatini, M.B. Molento, E.L. Bordin and G.J. Arantes, 2005. Tick control: an industry point of view. Parasitology, 129: S427-S442.
36. Desalegn, T., A. Fikru and S. Kasaye, 2015. Survey of Tick Infestation in Domestic Ruminants of Haramaya District, Eastern Hararghe, Ethiopia. Journal of Bacteriology and Parasitology.
37. Mekonnen, S., I. Hussein and B. Bedane, 2001. The distribution of ixodid ticks (Acari: Ixodidae) in central Ethiopia. The Onderstepoort journal of veterinary research, 68(4): 243.
38. Regassa, A., 2001. Tick infestation of Borana cattle in the Borana Province of Ethiopia. Onderstepoort Journal of Veterinary Research, 68(8): 41-45.
39. de Castro, J.J., 1994. A survey of the tick species in western Ethiopia including the previous findings and recommendation, for further tick survey in Ethiopia. Technical report AG: DP/ETH/83/023, FAO. Rome, pp: 1-83.
40. Seyoum, Z., 2001. Study of ticks and tick-borne diseases on cattle at Girana valley in the North Wollo Zone. Proceeding of the Ethiopian Veterinary Association, Vol. 15.
41. Gebre, S., S. Mekonnen, P. Kaaya Godwin, T. Tekle and Y. Jobre, 2004. Prevalence of ixodid ticks and trypanosomosis in camels in southern rangelands of Ethiopia.
42. Berhane, M., 2004. Distribution of livestock tick species in Awassa area. DVM thesis, AAU, FVM, Debre-Zeit, pp: 1-16.
43. Bekele, T., 2002. Studies on seasonal dynamics of ticks of Ogaden cattle and individual variation in resistance to ticks in Eastern Ethiopia. J. Vet. Med. 49: 285-288.
44. Bedasso, M., B. Abebe and H. Degefu, 2014. Species composition, prevalence and seasonal variations of Ixodidae cattle ticks in and around Haramaya town, Ethiopia Full Length Research Paper, 6(5): 131-137.
45. Assefa, B., 2004. A survey of ticks and tick-borne blood protista in cattle at Asela, Arsi Zone. DVM thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, pp: 25-36.
46. Dessie, S., 2005. Cattle Tick Dynamics in Different Agro-Ecological Zones of Wolayta, Southern Ethiopia. Master thesis, Addis Ababa University, Faculty of Veterinary Medicine, Debre Zeit, Ethiopia, pp: 18-83.
47. Fantahun, B. and A. Mohamed, 2012. Survey on the Distribution of Tick Species in and Around Assosa Town, Ethiopia. Research Journal of Veterinary Sciences, 5: 32-41.
48. Bowman, D.D., 2009. Georgis Parasitology for veterinarian 9th editon. Newyork, pp: 1-465.
49. Jabbar, A., T. Abbas, Z.U.D. Sandhu, H.A. Saddiqi, M.F. Qamar and R.B. Gasser, 2015. Tick-borne diseases of bovines in Pakistan: major scope for future research and improved control. Parasit Vector, 8: 283.
50. Eygelaar, D., F. Jori, M. Mokopasetso, K.P. Sibeko, N.E. Collins, I. Vorster and M.C. Oosthuizen, 2015. Tick-borne haemoparasites in African buffalo (*Syncerus caffer*) from two wildlife areas in Northern Botswana. Parasites and vectors, 8(1): 1-11.

51. Simuunza, M.C., 2009. Differential Diagnosis of Tick-borne diseases and population genetic analysis of *Babesia bovis* and *Babesia bigemina* (PhD Thesis, University of Glasgow).
52. Mekonnen, S., J. de Castro, S. Gebre, I. Hussein and A. Regassa, 1992. Ticks, tick-borne diseases and their control in Western Ethiopia. *Int. J. Trop. Ins. Sci.*, 13: 661-664.
53. Mekonnen, S., 1996. Epidemiology of ticks and tick-borne diseases in Ethiopia: future research needs and priorities. In: Proceedings of a Workshop Held in Harare, Natl. Anim. Health Res. Centre, Ethiopia, 12-13 March.
54. Ghosh, S., P. Azhahianambia and M.P. Yadav, 2007. Upcoming and future strategies of tick control: a review. *J. Vect. Born. Dis.*, 44: 79-89.
55. Pegram, R.G., 2001. Getting a handle on tick control: A modern approach may be needed. *Vet. J.*, 161: 227-228.
56. Sileshi, M., 1994. Tick and TBDs Survey and control in Ethiopia. In T.T. Dolan and Musisi F-TBD control in Eastern, Central and Southern Africa. In: Proceeding of 16th conference of EVA, Addis Ababa, Ethiopia, pp: 120-151.
57. Gebrekidan, H., A. Hailub, A. Kassahunb, I. Rohousova, C. Maiac, D. Talmi-Frank, A. Warburge and G. Baneth, 2014. *Theileria* infection in domestic ruminants in northern Ethiopia. *Veterinary Parasitology*, 200(2): 31-38.
58. Jewaro, A., 1986. A survey of tick and tick born disease in Gamo Gofa administrative region DVM thesis. Faculty of Veterinary Medicine, Addis Ababa University, Debre zeit, Ethiopia.
59. Wollega, D., 1997. Survey of ticks and tick born disease in eight domestic animals in and round Deberzeit. Addis Ababa University, Faculty of Veterinary Medicine Deberzeit. DVM Thesis, Eastern Showa, pp: 1-79.
60. Gebre, S., G.P. Kaaya, G. Feseha and G. Tilahun, 1998. Insect Science and its application. *Journal of Ethiopian Veterinary Association*, 18(2, 3): 59-66.