

Effect of Citalopram on Reducing Transportation Stress in Rainbow Trout (*Oncorhynchus mykiss*)

¹M.M. Esfandanikeysomi, ¹M. Sudagar and ²Nasirir Asl

¹Department of Fishery, University of Agricultural Sciences and Natural Resources, Gorgan, Iran

²Department of Pharmacology, School of Medicine, University of Medical Sciences, Qazvin, Iran

Abstract: It is often necessary to transfer live fish between aquaculture facilities to permit on-growing or fishing, handling and the physical disturbances associated with loading, transport and discharge could be regarded as the potential to cause distress and injury and then leading to possible long term health impairment. Fish in different ways keep its stability (homeostasis) after stress responds. Stress response is involved in many physiological changes. One of the main reactions to stress is increasing plasma cortisol level. Purpose of this study was to examine the effect of a selective serotonin reuptake inhibitor (citalopram) on plasma cortisol and handling stress of rainbow trout. Immature rainbow trout (*Oncorhynchus mykiss*) with an average of 50 ± 7 g weight set in three treatments, control, acute (5 mg/l for 48 hours) and chronic (5 μ g/l for 10 days). Plasma cortisol before transport in control group was 22.11 ± 5.33 (ng/ml), chronic dose was 15.99 ± 5.85 (ng/ml) and in the acute treatment was 18.81 ± 7.42 (ng/ml). Mean plasma cortisol after transportation in the control, acute and chronic treatment respectively was 286.01 ± 54.26 , 107.12 ± 25.53 and 239.89 ± 57.56 ng/ml. Based on the results, significant difference observed ($p < 0.05$) between mean plasma cortisol before and after transportation in chronic treatment. Also in acute dose, there was a significant difference in mean plasma cortisol before and after transportation ($p < 0.05$).

Key words: Citalopram • Stress • Transport Stress • Rainbow Trout

INTRODUCTION

During transport, significant changes in water quality may occur and these may have numerous adverse effects on the vital physiological processes of fishes [1]. Continuously Fish is exposing to management practices such as handling, transportation or confinement which can elicit stress responses in intensive rearing facilities [2]. It was shown that fishes respond in different ways to maintain homeostasis after stress [3]. Many physiological changes are involved in such a stress response including hematology [4] osmoregulation [5], hormone release and energetic metabolism [6, 7].

Increasing plasma levels of cortisol as one of the most accepted primary response to stress has been previously reported [7, 8, 9]. Air exposure produced a 50-fold increase in plasma cortisol concentrations in gilthead sea bream (*Sparus aurata*) [10]. Two days of confinement procedures increased three-fold plasma

cortisol in gilthead sea bream, *S. aurata* [11]. In another study turbot, (*Scophthalmus maximus*), net confinement procedures provoked a 10-fold increase in plasma cortisol [12]. The stress response is highly comparable in all vertebrates and the secretion of both the glucocorticoid hormone corticosterone and cortisol is the main component [13]. In fish, the secretion of cortisol is mainly controlled by the hypothalamus-pituitary-interrenal (HPI) axis [14].

Corticosteroids affect behavior through genomic (slow) and non-genomic (fast) mechanisms in the central nervous system [15, 16]. In mammals the immediate increase in corticosteroids leads to escalated aggression propagated by a fast feed forward mechanism [17]. Moreover, regulation of glucocorticoid levels is influenced by behavior [18]. Sloman *et al.* [19] observed that the basal levels of circulating cortisol in rainbow trout (*Oncorhynchus mykiss*) prior to social interaction were associated with the outcome of fights for social

Corresponding Author: M.M. Esfandanikeysomi, Department of Fishery Sciences Gorgan University of Agricultural Sciences and Natural Resources, Beheshti Ave., Gorgan 49138-15739, Iran. Tel: +98(171) 245965, Ext: 464, Mob: +98-911-170-8928, Fax: +98(171) 2245884.

dominance. Schjolden *et al.* [18] perceived that there is a relevance between aggressive behavior and cortisol content. Adding anesthetic and hypnotic drugs to the transport tank water may be significantly useful to reduce the physiological stress associated with handling [20]. Drugs are primarily used to slow down metabolism, thus this will lead to reducing oxygen consumption and ammonia and carbon dioxide production. Furthermore they will mitigate the stress response caused by excitement and handling and reduce swimming [21, 22]. Serotonin is an important neurotransmitter in the regulation of social interactions in many animals. Correlative studies in numerous vertebrate species, including fish, indicated that, aggressive males have lower relative serotonergic activity than less aggressive males [23]. In the last decade, it was established that SSRIs such as fluoxetine and sertraline could be effective tools for chronic elevating serotonin activity. In recent studies the mentioned drugs are used to study the role of serotonin in aggressive behavior [24, 25]. The relationship between the brain serotonergic systems, social status and aggression has been established through a large number of correlative and experimental studies [26-28]. Thus, the aim of this study is to examine the effects of citalopram (a selective serotonin reuptake inhibitor) that caused reduced aggression in many species, on stress and plasma cortisol levels in rainbow trout.

MATERIALS AND METHODS

Immature rainbow trout (*Oncorhynchus mykiss*) with 50 ± 7 g bodyweight were obtained from a fish farm in Iran (Golestan, Zarringol Farm). Then they were maintained in small raceways in freshwater. Experiments were conducted from July to August of 2011.

The fish randomly selected and placed in three separate tanks (200 L) and adaptation was carried out for one week. Fish were fed once a day with a commercial diet until 24 h before the experiments. Physical and chemical water parameters were measured at the beginning and during the experiments. Water temperature was 16°C , pH was about 7.2 and water hardness was 145-155 (DH).

The fish was set in three doses of citalopram including acute (5 mg/l for 48 hours), chronic (5 μg /l for 10 days) and control treatment. We set 45 fish for each treatment, due to a wide individual difference in fish, each blood sample collected from 3 fishes and then mixed together. For inducing stress to fish, individuals were netted and carried (air exposure 10-15 s) to transport tanks. Fish were transported for 2 h with constant

oxygenation. After anesthesia with clove oil blood sample were immediately taken (with 1-ml heparinized syringes) from each fish without any anticoagulant material. The blood collection lasted less than 3 min. In order to avoid cortisol rise induced by the manipulation during sampling. The extracted blood was left to clot at 4°C and centrifuged at 6000 rpm for 5 min at room temperature. The collected serum was stored at 80°C for cortisol assay and then samples were transported to laboratory for plasma cortisol measurements. Data analysis performed with SPSS and the p-value = 0.05.

RESULTS

Mean plasma cortisol levels in control group before transport was 22.11 ± 5.33 ng/ml and after transport was 286.01 ± 54.26 ng/ml (Fig. 1). In the acute dose mean plasma cortisol was 18.81 ± 7.42 ng/ml and after transport was 107.12 ± 25.53 ng/ml (Fig. 2). Mean plasma cortisol in chronic treatment was 15.99 ± 5.85 ng/ml, before transport was 239.89 ± 57.56 ng/ml. Based on the results, the difference between mean **plasma cortisol in chronic dose before and after transport** was significant ($p < 0.05$). Also, there were significant differences ($p < 0.05$) between **the mean plasma cortisol before** and after transportation in acute treatment (Fig. 3).

DISCUSSION

The relationship between brain serotonergic systems, social status and aggression has been established through a large number of correlative and experimental researches or studies [23]; on the other hand, in other researches (Schjolden *et al.* [18]) the relevance between stress and aggression has been demonstrated. The results showed that artificial modification of the serotonin, which is an important neurotransmitter, can affect the stress behavior in rainbow trout. Because this drug is absorbed through the fish gills and its absorption is depending on temperature and fish metabolism, (e.g. in lower temperatures drug absorption is less) and because in the prolonged duration of exposure in chronic dose, the bioavailability of drug in the fish body will increase; led us to consider the chronic dose more effective than acute dose.

On the other hand, various drug onset times may be the reason for variant results in experiment. Notably, this study did not survey other effects of citalopram on various biological factors of (fish), therefore more research should be carried out in order to make perfect this study perfect.




Fig. 1: Mean plasma cortisol before moving




Fig. 2: Mean plasma cortisol after moving




Fig. 3: Comparison between all treatments

REFERENCES

- King, R.H., 2009. Fish transport in the aquaculture sector: An overview of the road transport of Atlantic salmon in Tasmania. *Veterinary Behavior*, 4: 163-168.
- Davis, C.R., M.S. Okihiro and D.E. Hinton, 2002. Effects of husbandry practices, gender and normal physiological variation on growth and reproduction of Japanese medaka, *Oryzias latipes*. *Aquatic Toxicology*, 60(3-4): 185-201.
- Wedemeyer, G.A., B. Barton and D.J. McLeay, 1990. Stress and Acclimation. In: C.B. Schreck and P.B. Moyle, (Eds.), *Methods for Fish Biology*. American Fisheries Society, Bethesda, Maryland, pp: 451-489. Chapter 14.
- Dethloff, G.M., D. Schlenk, S. Khan and H.C. Bailey, 1999. The effects of copper on blood and biochemical parameters of rainbow trout (*Oncorhynchus mykiss*). *Archives of Environmental Contamination and Toxicology*, 36(4): 415-423.
- McDonald, G. and L. Milligan, 1997. Ionic, Osmotic and Acid-Base Regulation in Stress. In: G.K. Iwama, A.D. Pickering, J.P. Sumpter and C.B. Schreck, (Eds.), *Fish Stress and Health in Aquaculture*. Cambridge University Press, Cambridge, U.K, pp: 119-144.
- Carragher, J.F. and C.M. Rees, 1994. Primary and secondary stress responses in golden perch, *Macquaria ambigua*. *Comparative Biochemistry and Physiology*, 107A: 49-56.
- Barton, B. and G. Iwama, 1991. Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of Fish Diseases*, 1: 3-26.
- Ortun, J., A. Esteban and J. Meseguer, 2002. Lack of effect of combining different stressors on innate responses of sea bream (*Sparus aurata* L.). *Veterinary Immunology and Immunopathology*, 84: 17-27.
- Barton, B., 2002. Stress in fishes: a diversity with particular reference in changes in circulating corticosteroids. *Integrative and Comparative Biology*, 42: 517-525.
- Arends, R.J., J.M. Mancera, J.L. Munoz and S.E. Wendelaar Bonga, 1999. The stress response of the gilthead seabream (*Sparus aurata* L.) to air exposure and confinement. *Endocrinol.*, 163: 149-157.
- Tort, L., E. Gomez, D. Montero and O. Sunyer, 1996. Serum hemolytic and agglutinating activity as indicators of fish immunocompetence: their suitability in stress and dietary studies. *Aquaculture International*, 4: 31-41.
- Waring, C.P., R.M. Stagg and M.G. Poxton, 1996. Physiological responses to handling in the turbot. *Fish Biology*, 48: 161-173.
- Mommsen, T.P., M.M. Vijayan and T.W. Moon, 1999. Cortisol in teleosts: dynamics, mechanisms of action and metabolic regulation. *Fish Biology Review*, 9(3): 211-68.
- Donaldson, E.M., 1981. The pituitary-interrenal axis as an indicator of stress in fish. In: A.D. Pickering, editor. *Stress in Fish*. London: Academic Press, pp: 11-47.
- Haller, J., J. Halasz, G.B. Makara and M.R. Kruk, 1998. Acute effects of glucocorticoids: behavioral and pharmacological perspectives. *Neuroscience and Biobehavioral Review*, 23(2): 337-44.
- Di Battista, J.D., H. Anisman, M. Whitehead and K.M. Gilmour, 2005. The effects of cortisol administration on social status and brain monoaminergic activity in rainbow trout (*Oncorhynchus mykiss*). *Exp. Biol.*, 208(14): 2707-18.

17. Kruk, M.R., J. Halasz, W. Meelis and J. Haller, 2004. Fast positive feedback between the adrenocortical stress response and a brain mechanism involved in aggressive behavior. *Behav. Neurosci.*, 118(5): 1062-1070.
18. Schjolden, A., D. Basic and S. Winberg, 2009. Aggression in rainbow trout is inhibited by both MR and GR antagonists. *Physiology and Behavior*, 98: 625-630.
19. Sloman, K., N.B.A. Metcalfe, A.C. Taylor and K.M. Gilmour, 2001. Plasma cortisol concentrations before and after social stress in rainbow trout and brown trout. *Physiology Biochemical Zoology*, 74(3): 383-9.
20. Finstad, B., M. Iversen and R. Sandodden, 2003. Stress-reducing methods for releases of Atlantic salmon (*Salmosalar*) smolts in Norway. *Aquaculture*, 222: 203-214.
21. Wedemeyer, G.A., 1997. Rearing conditions: effects on fish in intensive culture. In: G.K. Iwama, A.D. Pickering, J.P. Sumpter and C.B. Schreck, (Eds.), *Fish Stress and Health in Aquaculture Society. Experimental. Biology. Seminar Series*, University Press, Cambridge, 62: 35-71.
22. Ross, L.G. and B. Ross, 1999. *Anesthetic and sedative techniques for aquatic animals*, 2nd ed. Blackwell, London, UK, pp: 159.
23. Perreault, Heidi A.N., K. Semsar and J. Godwin, 2003. Fluoxetine treatment decreases territorial aggression in a coral reef fish. *Physiology and Behavior*, 79: 719-724.
24. Fuller, R.W., 1996. The influence of fluoxetine on aggressive behavior. *Neuro Psychopharmacology*, 14: 77-81.
25. Larson, E.T. and C.H. Summers, 2001. Serotonin reverses dominant social status. *Behavioral Brain Research*, 121: 95-102.
26. Blanchard, D.C., P. Cholvanich, R.J. Blanchard, D.W. Clow, R.P. Hammer and J. Rowlett, 1991. Serotonin, but not dopamine, metabolites are increased in selected brain-regions of subordinate male rats in a colonyenvironment. *Brain Research*, 568: 61-6.
27. Raleigh, M.J., M.T. McGuire, G.L. Brammer, D.B. Pollack and A. Yuwiler, 1991. Serotonergic mechanisms promote dominance acquisition in adultmale vervet monkeys. *Brain Res.*, 559: 181-90.
28. Winberg, S., C.G. Carter, J.D. McCarthy, Z.Y. He, G.E. Nilsson and D.F. Houlihan, 1993. Feeding rank and brain serotonergic activity in rainbow trout *Oncorhynchus mykiss*. *Experimental Biology*, 179: 197-211.