

## Flag Leaf Role in N Accumulation and Remobilization as Affected by Nitrogen in a Bread and Durum Wheat Cultivar

<sup>1</sup>A. Bahrani and <sup>2</sup>M. HaghJoo

<sup>1</sup>Department of Agriculture, Islamic Azad University, Ramhormoz branch, Khuzestan, Iran

<sup>2</sup>Department of Agriculture, Former Graduated Student in M.Sc,  
Islamic Azad University, Arsanjan branch, Fars, Iran

**Abstract:** Understanding the physiological basis of absorption and transportation of nitrogen by plants has specific importance. In this experiment, a bread and durum wheat cultivar were treated with different rates and timing of nitrogen application, using split factorial using a randomized complete block design with three replications at Khuzestan region during 2007-2008. Main plots consisted of two wheat cultivars (Cahmran, *Triticum aestivum* L. and Yavaros, *Triticum durum* L.) and sub plots were the nitrogen rates (40, 80 and 160 kg ha<sup>-1</sup>) and times of nitrogen application (T1= all N fertilizer at planting, T2= 1/2 at planting + 1/2 during booting stage and T3= 1/3 at planting + 1/3 during booting stage + 1/3 at heading stage). The results showed that there were significant differences between cultivars in flag leaf nitrogen content at maturity stage, N remobilization and its efficiency from flag leaf to the grains and also grain protein percentage. Durum wheat was more efficient in nitrogen remobilization and therefore, had higher grain protein percentage. Increasing rates and timing of nitrogen application had significant effect on most of the traits. There were significant interactions between cultivars, rates and times of N application, indicating that durum wheat was more efficient in N remobilization from flag leaf to the grain. In general, it appeared that N remobilization efficiency was the main factor affecting the grain protein percentage under the conditions of low N absorption in this experiment.

**Key words:** Wheat • Nitrogen remobilization efficiency • Grain protein

### INTRODUCTION

Nitrogen is the most expensive fertilizer nutrient used to raise crop plants [1,2]. A sizable portion of applied N is lost by leaching and denitrification [3,4]. The increased demand for N fertilizers has also raised farm input costs. Therefore, plant breeders need to develop cultivars that can absorb N more efficiently from the soil and partition most of the absorbed N into the grain. Such cultivars would minimize loss of N from the soil and make more economic use of the absorbed N.

Nitrogen transportation in plants, especially in growth and development stages, where N absorption from the soil is limited, is one of the questionable phenomena in fundamental crop physiology. For example, several studies have indicated that grain N in wheat primarily originates as a result of translocation from vegetative parts after anthesis [5-7]. This translocation depends on environment conditions [8], is under genetic control [9-11] and also can be affected by fertilizer N application [12].

Nitrogen remobilization efficiency from vegetative organs to grains was more significant in high protein cultivars [13]. It seems that, the greater N remobilization efficiency has been the main factor in increasing grain protein in high protein cultivars [14]. Austin *et al.* [15] observed significant genotypic differences for vegetative N concentration at anthesis, yet genotypic variation for total N content at anthesis was associated primarily with variation in dry matter. Several authors have reported significant variation for postanthesis N uptake [16, 15, 17] and in several studies N uptake during grain fill accounted for as much as 50 % of the grain N content at maturity [18-21]. Terman [21] and Perez *et al.* [22] reported that, grain protein content maybe improved by selection of genotypes which transport more nitrogen from vegetative organs to grain. Lotfali Ayeneh and Radmehr [23] found that, there were significant differences between nitrogen application rates and genotypes in nitrogen absorption at anthesis, amount of N remobilization and N absorption at maturity stage. Patric and Smith [24] demonstrated that

N fertilizer partitioning increase remobilization efficiency. Because the most part of nitrogen remobilization to grain raised from early nitrogen application treatment. Rostami and Jereai [19] showed that nitrogen concentration in above ground organs of high protein cultivars was significantly more than low protein cultivars. Austin *et al.* [15] in a study with 47 genotypes of winter wheat reported that, nearly 50 percent of nitrogen in spike absorbed from the soil after anthesis stage and the rest from N remobilization. Van sanford and Mackown [10] and Tindal *et al.* [25] stated that flag leaf at maturity stage contained lower portion of nitrogen in low-N environments than enough-N environments. This probably indicates that, flag leaf nitrogen content under nitrogen stress conditions was highly transported to the grain. Waldren and Flowerdy [26] and Wada *et al.* [27] showed that, about 80 percent of nitrogen at maturity stage absorbed at anthesis stage and two third of the nitrogen was transported from leaves to the grains.

Wheat grown under water deficit condition in the Mediterranean climatic region of Iran commonly fill their grain under increasing soil water deficits which restrict the uptake of N from the soil. Under these circumstances, most grain N in wheat may be derived from pre-anthesis stored N [28, 7].

Because of the importance of protein content and indirectly fertilizer N management to grain quality, this experiment was conducted to evaluate the effect of N fertilizer rate and time on the accumulation, remobilization and remobilization efficiency of nitrogen from flag leaf to the grain and also studying the relation between nitrogen concentration in flag leaf with grain protein.

**MATERIALS AND METHODS**

This study was conducted at a field experiment located at the Islamic Azad University of Ramhormoz, Khuzestan, Iran (31° 16' N, 49° 36' E and 150.5 m above the sea level) during 2007-2008. Total precipitation was 310.85 mm, which is shown in Fig. 1. The experiment was complete randomized block, split factorial design with three replications. Main plots consisted of two levels of cultivars (“Chamran” *Triticum aestivum L.* and “Yavarous” *Triticum durum L.*) and sub plots including nitrogen rates (40, 80 and 160 kg ha<sup>-1</sup>) and times (T1= all N fertilizer at planting, T2=1/2 at planting +1/2 during booting stage and T3= 1/3 at planting + 1/3 during booting stage + 1/3 at heading stage ), arranged as factorial. To determine the soil characteristics 15 samples from 30 cm depth were collected and analyzed for physical




Fig. 1: Monthly values of temperature and precipitation in the field experiment

Table 1: Soil physical and chemical characteristics of the experimental site

Variable	
Texture	Si-C
PH	8.96
Ecd S m <sup>-1</sup>	3.88
Organic matter (%)	0.55
N(%)	0.042
P mg Kg <sup>-1</sup>	5.4
K mg Kg <sup>-1</sup>	340
Fe	2.6
Zn	0.56
Mn	5.2
Cu	0.66

and chemical properties (Table 1). P and K fertilizer were applied according to recommendations of soil testing in forms of superphosphates and potassium sulfate, respectively. Plots were sown on 11 November 2007 with a cone seeder and were 8 m long and 1.5 m wide, with 6 rows 0.2 m apart. Plots were plowed and disked after winter wheat harvest in July. The plots were disked again before seeding in November. Apirus was applied in early April to the crop to control both broad and narrow leaved weeds. Twenty main stems that headed on same day were tagged for each treatment. As the tagged main stems of each cultivar reached anthesis, 10 plants in each plot were removed and flag leaves blade of main stems was cut. Tillers were discarded. At maturity, 10 additional tagged plants were removed and flag leaves blade of main stems was cut. Anthesis was scored when anthers in the central florets of 50% of the spikes in plant had dehiscid and maturity when almost all the spikes in plot showed complete loss of green color. Samples were dried to constant weight at 70°C, weighed and ground to pass a

0.4-mm screen. Grain samples were ground using the same screen. Total nitrogen concentration was determined by standard macro-Kjeldhal procedure. Grain protein concentration was calculated by multiplying grain N concentration by 5.7 [29].

The various parameters referring to N movement within the flag leaf of wheat plant that are discussed in this paper were evaluated as follows: [30].

Flag leaf nitrogen content at anthesis stage ( $\text{mg plant}^{-1}$ ).

Flag leaf nitrogen content at maturity stage ( $\text{mg plant}^{-1}$ ).

Nitrogen remobilization ( $\text{mg plant}^{-1}$ ) = N content at anthesis – N content at maturity.

Nitrogen remobilization efficiency (%) = (N remobilization/N content at anthesis)  $\times$  100.

Data were analyzed by analysis of variance [31]. When significant differences were found ( $P=0.05$ ) among means, Duncan's multiple range test (DMRT) were applied.

## RESULTS AND DISCUSSION

The results showed significant differences between two wheat cultivars in nitrogen remobilization, its efficiency and grain protein content (Table 2). Mean comparisons (Table 3) indicated that, there were no significant differences between cultivars in nitrogen content of flag leaf at anthesis stage. This probably meant that both cultivars absorbed the same amount of nitrogen from the soil. But, at maturity stage, durum wheat reserved less nitrogen content in flag leaf than bread cultivar, indicating more N remobilization in durum wheat than bread wheat. Consequently, grain protein percentage in durum wheat increased. Branlard *et al.* [32] and Cox *et al.* [14] found that differences among wheat cultivars in the amount of residual N in the straw resulted mainly from the differences in straw DM, since the differences in straw N concentrations were small. Rostami and Jereiae [19] reported that N remobilization was the reason of increasing grain protein percent. It seems that a genetic difference between two cultivars also is another factor in N remobilization from vegetative organs to the grains. Significant differences for N remobilization after anthesis were reported in a study conducted on 95 F<sub>2</sub> breed wheat lined derived from a single cross [33]. Grain protein in low protein cultivars depended on translocation of N already present in lower leaves at anthesis. In contrast, high protein cultivars required continued assimilation of N by

flag leaves after anthesis [34,13]. One possible avenue for increasing the grain nitrogen yield in wheat is through improving the efficiency of remobilization of nitrogen in the plant to the grain. Other studies have shown that the relationship between grain protein concentration and N translocation or N-translocation efficiency is not consistent [35, 36]. Conversely, Robert *et al.* [37], McNeal *et al.* [38] and Bhatia and Rabson [39] have reported that protein concentration in grain might be improved by selecting genotypes that translocate a higher percentage of N from vegetative organs to the grain. Positive correlations have been observed in wheat between grain protein concentration and nitrogen harvest index [40-43]. The descending ranking of sources of advantage of high protein wheat cultivars would seem to be: viability and continued assimilation of N by flag leaves after anthesis; efficiency of translocation of vegetative N present at anthesis; and amount of vegetative N present at anthesis [44, 9, 43, 21, 25].

Nitrogen application rates had significant effect on all traits (Table 2). By increasing nitrogen levels all traits increased, except N remobilization efficiency (Table 4). Increasing N remobilization due to N fertilization was also reported by Pritchard and Bhandari, [2] Spiertz and Ellen [45] and Cox *et al.* [14]. N content of flag leaf enhanced at anthesis and maturity stages by increasing of nitrogen. N remobilization from flag leaf to the grain increased by increasing N application rate, resulted in increasing grain protein percent. However, N remobilization efficiency decreased by increasing nitrogen application rate. Grain N content in wheat depends on uptake of soil nitrate prior to flowering, continued uptake of nitrate during grain fill and remobilization of stored vegetative N accumulated prior to flowering. However, under condition of high N fertility and available soil moisture, the contribution of post anthesis N uptake may increase substantially [5, 46]. Palta and Filery [4] reported that N remobilization efficiency decreased, when the amount of N increased. N absorption after flowering in Mediterranean climate conditions, similarly climate of Iran, is slow because of drought conditions, which is generally happens after anthesis stage. Other researchers found that N absorption reduced 25% during hot and drought summer [47-49]. But when the summer was moderate this amount was about 50% [20, 28, 50]. It seems that environmental differences lead to different patterns to N remobilization from flag leaves to the grains. In general, when plants exposed to low nitrogen in the soil the mechanism of N remobilization was more efficient.

Table 2: Analysis of variance of cultivars, rate and time of nitrogen application and interaction between them on nitrogen content of flag leaf at anthesis and maturity stage, N remobilization, N remobilization efficiency and grain protein percent

s.o.v	d.f	N content of flag leaf at anthesis (mg g <sup>-1</sup> )	N content of flag leaf at maturity (mg g <sup>-1</sup> )	N remobilization (mg g <sup>-1</sup> )	N remobilization efficiency (%)	Grain protein (%)
Replication	2	0.457 ns	0.14 ns	0.003 ns	0.574 ns	0.211 ns
Varieties	1	4.932**	14.97**	131.7**	1633.5**	32.3**
Error (a)	2	8.65	0.035	0.033	0.722	0.335
N Rates	2	69.73**	14.23**	20.4*	43.6**	22.7**
N Times	2	69.23**	11.35**	3.99**	3.019*	8.13**
Varieties & N Rates	2	33.22**	0.16**	3.20**	12.06*	0.65**
Varieties & N Times	2	15.01**	1.57**	3.90**	100.05**	5.72**
N Rates & N Times	4	11.70**	0.85**	10.81**	59.96**	1.39**
Varieties & N Rates & N Times	4	16.32**	4.43**	13.78**	74.94**	6.38**
Error (bc)	32	2.644	0.005	0.002	0.419	0.122

Ns (Non significant), \*(Significant at 5%) and \*\*, 1% (Significant at 1%) levels

Table 3: Mean nitrogen content of flag leaf at anthesis and maturity stage, N remobilization, N remobilization efficiency and grain protein percent in bread and durum wheat cultivar

Cultivars	N content of flag leaf at anthesis (mg g <sup>-1</sup> )	N content of flag leaf at maturity (mg g <sup>-1</sup> )	N remobilization (mg g <sup>-1</sup> )	N Remobilization Efficiency (%)	Grain protein (%)
Chamran	16.44 a	6.51 a	9.67 b	58.7 b	12.29 b
Yavarous	16.97 a	5.46 b	11.59 a	69.7 a	13.70 a

Means with similar letters in each column are not significantly different at 5 % level of probability. (Duncan)

Table 4: Effect of nitrogen application rates on mean nitrogen content of flag leaf at anthesis and maturity stage, N remobilization, N remobilization efficiency and grain protein percent in bread and durum wheat cultivar

N application Rates (kg ha <sup>-1</sup> )	N content of flag leaf at anthesis (mg g <sup>-1</sup> )	N content of flag leaf at maturity (mg g <sup>-1</sup> )	N remobilization (mg g <sup>-1</sup> )	N Remobilization Efficiency (%)	Grain protein (%)
s40	15.12 b	5.15 c	10.45 b	66 a	11.93 c
80	16.22 b	5.88 b	10.29 c	63.39 b	12.95 b
160	18.85 a	6.92 a	12.21 a	63.22 b	13.48 a

Means with similar letters in each column are not significantly different at 5 % level of probability. (Duncan)

Table 5: Effect of nitrogen application times on mean nitrogen content of flag leaf at anthesis and maturity stage, N remobilization, N remobilization efficiency and grain protein percent in bread and durum wheat cultivar

N application Time *	N content of flag leaf at anthesis (mg g <sup>-1</sup> )	N content of flag leaf at maturity (mg g <sup>-1</sup> )	N remobilization (mg g <sup>-1</sup> )	N Remobilization efficiency (%)	Grain protein (%)
T1	14.48 b	5.19 c	9.56 c	64.06 ab	11.97 c
T2	17.38 a	5.98 b	11.22 b	64.67 a	12.87 b
T3	18.22 a	6.78 a	12.16 a	63.89 b	13.63 a

Means with similar letters in each column are not significantly different at 5 % level of probability. (Duncan)

\* T1= All N fertilizer at planting, T2= 1/2 at planting + 1/2 during booting stage and T3= 1/3 at planting + 1/3 during booting stage + 1/3 at heading stage

N content of the flag leaf at anthesis and maturity stages increased by N split application (Table 2). Also N remobilization increased by N split application, resulted in higher grain protein percentage. However, N remobilization efficiency decreased by N split application (Table 5). It was estimated that two-thirds of the grain N

in wheat is derived from N assimilated before anthesis and one-third from assimilation during grain development [51, 52]. However, this may vary greatly depending on soil moisture and N availability during the grain-filling period. Vansanford and Mackown [10] reported that N remobilization efficiency in flag leaf was greater under

nitrogen deficiency condition. Radmehr *et al.* [53] also, reported that splitting of N fertilizer raised N remobilization efficiency. Knowles and Watkins [54] found that most of the N taken up by the wheat plants was translocated to the grain either directly or by mobilization from other plant parts. When N was probably abundantly available, no significant difference in remobilization efficiency was observed. One explanation for differences in remobilization efficiency is that during grain filling period, the plant retains an amount of N at anthesis that is essential for survival and various biological functions, while the remainder is available for remobilization. It appears that N retained depends on cultivars and prevailing growth conditions, although genetic variability in nitrogen remobilization has been reported [55-59].

Interaction between cultivars and nitrogen application rates, cultivars and N application times and rates and times of N application were significant in all traits (Table 2). It seems that, durum wheat was more efficient in low N application rates, probably due to genetic characteristic of durum wheat. N remobilization in durum wheat was higher than bread wheat by increasing of N application rate. Lotfaly ayeneh and Radmehr [23] and Le Gouis *et al.* [60] reported that there are significant differences between durum wheat genotypes and N application rates in N remobilization efficiency. Interaction between cultivars and N application time showed that by split N application durum wheat had more N remobilization efficiency than bread wheat. Indicated under N deficiency conditions in the soil, the mechanism of N remobilization was more efficient in durum wheat. The results also showed that, N remobilization increased by split N application in two wheat cultivars. As, durum wheat had better response to split N application. The highest N remobilization efficiency obtained in durum wheat with 80 kg N application at the first time of N application level.

### CONCLUSION

In summary, N content was always greater at anthesis than maturity for the Mediterranean climate conditions of this study. Wheat grown under Hot and dry weather condition in the Mediterranean climatic region of Iran commonly fills their grain under increasing soil water deficits which restrict the uptake of N from the soil. Under these circumstances, most grain N in wheat may be derived from pre-anthesis stored N.

Remobilization efficiency of plant nitrogen to grain differed between cultivars over the range of nitrogen levels used in this study. As durum wheat was more efficient than beard wheat in nitrogen remobilization

efficiency. There was a general trend for a decreased efficiency in flag leaf remobilization with increasing of soil nitrogen in beard and durum wheat cultivars. The minimum level of nitrogen obtained in flag leaf tissue of the durum wheat under the lowest nitrogen level at maturity in the present study enables speculation to be made on the potential for increasing nitrogen transport out of these tissues by breeding for increased remobilization efficiency.

In comparing the nitrogen remobilization efficiency values for this tissue obtained at low soil nitrogen level with the values at the higher nitrogen levels, a large potential appears for increasing remobilization efficiency. Harper *et al.* [61] reported that decrease in grain N of the high-protein cultivars when flag leaves were removed at anthesis indicated that the importance of flag leaves in mobilizing and translocation vegetative N to the developing grain.

### ACKNOWLEDGMENTS

The authors wish to express their appreciation to Islamic Azad University of Ramhormoz for their cooperation in conducting the field experiment.

### REFERENCES

1. Anonymous, 1985. FAO fertilizer yearbook. Food and Agricultural Organization of United Nations, Rome, Italy.
2. Pritchard, P.E. and D.G. Bhandari, 1996. The influence of nitrogen fertilizers on expression of functional proteins in wheat. HGA-project-Report, 121: 35.
3. Huber, D.M., H.L. Waren, D.W. Nelson and C.Y. Tsai, 1977. Nitrification inhibitors-new tools for food production. *Bio-Science*, 27: 523-529.
4. Palta, J.A. and I.R.P. Filery, 1995. Nitrogen application enhances remobilization and reduces losses of preanthesis N in wheat. *CAB Abstract*. 4(45): 4.
5. Barbottin, A., C. Lecompte, C. Bouchard and M.H. Jeuffroy, 2005. Nitrogen remobilization during grain filling in wheat: genotypic and environmental effects. *Crop Sci.*, 45: 1141-1150.
6. Heinrich, G.M., 1979. Accumulation and distribution of nitrogen in four winter wheat cultivars. M.S. Thesis. Michigan State Univ., East Lansing.
7. Simmons, S.R. and D.M. Moss, 1978. Nitrate reductase as a factor affecting nitrogen assimilation during the grain filling period in spring wheat. *Crop Sci.*, 18: 584-586.

8. Spiertz, J.H.J., 1977. The influence of temperature and light intensity on grain growth in relation to the carbohydrate and nitrogen economy of the wheat plant. *Neth. J. Agric. Sci.*, 25: 192-197.
9. Charmet, G., N. Robert and G. Branlard, 2005. Genetic analysis of dry matter and nitrogen accumulation and protein composition in wheat kernels. *Theoretical and Applied Genetics*, 111(3): 540-550.
10. Van Sanford, D.A. and C.T. Mackown, 1987. Cultivar differences in nitrogen remobilization during grain fill in soft red winter wheat. *Crop Sci.*, 27: 295-300.
11. Halloran, G.M., 1981. Cultivar differences in nitrogen translocation in wheat. *Aust. J. Agric. Res.*, 32: 534-544.
12. Perez, P., R. Martinez-Carrasco and L. Sanchez de la Puente, 1983. Uptake and distribution of nitrogen in wheat plants supplied with different amounts of nitrogen after stem elongation. *Ann. Appl. Biol.*, 102: 399-406.
13. Kade, M., A.J. Barneix and S. Olmos, 2005. Nitrogen uptake and remobilization in tetraploid 'Langdon' durum wheat and a recombinant substitution line with the high grain protein gene *Gpc-B1*. *Plant Breeding*, 124(4): 343-349.
14. Cox, M.C., C.O. Qualset and D.W. Rains, 1990. Genetic variation for nitrogen assimilation and translocation in wheat. III. Nitrogen translocation in relation to grain yield and protein. *Crop Sci.*, 26: 737-740.
15. Austin, R.B., J.A. Edrich, M.A. Ford and R.D. Blackwell, 1977. The fate of dry matter, carbohydrates and C14 lost from the leaves and stems of wheat during grain filling. *Ann. Bot. (London)*, 41: 1309-1321.
16. Fathi, G.H., 2005. Effect of drought and nitrogen on remobilization of dry matter and nitrogen in six wheat cultivars. *Iranian J. Agric. Sci.*, 36(5): 1093-1101.
17. Cox, C.M., C.O. Qualset and D.W. Rains, 1985a. Genetic variation for nitrogen assimilation and translocation in wheat. I. Dry matter and nitrogen accumulation. *Crop Sci.*, 25: 430-435.
18. Gooding, M.J., P.J. Gregory, K.E. Ford and R.E. Ruske, 2007. Recovery of nitrogen from different sources following applications to winter wheat after anthesis. *Field Crops Res.*, 100: 143-154.
19. Rostami, M.A. and H. Jiriaei, 1998. Nitrogen concentration profiles in wheat and its relationship to grain protein. *Iranian J. Agric. Sci.*, 29: 447-453.
20. Gregory, P.J., B. Marshal and P.V. Biscoe, 1981. Nutrient relations of winter wheat. 3. Nitrogen uptake, Photosynthesis of flag leaves and translocation of nitrogen to grain. *J. Agric. Sci.*, 96: 539-547.
21. Terman, G.L., 1979. Yield and protein content of wheat grain as affected by cultivars, N and environmental growth factors. *Agron. J.*, 71: 437-440.
22. Perez, C.M., G.B. Gagampang, B.V. Esmama, R.U. Monserrate and B.Q. Juliano, 1973. Protein metabolism in leaves and developing grains for rice differing in grain protein content. *Plant Physiol.*, 51: 537-542.
23. Lotfaly Ayeneh, G. and M. Radmehr, 2000. The diversity of durum genotypes in nitrogen remobilization in five levels of nitrogen application. *Proceeding of 6<sup>th</sup> Iranian congress of crop production and plant breeding*. Babol. Iran, pp: 325-326.
24. Patric, B. and D.L. Smith, 1993. Accumulation and redistribution of dry matter and nitrogen by spring barley. *Agron. J.*, 85: 1114-1121.
25. Tindal, T.A. C.S. Jeffrey and R.H. Brooks, 1995. Irrigated spring wheat responses to top dressed nitrogen as predict by flag leaf nitrogen concentration. *J. Prod. Agric.*, 8: 46-52.
26. Waldren, R.D. and A.D. Flowerday, 1979. Growth stage and distribution of dry matter, N, P and K in winter wheat. *Agron. J.*, 71: 391-397.
27. Wada, Y., K. Miura and K. Watanabe, 1993. Effect of source-to-sink ratio on carbohydrate production and senescence of flag leaves during the ripening period. *Japanese J. Crop Sci.*, 62: 578-553.
28. Kirda, C., M.R. Dericci and J.S. Schepers, 2001. Yield response and N- fertilizer recovery of rained wheat growing in the Mediterranean region. *Field Crops Res.*, 71: 113-122.
29. A.A.C.C., 1971. *Cereal laboratory methods*, 8<sup>th</sup> ed. Amer. Assoc. of Cereal Chem., St. Paul, Minn.
30. Papakosta, D.K. and A.A. Gagianas, 1991. Nitrogen and dry matter accumulation, remobilization and losses of Mediterranean wheat during grain filling. *Agron. J.*, 83: 564-870.
31. SAS Institute, Inc. 2004. *SAS/STAT 9.1 User's guide*. SAS Institute Inc., Cary, NC.
32. Branlard, G., M. Dardevet and R. Saccomano, 2001. Genetic diversity of wheat storage proteins and bread wheat quality. *Euphytica.*, 119(1/2): 59-67.

33. Cox, C.M., C.O. Qualset and D.W. Rains, 1985b. Genetic variation for nitrogen assimilation and translocation in wheat. II. Nitrogen assimilation in relation to grain yield and protein. *Crop Sci.*, 25: 435-440.
34. Asseng, S. and S.P. Milroy, 2006. Simulation of environmental and genetic effects on grain protein concentration in wheat. *European J. Agronomy*, 25(2): 119-128.
35. Mikesell, M.E. and G.M. Paulsen, 1971. Nitrogen translocation and the role of the individual leaves in protein accumulation in wheat. *Crop Sci.*, 11: 919-925.
36. Peterson, D.M., L.E. Schrader, D.A. Cataldo and D. Smith, 1975. Assimilation and remobilization of nitrogen and carbohydrate in oats, especially related to growth protein concentration. *Can. J. Plant Sci.*, 55: 19-28.
37. Robert, N., C. Hennequet and P. Bérard, 2001. Dry matter and nitrogen accumulation in wheat kernel: genetic variation in rate and duration of grain filling. *J. Genetics and Breeding*, 55(4): 297-305.
38. McNeal, F.H., M.A. Berg, C.F. MacGuire, V.R. Stewart and D.E. Baldrige, 1972. Grain and Nitrogen Relationships in Eight Spring Wheat Crosses. *Crop Sci.*, 12: 599-602.
39. Bhatia, C.R. and R. Rabson, 1976. Bioenergetics consideration in cereal breeding for protein improvement. *Science (Washington)*, 194: 1418-1421.
40. Loffer, C.M., I.L. Rouch and R.H. Bush, 1985. Grain and plant protein relationship in hard red spring wheat. *Crop Sci.*, 25: 521-525.
41. Paccaud, F.X., A. Fosati and Hong Sheng Cao, 1985. Breeding for yield and quality in winter wheat: Consequences for nitrogen uptake and partitioning efficiency. *Z. Pflanzenzuecht.*, 94: 89-100.
42. Desai, R.M. and C.R. Bhatia, 1976. Nitrogen uptake and nitrogen harvest index in durum wheat cultivars varying in their grain protein concentration. *Euphytica*, 27: 561-566.
43. Bhatia, T., 1976. Some effects of nitrogen fertilizer on winter wheat. *J. Sci. Food Agric.*, 27: 287-97.
44. Stoddard, F.L. and D.R. Marshall, 1990. Variability in grain protein in Australian hexaploid wheat. *Aust. J. Agri. Res.*, 41: 277-288.
45. Spiertz, J.H.J. and J. Ellen, 1978. Effects of nitrogen on canopy development and grain growth of winter wheat in relation to assimilation and utilization of assimilates and nutrients. *Neth. J. Agric. Sci.*, 26: 210-231.
46. Perez, P., R. Martinez-Carrasco and L. Sanchez de la Puente, 1983. Uptake and distribution of nitrogen in wheat plants supplied with different amounts of nitrogen after stem elongation. *Ann. Appl. Biol.*, 102: 399-406.
47. Despo, K. and A.A. Gagianas, 1991. Nitrogen and dry matter accumulation, remobilization and losses for Mediterranean wheat during grain filling. *Agron. J.*, 83: 864-870.
48. Ho`rtensteiner, S. and U. Feller, 2002. Nitrogen Metabolism and Remobilization During Senescence. *J. Experimental Botany*, 53: 927-937.
49. Kichey, T., B. Hirel, E. Heumez, F. Dubois and J. Le Gouis, 2007. In winter wheat (*Triticum aestivum* L.), post-anthesis nitrogen uptake and remobilization to the grain correlates with agronomic and nitrogen physiological markers. *Field Crops Res.*, 102: 22-32.
50. Jamieson, P.D. and M.A. Semenov, 2000. Modeling nitrogen uptake and redistribution in wheat. *FieldCrops Res.*, 68(1): 21-29.
51. Ehdai, B. and J.G. Waines, 2001. Sowing date and nitrogen rate effects on dry matter and nitrogen partitioning in bread and durum wheat. *Field Crops Res.*, 73: 47-61.
52. Paolov, A.N. and T.I. Kolesnik, 1974. Factors determining different levels of protein accumulation in the grain high-and low protein varieties of wheat. *Plant Physio.*, 22: 267-272.
53. Radmehr, M., G. Lotfaly Ayeneh and A. Kajbaf, 1997. The effects of planting date on growth and yield of wheat in Kozestan climate. I- Accumulation and redistribution of dry matter. *Seed Plant*, 13: 22-33.
54. Knowles, R. and J.E. Watkins, 1993. The assimilation and translocation of plant nutrients in wheat during growth. *J. Agric. Sci.*, 21: 612-637.
55. Mickelson, S., D. See and F.D. Meyer, 2003. Mapping of QTL associated with nitrogen storage and remobilization in barley (*Hordeum vulgare* L.) leaves. *J. Experimental Botany*, 54(383): 801-812.
56. Thompson, R.L., T. Leavitt and S. Mattias, 2000. Leaf nitrogen concentration of wheat subjected to elevated [CO<sub>2</sub>] and either water or N deficits. *Agr. Ecosyst. Environ.*, 79: 53-60.
57. Pethes, J., E. Kiss and B. Debreczeni, 1996. Effect of split top dressings of N on the grain N concentration of winter wheat cultivars. *Agrokemia-es- Talajtan*, 45(1-2): 77-94.

58. Ghoneim, A.M., 2008. Nitrogen Dynamics and Fertilizer Use Efficiency in Rice Using the Nitrogen-15 Isotope Techniques. *World Appl. Sci. J.*, 3(6): 869-874.
59. Mehdi, S.M. M. Sarfaraz and M. Ibrahim, 2007. Fertilizer requirement of wheat in recently reclaim soils. *World Appl. Sci. J.*, 2(6): 559-568.
60. Le Gouis, J., D. Béghin and E. Heumez, 2000. Genetic differences for nitrogen uptake and nitrogen utilization efficiencies in winter wheat. *European J. Agronomy*, 12(3/4): 163-173.
61. Harper, L.A., R.R. Sharpe, G.W. Langdale and J.E. Gidens, 1987. Nitrogen cycling in a wheat crop, soil and arial nitrogen transport. *Agron. J.*, 79: 965-973.